

Historic and Contemporary Mercury Exposure and Potential Risk to Yellow-Billed Loons (*Gavia adamsii*) Breeding in Alaska and Canada

Authors: Evers, David C., Schmutz, Joel A., Basu, Niladri, DeSorbo, Christopher R., Fair, Jeff, et al.

Source: *Waterbirds*, 37(sp1) : 147-159

Published By: The Waterbird Society

URL: <https://doi.org/10.1675/063.037.sp117>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Historic and Contemporary Mercury Exposure and Potential Risk to Yellow-billed Loons (*Gavia adamsii*) Breeding in Alaska and Canada

DAVID C. EVERS^{1,*}, JOEL A. SCHMUTZ², NILADRI BASU³, CHRISTOPHER R. DESORBO¹, JEFF FAIR⁴,
CARRIE E. GRAY¹, JAMES D. PARUK¹, MARIE PERKINS³, KEVIN REGAN¹, BRIAN D. UHER-KOCH²
AND KENNETH G. WRIGHT^{1,2}

¹Biodiversity Research Institute, 19 Flaggy Meadow Road, Gorham, ME, 04038, USA

²U.S. Geological Survey, Alaska Science Center, 4210 University Drive, Anchorage, AK, 99508, USA

³University of Michigan, School of Public Health, 1415 Washington Heights, Ann Arbor, MI, 48109, USA

⁴Fairwinds Wildlife Services, P.O. Box 2947, Palmer, AK, 99645, USA

*Corresponding author: E-mail: david.evers@briloon.org

Abstract.—The Yellow-billed Loon (*Gavia adamsii*) is one of the rarest breeding birds in North America. Because of the small population size and patchy distribution, any stressor to its population is of concern. To determine risks posed by environmental mercury (Hg) loads, we captured 115 Yellow-billed Loons between 2002 and 2012 in the North American Arctic and sampled their blood and/or feather tissues and collected nine eggs. Museum samples from Yellow-billed Loons also were analyzed to examine potential changes in Hg exposure over time. An extensive database of published Hg concentrations and associated adverse effects in Common Loons (*G. immer*) is highly informative and representative for Yellow-billed Loons. Blood Hg concentrations reflect dietary uptake of methylmercury (MeHg) from breeding areas and are generally considered near background levels if less than 1.0 µg/g wet weight (ww). Feather (grown at wintering sites) and egg Hg concentrations can represent a mix of breeding and wintering dietary uptake of MeHg. Based on Common Loon studies, significant risk of reduced reproductive success generally occurs when adult Hg concentrations exceed 2.0 µg/g ww in blood, 20.0 µg/g fresh weight (fw) in flight feathers and 1.0 µg/g ww in eggs. Contemporary mercury concentrations for 176 total samples (across all study sites for 115 Yellow-billed Loons) ranged from 0.08 to 1.45 µg/g ww in blood, 3.0 to 24.9 µg/g fw in feathers and 0.21 to 1.23 µg/g ww in eggs. Mercury concentrations in blood, feather and egg tissues indicate that some individual Yellow-billed Loons in breeding populations across North America are at risk of lowered productivity resulting from Hg exposure. Most Yellow-billed Loons breeding in Alaska overwinter in marine waters of eastern Asia. Although blood Hg concentrations from most breeding loons in Alaska are within background levels, some individuals exhibit elevated feather and egg Hg concentrations, which likely indicate the uptake of MeHg originating from eastern Asia. Feather Hg concentrations tended to be highest in individuals overwintering farthest west (closer to Asia). A retrospective analysis of museum specimens ($n = 25$) found a two-fold increase in Yellow-billed Loon feather Hg concentrations from the pre-1920s (as early as 1845) to the present. The projected increase in Hg deposition (approximately four-fold by 2050) along with the uncertainty of Hg being released through the thawing of permafrost and Arctic sea ice suggest that Hg body burdens in Yellow-billed Loons may increase. These findings indicate that Hg is a current and potentially increasing environmental stressor for the Yellow-billed Loon and possibly other Nearctic-Palaearctic migrant birds. Received 19 June 2013, accepted 4 July 2013.

Key words.—Alaska, Asia, Common Loon, *Gavia adamsii*, *Gavia immer*, Mercury, Yellow-billed Loon.

Waterbirds 37 (Special Publication 1): 147-159, 2014

The Yellow-billed Loon (*Gavia adamsii*) is one of the rarest breeding birds in North America (Earnst 2004) and the rarest of the world's five species of loons (Family Gavidae). Approximately 3,000 Yellow-billed Loon individuals occur in Alaska, the only state in the United States where they breed (Stehn *et al.* 2013). The majority of the Alaska breeding population (80%) is found on the Arctic Coastal Plain (ACP), with the remaining (20%) nesting in the northern half of the Seward Peninsula. In an average year, < 1,000 pairs nest on the

ACP, where their population is patchy and unevenly distributed (Earnst *et al.* 2005). Monitoring surveys, initiated in 1985, indicate that the population appears stable (Earnst *et al.* 2005; Stehn *et al.* 2013). Because of the small population size and patchy distribution, any habitat changes or anthropogenic stressors are cause for concern. Potential population threats include expansion of the oil industry into relatively high density breeding areas (90% of the Yellow-billed Loon breeding population in the ACP is within the National Petroleum

Reserve - Alaska) (Earnst 2004). Other population threats include climate change, subsistence hunting (Schmutz 2009), commercial fishnets (bycatch; Żydulis *et al.* 2009), and contaminants such as mercury (Hg) and other pollutants along nearshore marine ecosystems where loons forage, especially in eastern Asia (Agusa *et al.* 2007) where the majority of Alaska's Yellow-billed Loon population overwinters (North 1994; Fair 2002; Schmutz *et al.* 2014). As such, the Yellow-billed Loon is being considered for listing under the Endangered Species Act (U.S. Fish and Wildlife Service 2009). The potential anthropogenic threat to Yellow-billed Loons addressed in this study is the exposure to methylmercury (MeHg) available in ecosystems.

Recent evidence suggests that Hg exposure in piscivorous birds is increasing in the North American Arctic (Braune 2007; Rigét *et al.* 2011). The origin for the increase of environmental Hg loads is related to atmospheric Hg deposition from Asian emission sources (Sunderland *et al.* 2009; Blum *et al.* 2013), and possible changes in climate patterns that are resulting in thawing of permafrost and melting of Arctic sea ice with subsequent releases of bromine and increased Hg deposition (Brooks *et al.* 2006).

While specific adverse effects of Hg on the Yellow-billed Loon population in North America have not yet been measured, they are well established for the closely related Common Loon (*G. immer*). The two species are considered a super-species (American Ornithologists' Union 1998) and they share similar plumages, vocalizations, behaviors, and ecology (Sjölander and Ågren 1976; North 1994; Evers *et al.* 2010). Yellow-billed Loon weights range from 3.7 to 6.4 kg (this study), and Common Loon weights range from 3.5 to 7.6 kg (Gray *et al.* 2014). Because of these strong ecological and morphometric similarities between species, the robust literature on the exposure and effects of Hg on the Common Loon can be used to make relevant inferences about Hg exposure to Yellow-billed Loons (Evers *et al.* 1998, 2008b; Burgess and Meyer 2008; Schoch *et al.* 2014).

Specific Hg concentrations in blood, feather and egg tissues of Common Loons can now be directly related to levels of reduced fledging success in breeding populations in Maine and New Hampshire (Evers *et al.* 2008b), Wisconsin and New Brunswick (Burgess and Meyer 2008), and New York (Schoch *et al.* 2014). These studies found close agreement among tissue Hg concentrations and subsequent effects, with levels causing significant reproductive concern generally above 2.0 µg/g wet weight (ww) in the blood, 1.0 µg/g ww in the egg and 20.0 µg/g fresh weight (fw) in the feather. Hg effect concentrations based on these studies are believed to be physiologically relevant for use with the similar-sized and closely related Yellow-billed Loon as the pharmacokinetics and overall ecology are comparable.

We examined current and historic tissues from Yellow-billed Loons while on their breeding territories in Alaska to assess levels of exposure to Hg and to compare these values with effect concentrations in Common Loons. Our specific objectives were to: 1) document current patterns of Hg exposure to Yellow-billed Loons through examination of multiple tissue types that portray different time periods of exposure; 2) infer the magnitude of possible reproductive effects of Hg exposure on Yellow-billed Loons; and 3) examine evidence for change in Yellow-billed Loon exposure to Hg over the past 170 years. The contemporary and retrospective assessment conducted here is timely because environmental Hg loads have increased over the past century and further increases are expected over the next several decades (Sunderland *et al.* 2009).

METHODS

Study Area

The primary study area is in Alaska within the ACP, roughly 100 km southeast of Barrow, interior from the Beaufort Sea coast, and 60 km southwest of Teshekpuk Lake (Fig. 1). Here exists an extensive system of variable-sized fish-bearing lakes and several major drainage rivers. This study area is within the high-density region for breeding Yellow-billed Loons



Figure 1. Study sites for sample collection in Yellow-billed Loons, 2002 to 2012.

(Earnst *et al.* (2005). The landscape is composed of a continuous permafrost environment, with a shallow active layer under tundra plant communities (< 1 m) and somewhat deeper active layers (taliks) underneath lakes. Unlike boreal forest areas, such as those within interior Alaska, where discontinuities in permafrost allow deep subsurface flow and connectivity, the shallow active layers of the ACP and the impermeability of the permafrost trap labile water near or at the surface. Thus, despite the ACP receiving relatively minimal annual precipitation (15-25 cm/year), it is replete with many small and large lakes (> 100,000), as well as highly saturated soils in many plant communities (Walker *et al.* 2005).

Our secondary, less intensive study sites were southwest and southeast of the ACP (Fig. 1). We sampled Yellow-billed Loons on the northern part of the Seward Peninsula, Alaska, which is similar to the ACP but with a slightly deeper active layer (for further description, see Schmutz *et al.* 2014). In Canada, our study site was the Daring Lake region of the Northwest Territories (65° 50' N and 111° 38' W), approximately 300 km northeast of Yellowknife. This area, just north of tree line, consisted of multiple oligotrophic lakes with irregular shorelines and coves that harbor multiple breeding pairs of Yellow-billed Loons. The lakes are larger and

deeper than those in Alaska and surrounded by Arctic tundra, but with prevalent rock outcroppings.

Contemporary Capture and Field Tissue Sampling

We evaluated Hg exposure in Yellow-billed Loons using blood, feathers, and eggs from 2002 to 2012, but not all tissue types were collected at every site (Table 1) or in every year. To locate individuals for capture and sampling purposes, we conducted aerial surveys using fixed-wing and rotary-wing aircraft as well as ground surveys. We captured Yellow-billed Loons during the mid to late incubation period using two methods: 1) nest trapping; and 2) off-nest trapping. For nest trapping, we used an approximately 1-m diameter spring-loaded aluminum bow-net. To prevent accidental breakage, we replaced Yellow-billed Loon eggs with wooden dummy eggs during the capture process. We captured approximately 15% of the Yellow-billed Loons while off nest by luring them into mist nets with decoys and call playbacks. Once captured, we recorded body and bill measurements, attached bands to their legs (one or two color bands on each leg along with a U.S. Geological Survey aluminum band), and collected blood and feather samples following protocols of Evers *et al.* (1998, 2008b). We collected partially-incubated eggs in the ACP (one

Table 1. Mercury (Hg) concentrations in blood, feather, and eggs from 115 Yellow-billed Loons sampled ($n = 176$) on breeding territories in Alaska, USA, and Northwest Territories, Canada, 2002-2012.

Region	Blood Hg ($\mu\text{g/g ww}$)			Feather Hg ($\mu\text{g/g fw}$)			Egg Hg ($\mu\text{g/g ww}$)		
	n	Mean (SE)	Range	n	Mean (SE)	Range	n	Mean (SE)	Range
Arctic Coastal Plain, Alaska	58	0.38 (0.03)	0.08-1.45	77	8.18 (0.57)	3.01-24.92	9	0.49 (0.13)	0.21-1.23
Daring Lake, Northwest Territories	13	0.61 (0.05)	0.30-0.96	13	6.22 (0.36)	4.09-8.12	—	—	—
Seward Peninsula, Alaska	—	—	—	6	7.37 (1.31)	3.62-11.09	—	—	—

egg per nest), wrapped them in aluminum foil that was cleansed with acetone, and kept them cool until they could be frozen.

Field Tissue Analyses

We followed analytical protocols previously outlined for blood and feathers (Evers *et al.* 1998) and for eggs (Evers *et al.* 2003). Blood and feather samples for 2007-2012 were analyzed for total Hg concentrations at the Biodiversity Research Institute Wildlife Mercury Research Lab (Gorham, Maine). Samples included secondary feathers (5 cm tips) and whole blood. We placed samples into nickel boats that were then weighed and analyzed for total Hg concentration using a thermal decomposition technique with an automated direct Hg analyzer via the U.S. Environmental Protection Agency Method 7473 (U.S. Environmental Protection Agency 2007). Before and after every set of 30 samples, we included one sample each of two standard reference materials (Dorm-3 and Dolt-4), two methods blanks, and one sample blank. After every 20 samples, a duplicate was analyzed. Mean percent recoveries for total Hg of standard reference materials were within acceptable levels (U.S. Environmental Protection Agency Method 7473). Blood and egg samples from 2002-2003 were analyzed for total Hg at the Research Triangle Institute (Research Triangle Park, North Carolina) using standard and comparable analytical procedures with cold vapor atomic adsorption. We report Hg results in $\mu\text{g/g}$ for all tissues and reported as ww for blood, fw for feathers, and dry weight (dw) converted to ww for eggs.

Museum Tissue Analyses

To gain inference into historical Hg levels, we obtained secondary covert feathers from Yellow-billed Loon specimens at the Harvard University Museum of Comparative Zoology ($n = 19$) and the University of Michigan Museum of Zoology ($n = 6$). Museum specimens originated from the ACP, Alaska ($n = 17$), Northwest Territories ($n = 6$), and the Seward Peninsula, Alaska ($n = 2$) between 1845 and 1949. We analyzed these specimens at the University of Michigan for organic Hg since inorganic Hg has sometimes been used as a preservative in museum specimens and more than 90% of total Hg in the feather exists in an organic form (Head *et al.* 2011). Feathers were washed to remove any surface contamination and homogenized whole with a grinder using a stainless steel vial and ball pestle. We then digested the ground feathers and extracted organic Hg using the method described by Head *et al.* (2011) and Nam and Basu (2011). Every 10 samples included a standard reference material (Dolt-4, Dogfish Liver Certified Reference Material for Trace Metals) and a method blank.

Statistical Analysis

Spatial patterns of mercury concentrations. Blood and feather Hg values were log-transformed to normalize data and reduce heteroscedasticity. Normality was checked with the Shapiro-Wilk test and homogeneity of variance was checked with the Bartlett test. Using

the nonparametric Spearman's rank correlation test, correlations between normally distributed blood Hg and non-normally distributed feather Hg concentrations were determined for individuals where both sample types were collected. One of our analytical goals was to evaluate the effects of sex and body mass on Hg concentration. However, since males are consistently heavier than females, sex and body mass are confounded if one uses the raw data in analysis. Thus, we transformed body mass data into normalized z-scores by region and sex. We then statistically examined the effects of sampling year, sampling region, sex, and body mass on blood and feather Hg concentrations using a general linear model framework. We distinguished among competing models of suites of covariates by ranking the relative fit of each model with the Akaike Information Criterion adjusted for small sample size (AIC_c). The model with the lowest AIC_c and those having $\Delta AIC_c < 2$ had the most statistical support, those between 4 and 7 had considerably less support, and those > 10 had virtually no support (Burnham and Anderson 2002). Additional insight to the relative amount of statistical support for a given model was provided by each model's Akaike weight. Statistical analyses were performed in Microsoft EXCEL and JMP SAS (SAS Institute, Inc. 2008).

Male and female Yellow-billed Loons are monomorphic and were sexed using HINTZ and CHD primers, following the molecular genetic techniques outlined in Guzzetti *et al.* (2008). However, DNA blood samples for a subset of individuals ($n = 33$) were not collected. To include these samples where sex data were unavailable, we used a logistic regression function to predict the sex of Yellow-billed Loons based on morphometric measurements of individuals of known sex. Morphometric measurements included diagonal tarsus (mm), tarsus width (mm), tarsus breadth (mm), toe length (mm), culmen length (mm), culmen width (mm), culmen depth (mm), and body mass (g). Analyses were performed in Program R (R Development Core Team 2012). Diagonal tarsus ($P = 0.02$) and body mass ($P = 0.02$) were significant predictors of sex and so individuals were included in the Hg analysis if their combination of diagonal tarsus and mass measurements resulted in a 90% or higher probability of correct sex assignment. Specifically, individuals weighing less than 5,145 g had a 90% probability of being female, and individuals weighing more than 5,924 g had a 90% probability of being male (for Alaska individuals only). This function allowed the inclusion of an additional 21 Yellow-billed Loons in the blood and feather Hg analyses.

To evaluate the relationship between Hg in secondary flight feathers and wintering area, we used the most westerly location (degree of longitude) estimate (Douglas *et al.* 2012) from 48 Yellow-billed Loons marked with satellite transmitters (Schmutz *et al.* 2014). We used a quantile regression approach (Cade *et al.* 1999), PROC QUANTREG (SAS Institute Inc. 2008), which is appropriate for situations where limiting factors may exert their influence near the tails of the dependent variable's distribution and thus are not easily detected by ordinary least squares regression, which focuses on the mean response. We evaluated the relationship of wintering longitude to Hg concentration by examining estimates from the 50, 55, 60, 65, 70, 75, 80, 85, 90, and 95% quantiles (Fig. 2).




Figure 2. Feather mercury (Hg) ($\mu\text{g/g}$, fw) concentrations in Yellow-billed Loons compared longitudinally to overwintering territories (smaller or more westerly longitudes are along the Asian coast, while larger or more easterly longitudes are along the North American coast). Longitudes east of the international dateline were recast as 180° plus the number of degrees east of the dateline. Darkened prediction lines reflect slopes whose confidence intervals did not overlap zero.

Long-term patterns of feather mercury concentrations. To compare Hg concentrations in museum secondary coverts with secondary flight feathers collected in the field since 2007, it was necessary to determine the correlation in Hg between secondary coverts and secondary flight feathers. Log-transformed Hg concentrations found in secondary coverts and secondary flight feathers collected from the same individual were plotted in a linear regression to determine the Hg relationship between the two feather types. The regression calculated from this analysis was then used to create post hoc estimates of secondary flight feather Hg concentrations for museum specimens where only secondary coverts were available for Hg analysis. Based on available data, sample collection periods were divided into three groups: Pre-1920, 1930-1950, and Post-2000. Differences in Yellow-billed Loon secondary flight feather Hg concentrations among the three periods were examined with analysis of variance (ANOVA) and pairwise comparisons were tested with Tukey's honestly significant difference (HSD) test.

RESULTS

Contemporary Patterns of Mercury Exposure

We sampled 115 Yellow-billed Loons or 176 tissues (10 for blood only, 47 for feathers only, 9 for blood and egg only, and 49 for both blood and feathers) in the three study areas during the summers of 2002 to 2003 and 2007 to 2012 (Table 1). Most samples originated on the ACP.

Overall, blood Hg concentrations ranged from 0.08 to 1.45 µg/g ww, feather Hg concentrations ranged from 3.01 to 24.92 µg/g fw, and egg Hg concentrations ranged from 0.21 to 1.23 µg/g ww (Table 1). Blood and feather Hg were positively correlated in individuals where both sample types were available (Spearman rank correlation coefficient $\rho = 0.45$, $P = 0.001$, $n = 49$). Body mass ranged from 3,700-6,450 g among all sampling regions (Table 2). Male Yellow-billed Loons in the ACP were significantly larger than females, averaging 17% greater body mass ($t_{71} = 14.26$, $P < 0.001$), and males in the Daring Lake region averaged 22% greater body mass than females ($t_{12} = 6.41$, $P < 0.001$). Small sample size precluded comparison of male and female body masses in the Seward Peninsula region.

Spatial Patterns of Mercury Concentrations

We had no blood data from Seward Peninsula and only 1 year of blood data from Daring Lake in Canada. Thus, we conducted two analyses of variation in blood Hg: one examining year, sex, and body mass effects for loons sampled in the ACP over 6 years and one examining regional variation (ACP vs. Daring Lake) in one year (2010). For the time series of data from the ACP, blood Hg concentrations were highest in 2002 and 2003 ($\bar{x} = 0.69 \pm 0.15$ µg/g ww) and lowest in 2012 ($\bar{x} = 0.32 \pm 0.05$ µg/g ww). While blood Hg tended to be higher in males than females (Table 2), much of the variation in the data remained unexplained ($r^2 = 0.24$; Table 3). In 2010 (our lone sampling year for Daring Lake), blood Hg concentrations were greater at Daring Lake ($\bar{x} = 0.61 \pm 0.13$ µg/g ww) compared to the ACP ($\bar{x} = 0.38 \pm 0.03$ µg/g ww; $r^2 = 0.56$; Table 3). As with the previous analysis, the sex covariate fit substantially better than normalized body mass, suggesting that greater blood Hg in males is a function of ecological attributes other than body size or mass. Blood Hg concentrations were significantly higher in males ($\bar{x} = 0.56 \pm 0.08$ µg/g ww) compared to females ($\bar{x} = 0.41 \pm 0.05$ µg/g ww). The top supported model for feather Hg included no covariates. Thus, no differences in feather Hg concentrations

Table 2. Body mass (g) of Yellow-billed Loons sampled in Alaska, USA, and Northwest Territories, Canada, 2002-2012.

Region	Body Mass (g)													
	Female						Male						Unknown	
	n	Mean (SE)	Range	n	Mean (SE)	Range	n	Mean (SE)	Range	n	Mean (SE)	Range		
Arctic Coastal Plain, Alaska	49	4,945 (40)	4,032-5,450	24	5,893 (55)	5,300-6,450	29	5,369 (113)	4,325-6,360	—	—	—		
Daring Lake, Northwest Territories	8	4,150 (131)	3,700-4,900	6	5,183 (60)	4,900-5,300	—	—	—	—	—	—		
Seward Peninsula, Alaska	4	5,203 (214)	4,740-5,760	2	5,855 (205)	5,650-6,060	4	5,272 (390)	4,862-6,442	—	—	—		

Table 3. Model selection results examining the effect of sampling year, sampling region, sex, and body mass on blood and feather mercury (Hg) concentrations in breeding Yellow-billed Loons in Alaska, USA, and Northwest Territories, Canada from 2002 to 2012. Number of estimated parameters (K), differences between model Akaike Information Criterion adjusted for small samples size (ΔAIC_c) values <10, and AIC_c weights (w_i) are shown.

<i>Blood Hg across years on the ACP</i>			
Model	K	ΔAIC_c	w_i
Year (Temporal trend) + Sex	4	0	0.485
Year (Temporal trend)	3	1.153	0.273
Early vs. Late years + Sex	4	2.222	0.16
Year (Annual variation)	2	5.248	0.035
Sex	3	6.123	0.023
Null	4	6.769	0.016
Body mass	5	8.131	0.008

<i>Blood Hg between regions</i>			
Model	K	ΔAIC_c	w_i
Region + Sex	3	0	0.959
Region	2	6.23	0.041

<i>Feather Hg</i>			
Model	K	ΔAIC_c	w_i
Null	1	0	0.262
Body mass	2	0.892	0.168
Year (Temporal trend)	2	1.484	0.125
Sex	2	1.563	0.12
Body mass + Region	3	2.441	0.077
Body mass+ Sex	3	2.668	0.069
Sex + Region	3	2.836	0.063
Region	3	3.12	0.055
Body mass+ Sex + Region	4	4.151	0.033
Year (Annual variation)	6	4.449	0.028

were observed between sexes or among sampling regions (Table 3).

Estimates for the 70, 75, and 80% quantiles had confidence intervals that did not overlap zero and indicated that more westerly wintering loons (i.e., in eastern Asia) were exposed to more Hg than those wintering along the coast of North America (Fig. 2). Point estimates for the 85, 90, and 95% quantiles were even greater, but were imprecisely estimated due to sparser data.

Long-term Trends of Mercury Exposure

Mercury concentrations in secondary coverts and secondary feathers in Yellow-billed

Loons ($n = 15$) were strongly correlated ($r^2 = 0.97$, $F_{14} = 410.21$, $P < 0.001$). The regression calculated from this equation [LOG Secondary Hg = $-0.032575 + 1.0861684 \cdot \text{LOG Secondary Covert Hg}$] (SE of $m = 0.05$) was used to convert 25 historical secondary covert samples collected from museum specimens to secondary feather Hg concentrations. Feather Hg concentrations in Yellow-billed Loons varied significantly among sampling periods ($F_{128} = 20.10$, $P < 0.001$) (Fig. 3). Pairwise comparisons between sampling periods using Tukey's HSD test indicated that feather samples collected after 2000 ($\bar{x} = 8.01 \pm 0.45 \mu\text{g/g fw}$; $n = 104$) were significantly higher than those collected Pre-1920 ($\bar{x} = 3.81 \pm 0.40 \mu\text{g/g fw}$; $n = 19$; $q = 1.96$, $P < 0.001$). The other two pairwise comparisons, involving the middle time period ($\bar{x} = 4.97 \pm 0.76 \mu\text{g/g fw}$; $n = 6$) were not significant ($P > 0.05$).

DISCUSSION

Our results indicate current Hg concentrations for Yellow-billed Loons on their breeding range are generally below background levels (i.e., blood Hg $1.0 \mu\text{g/g ww}$) and are consistent with earlier work




Figure 3. Median feather mercury (Hg) ($\mu\text{g/g fw}$) concentrations in Yellow-billed Loons sampled during three time periods, Pre-1920 ($n = 19$), 1930-1950 ($n = 6$), and Post-2000 ($n = 104$, this study), across the three study areas in the Arctic Coastal Plain and Seward Peninsula, Alaska, USA, and east to the Daring Lake region, Northwest Territories, Canada. Each of these three study regions are represented in each of the three sampling periods.

conducted on breeding Common Loons in southern, boreal Alaska (Evers *et al.* 1998). Arctic environmental Hg loads primarily originate from atmospheric deposition. Historically, sources of deposition were natural (e.g., volcanic), whereas today the origin is primarily anthropogenic through current long-range transport (e.g., release from the burning of coal for industrial purposes) or re-emissions from ocean legacy sources (Corbitt *et al.* 2011; Mason *et al.* 2012). Once Hg is deposited in Arctic ecosystems, it is then made available through *in situ* methylation within lakes and their watersheds (St. Louis *et al.* 2005; Leitch *et al.* 2007). Methylmercury is transferred through the food web and biomagnifies as it moves into higher trophic level species and is well studied in temperate ecosystems (Driscoll *et al.* 2007) and increasingly in Arctic ecosystems (Kirk *et al.* 2012). Methylation and demethylation processes are dependent on many biogeochemical factors and can vary greatly among watersheds and even within watersheds and lakes. Such landscape variability creates significant challenges for determining spatial gradients of risk to Yellow-billed Loons. Mercury input from the surrounding watershed (Gantner *et al.* 2010b) and food web length appear to be important drivers for MeHg transfer to upper trophic organisms in Arctic lakes and subsequent variability (Ch  telat *et al.* 2008; Gantner *et al.* 2010a).

Ultimately, Yellow-billed Loon body burdens of MeHg are dictated by their prey, and as obligate piscivores, understanding preferred fish diet and determining Hg concentrations by fish species and size class will provide an ability to predict loon Hg concentrations based on models with Common Loons. For instance, among Common Loons Evers *et al.* (2008b) found that average prey fish Hg concentrations of 0.16 $\mu\text{g/g}$ ww (total Hg analyzed as whole body) relate to a 40% decline in the number of fledged chicks per territorial pair. Similarly, Depew *et al.* (2012) determined that 0.18 $\mu\text{g/g}$ ww of Hg in fish (whole body analyses) was responsible for a 50%

decline in overall productivity in Common Loons. Arctic lakes are known to contain fish of relevant prey size (< 25 cm; Barr 1996; Haynes *et al.* 2013) and species (e.g., cisco [*Coregonus* spp.]) that approach or exceed 0.16 $\mu\text{g/g}$ ww (Power *et al.* 2002), especially for land-locked Arctic char (*Salvelinus alpinus*) (Swanson *et al.* 2011).

Preliminary findings indicate that the majority of Yellow-billed Loons breeding on the ACP winter in eastern Asia (especially offshore of northern Japan), while half of the breeding population on the Seward Peninsula winter in eastern Asia and half along the Aleutian Islands, Alaska (J. A. Schmutz, pers. commun.). The majority (60%) of Yellow-billed Loons from the Daring Lake study site overwintered in Hecate Strait, British Columbia, and the remaining individuals overwintered as far west as the south-central shoreline of the Alaska Peninsula (Schmutz *et al.* 2014). These tracking efforts indicate that individuals from distinct breeding populations do not necessarily winter together, but the majority of the Yellow-billed Loons sampled within this study, principally those from the ACP, overwinter in eastern Asia. Because the Yellow-billed Loon undergoes a full remigial molt during winter (likely between December and February), the sampling of secondary feathers from individuals on their breeding territories allows a determination of dietary uptake of MeHg in marine waters of eastern Asia.

Past studies demonstrate that elevated body burdens of contaminants (e.g., polychlorinated biphenyls) in avian piscivores are accumulated while overwintering in eastern Asia (Kunisue *et al.* 2002; Minh *et al.* 2002; Schmutz *et al.* 2009). Yellow-billed Loons follow this contaminant loading pattern. Based on winter movements of trans-mitted individuals, there is a significant relationship with higher feather Hg concentrations in individuals overwintering to the west along the Asian coast (Fig. 2). Locally elevated atmospheric deposition (Jaffe and Strode 2008; Pan *et al.* 2008) and watershed input of Hg to eastern Asia marine waters likely contribute to this Hg exposure pattern.

Overall, feather Hg concentrations of Yellow-billed Loons sampled from 2007 to 2012 provided the most compelling evidence of risk. Seven percent of the feather samples exceeded Hg concentrations of approximately 20.0 $\mu\text{g/g}$ fw, which relate to projected lowered reproductive success in Common Loons (Evers *et al.* 2008b). However, exposure to Hg in Common Loons is greatest during the breeding season, whereas for Yellow-billed Loons it is currently greatest during the winter. Therefore, it is uncertain how MeHg loading during the winter may contribute to lowered reproductive output of Yellow-billed Loons.

Some insight into the contribution of wintertime exposure on breeding success lies within the blood-egg relationship of both loon species. There is a strong and significant relationship between blood and eggs for Common Loons ($r^2 = 0.79$, $n = 108$; Evers *et al.* 2003). Because adult female blood Hg concentrations of Common Loons strongly correlate with prey fish on their breeding territory (Burgess and Meyer 2008), it is well understood that egg Hg concentrations primarily reflect dietary uptake of MeHg from the breeding territory. However, there is a poor blood-egg relationship for Yellow-billed Loons (based on a limited number of eggs), which indicates that individual Hg body burdens accumulated in marine waters during winter and spring have a greater contribution to eggs than observed in Common Loons. Such observation and inference is consistent with the short interval between arriving at breeding areas and egg laying for Yellow-billed Loons (Schmutz *et al.* 2014) as compared to the long interval for Common Loons (Evers *et al.* 2010). Therefore, winter and spring exposure of Hg to Yellow-billed Loons may have a direct relationship to egg Hg concentrations, which may explain why breeding Yellow-billed Loons can have elevated egg Hg concentrations (i.e., $> 1.0 \mu\text{g/g}$ ww) while corresponding blood Hg concentrations are low (i.e., $< 1.0 \mu\text{g/g}$ ww). Larger sample sizes are necessary to determine if this observation is representative of the ACP and other areas with breeding Yellow-billed Loons.

Feather Hg concentrations measured in Yellow-billed Loons since 1845 have increased two-fold (Fig. 3). Our finding of this increase over this time period was expected and is consistent with sediment Hg concentrations over a similar time period from multiple sites in North America, where global atmospheric deposition was three to four times higher than pre-industrial times (Swain *et al.* 1992; Kamman and Engstrom 2002; Engstrom *et al.* 2007; Drevnick *et al.* 2012).

While contemporary blood Hg concentrations in Yellow-billed Loons generally remain under background levels defined for Common Loons for North America (Evers *et al.* 1998), egg and feather Hg concentrations for some individuals are elevated. Projections of water Hg concentrations in the North Pacific Ocean indicate a four-fold increase from current levels by 2050 (Sunderland *et al.* 2009; E. M. Sunderland, pers. commun.). Such increases could be even more pronounced within nearshore marine waters of the eastern Asian coast because of additional inputs from watersheds (Mason *et al.* 2012). Therefore, winter Hg body burdens could significantly increase over current levels and, because there may be a transfer of winter dietary uptake of MeHg to eggs, reproductive success could be adversely impacted. The future overall impact to Yellow-billed Loons from projected increasing environmental Hg emissions, deposition, and re-emissions (from legacy sources) will depend on changes within Arctic ecosystems related to permafrost thawing that are linked to changing climatic patterns (Schuur *et al.* 2009).

Loons are often-used as bioindicators of environmental quality because of their high trophic position and sensitivity to contaminants (Evers 2006). It is now well established that the Common Loon is experiencing adverse reproductive effects due to elevated environmental Hg loads (Burgess and Meyer 2008; Evers *et al.* 2003, 2008b, 2011; Schoch *et al.* 2014). Because of increases in global anthropogenic Hg emissions, spatial heterogeneity within a landscape, and changes in climatic patterns that may positively influ-

ence the remobilization of legacy environmental Hg loads and methylation rates, an assessment of the risk of Hg to Yellow-billed Loons within their breeding and wintering areas is timely. Biological Hg hotspots can form within areas of particular sensitivity to MeHg availability, even though the atmospheric input of Hg is relatively low (Driscoll *et al.* 2007; Evers *et al.* 2007). A better understanding for assessing risk to loons is now available, where effects concentrations in multiple tissues (i.e., blood, feathers and eggs) and prey can be confidently related to percent reduction in overall reproductive success (Depew *et al.* 2012) across broad landscapes.

Because of the uncertain relationship between Hg concentrations in air, water and sediment with high trophic level biota, the determination of spatial gradients and temporal trends on the magnitude of MeHg food web transfer requires suitable indicator organisms. The Yellow-billed Loon (and other loon species) are good high-trophic level indicators of Hg exposure across space (breeding and wintering areas) and time (short- and long-term). Blood provides a confident approach for examining short-term Hg exposure. Feathers collected from Yellow-billed Loons captured on their breeding territories indicate MeHg dietary uptake during the mid-winter molt are useful for determining Hg exposure along the eastern Asian near-shore waters during present and past time periods. Egg Hg for Yellow-billed Loons likely reflects a mixed Hg signal from wintering areas and breeding territories. The use of these three tissues provides a robust approach for monitoring Hg in both Arctic freshwater systems and marine ecosystems of eastern Asia. Such monitoring should be integrated into national programs of the United States (Mason *et al.* 2005; Evers *et al.* 2008a; Schmeltz *et al.* 2011) and Canada (Morrison 2011) and international efforts, such as the Global Mercury Observation System (Pirrone and Mason 2009) for the global Hg treaty (United Nations Environment Programme 2013), especially in light of increasing Hg emissions, watershed runoff of MeHg, and thawing of permafrost and pack ice.

ACKNOWLEDGMENTS

We thank the following individuals for field assistance: T. Arensberg, C. Bishop, B. Braden, H. Beutler, B. Courterier, D. Cummings, C. Eldermire, B. Estensen, T. Fondell, R. Gray, D. Heard, I. Isler, P. Lemons, T. Lemons, S. McCloskey, G. Meese, J. Morse, D. Mulcahy, D. Nigro, K. Overduijn, S. Reitsma, T. Ronningen, T. Shoemaker, M. Sorum, S. Stortz, H. Uher-Koch, C. VanStratt, and S. Wright. Feathers for retrospective analyses of Hg were provided by Jeremiah Trimble of the Harvard Museum of Comparative Zoology and Janet Hinshaw of the University of Michigan's Museum of Zoology. We also thank the National Fish and Wildlife Foundation for funding (2009-2012) and the U.S. Fish and Wildlife Service for aerial survey support. The Bureau of Land Management (D. Nigro) and National Park Service (B. Shults) provided in-kind aerial and logistical support. This work is part of the U.S. Geological Survey's Changing Arctic Ecosystem Initiative and is supported by funding from the Wildlife Program of the U.S. Geological Survey Ecosystem Mission Area. Permits were issued by relevant Federal and State agencies for Biodiversity Research Institute and the U.S. Geological Survey. We captured, banded, and sampled field tissues in accordance with Animal Care and Use protocols approved by the U.S. Geological Survey Alaska Science Center.

LITERATURE CITED

- Agusa, T., T. Kunito, A. Sudaryanto, I. Monirith, S. Kantireklap, H. Iwata, A. Ismail, J. Sanguansind, M. Muchtare, T. S. Tanaf and S. Tanabe. 2007. Exposure assessment for trace elements from consumption of marine fish in Southeast Asia. *Environmental Pollution* 145: 766-777.
- American Ornithologists' Union. 1998. Check-list of North American birds, 7th ed. American Ornithologists' Union, Washington, D.C.
- Barf, J. F. 1996. Aspects of common loon (*Gavia immer*) feeding biology on its breeding ground. *Hydrobiologia* 321: 119-144.
- Blum, J. D., B. N. Popp, J. C. Drazen, C. A. Choy and M. W. Johnson. 2013. Methylmercury production below the mixed layer in the North Pacific Ocean. *Nature Geoscience* 6: 879-884.
- Braune, M. B. 2007. Temporal trends of organochlorines and mercury in seabird eggs from the Canadian Arctic, 1975-2003. *Environmental Pollution* 148: 599-613.
- Brooks, S. B., A. Saiz-Lopez, H. Skov, S. E. Lindberg, J. M. C. Plane and M. E. Goodsite. 2006. The mass balance of mercury in the springtime Arctic environment. *Geophysical Research Letters* 33: L13812.
- Burgess, N. M. and M. W. Meyer. 2008. Methylmercury exposure associated with reduced productivity in common loons. *Ecotoxicology* 17: 83-91.
- Burnham, K. P. and D. R. Anderson. 2002. Model selection and multimodel inference: a practical information-theoretic approach. Second ed. Springer-Verlag, New York, New York.

- Cade, B. S., J. W. Terrell and R. L. Schroeder. 1999. Estimating effects of limiting factors with regression quantiles. *Ecology* 80: 311-323.
- Ch  telat, J., M. Amyot, L. Cloutier and A. Poulain. 2008. Metamorphosis in chironomids, more than mercury supply, controls methylmercury transfer to fish in High Arctic lakes. *Environmental Science and Technology* 42: 9110-9115.
- Corbitt, E. S., D. J. Jacob, C. D. Holmes, D. G. Streets and E. M. Sunderland. 2011. Global source-receptor relationships for mercury deposition under present-day and 2050 emissions scenarios. *Environmental Science and Technology* 45: 10477-10484.
- Depew D. C., N. Basu, N. M. Burgess, L. M. Campbell, D. C. Evers, K. A. Grasman, K. P. Kenow, M. W. Meyer, A. M. Scheuhammer and K. Williams. 2012. Derivation of screening benchmarks for the common loon (*Gavia immer*) for dietary methylmercury (MeHg): justification and rationale for use in ecological risk assessment. *Environmental Toxicology and Chemistry* 31: 2399-2407.
- Douglas, D. C., R. Weinzierl, S. C. Davidson, R. Kays, M. Wikelski and G. Bohrer. 2012. Moderating Argos location errors in animal tracking data. *Methods in Ecology and Evolution* 3: 999-1007.
- Drevnick, P. E., D. R. Engstrom, C. T. Driscoll, E. B. Swain, S. J. Balogh, N. C. Kamman and R. Rossmann. 2012. Spatial and temporal patterns of mercury accumulation in lacustrine sediments across the Laurentian Great Lakes region. *Environmental Pollution* 161: 252-260.
- Driscoll, C. T., Y. J. Han, C. Y. Chen, D. C. Evers, K. F. Lambert, T. M. Holsen, N. C. Kamman and R. Munson. 2007. Mercury contamination in remote forest and aquatic ecosystems in the northeastern U.S.: sources, transformations and management options. *Bioscience* 57: 17-28.
- Earnst, S. L. 2004. Status assessment and conservation plan for the Yellow-billed Loon (*Gavia adamsii*). Scientific Investigations Report 2004-5258, U.S. Department of the Interior, Geological Survey, Reston, Virginia.
- Earnst, S. L., R. A. Stehn, R. M. Platte, W. W. Larned and E. J. Mallek. 2005. Population size and trend of Yellow-billed Loons in northern Alaska. *Condor* 107: 289-304.
- Engstrom, D. R., S. J. Balogh and E. B. Swain. 2007. History of mercury inputs to Minnesota lakes: influences of watershed disturbance and localized atmospheric deposition. *Limnology and Oceanography* 52: 2467-2483.
- Evers, D. C. 2006. Loons as biosentinels of aquatic integrity. *Environmental Bioindicators* 1: 18-21.
- Evers, D. C., J. D. Paruk, J. W. McIntyre and J. F. Barr. 2010. Common Loon (*Gavia immer*). No. 313 in *The Birds of North America Online* (A. Poole, Ed.). Cornell Lab of Ornithology, Ithaca, New York. <http://bna.birds.cornell.edu/bna/species/313>, accessed 17 November 2013.
- Evers, D. C., K. M. Taylor, A. Major, R. J. Taylor, R. H. Poppenga and A. M. Scheuhammer. 2003. Common Loon eggs as indicators of methylmercury availability in North America. *Ecotoxicology* 12: 69-81.
- Evers, D. C., J. D. Kaplan, M. W. Meyer, P. S. Reaman, A. Major, N. Burgess and W. E. Braselton. 1998. Bioavailability of environmental mercury measured in Common Loon feathers and blood across North America. *Environmental Toxicology and Chemistry* 17: 173-183.
- Evers, D. C., Y. J. Han, C. T. Driscoll, N. C. Kamman, M. W. Goodale, K. F. Lambert, T. M. Holsen, C. Y. Chen, T. A. Clair and T. Butler. 2007. Identification and evaluation of biological hotspots of mercury in the northeastern U.S. and eastern Canada. *Bioscience* 57: 29-43.
- Evers, D. C., K. A. Williams, M. W. Meyer, A. M. Scheuhammer, N. Schoch, A. T. Gilbert, L. Siegel, R. J. Taylor, R. Poppenga and C. R. Perkins. 2011. Spatial gradients of methylmercury for breeding common loons in the Laurentian Great Lakes region. *Ecotoxicology* 20: 1609-1625.
- Evers, D. C., R. P. Mason, N. C. Kamman, C. Y. Chen, A. L. Bogomolni, D. H. Taylor, C. R. Hammerschmidt, S. H. Jones, N. M. Burgess, K. Munney and K. C. Parsons. 2008a. An integrated mercury monitoring program for temperate estuarine and marine ecosystems on the North American Atlantic Coast. *Ecology* 89: 426-441.
- Evers, D. C., L. Savoy, C. R. DeSorbo, D. Yates, W. Hanson, K. M. Taylor, L. Siegel, J. H. Cooley, M. Bank, A. Major, K. Munney, H. S. Vogel, N. Schoch, M. Pokras, W. Goodale and J. Fair. 2008b. Adverse effects from environmental mercury loads on breeding common loons. *Ecotoxicology* 17: 69-81.
- Fair, J. 2002. Status and significance of Yellow-billed Loon (*Gavia adamsii*) populations in Alaska. Unpublished report, Wilderness Society and Trustees for Alaska, Anchorage, Alaska.
- Gantner, N., M. Power, D. Iqaluk, M. Meili, H. Borg, M. Sundbom, K. R. Solomon, G. Lawson and D. C. Muir. 2010a. Mercury concentrations in landlocked Arctic char (*Salvelinus alpinus*) from the Canadian Arctic. Part I: insights from trophic relationships in 18 lakes. *Environmental Toxicology and Chemistry* 29: 621-632.
- Gantner, N., D. C. Murie, M. Power, D. Iqaluk, J. D. Reist, J. A. Babaluk, M. Meili, H. Borg, J. Hammar, W. Michaud, B. Dempson and K. R. Solomon. 2010b. Mercury concentrations in landlocked Arctic char (*Salvelinus alpinus*) from the Canadian Arctic. Part II: influence of lake biotic and abiotic characteristics on geographic trends in 27 populations. *Environmental Toxicology and Chemistry* 29: 633-643.
- Gray, C. E., J. D. Paruk, C. R. DeSorbo, L. J. Savoy, D. E. Yates, M. D. Chickering, R. B. Gray, K. M. Taylor, D. Long, IV, N. Schoch, W. Hanson, J. Cooley and D. C. Evers. 2014. Body mass in Common Loons (*Gavia immer*) strongly associated with migration distance. *Waterbirds (Special Publication 1)* 37: 64-75.
- Guzzetti, B. M., S. L. Talbot, D. F. Tessler, V. A. Gill and E. C. Murphy. 2008. Secrets in the eyes of Black

- Oystercatchers: a new sexing technique. *Journal of Field Ornithology* 79: 215-223.
- Haynes, T. B., A. E. Rosenberger, M. S. Lindberg, M. Whitman and J. A. Schmutz. 2013. Method- and species-specific detection probabilities of fish occupancy in Arctic lakes: implications for design and management. *Canadian Journal of Fisheries and Aquatic Sciences* 70: 1055-1062.
- Head, J. A., A. DeBofsky, J. Hinshaw and N. Basu. 2011. Retrospective analysis of mercury content in feathers of birds collected from the state of Michigan (1895-2007). *Ecotoxicology* 20: 1636-1643.
- Jaffe, D. and S. Strode. 2008. Sources, fate and transport of atmospheric mercury from Asia. *Environmental Chemistry* 5: 121-126.
- Kamman, N. C. and D. R. Engstrom. 2002. Historical and present fluxes of mercury to Vermont and New Hampshire lakes inferred from ^{210}Pb dated sediment cores. *Atmospheric Environment* 36: 1599-1609.
- Kirk, J. L., I. Lehnerr, M. Andersson, B. M. Braune, L. Chan, A. P. Dastoor, D. Durnford, A. L. Gleason, L. L. Loseto, A. Steffen and V. L. St. Louis. 2012. Mercury in Arctic marine ecosystems: sources, pathways, and exposure. *Environmental Research* 119: 64-87.
- Kunisue, T., T. B. Minh, K. Fukuda, M. Watanabe, S. Tanabe and A. M. Titenko. 2002. Seasonal variation of persistent organochlorine accumulation in birds from Lake Baikal, Russia and the role of the South Asian region as a source of pollution for wintering migrants. *Environmental Science and Technology* 36: 1396-1404.
- Leitch, D. R., J. Carrie, D. Lean, R. W. Macdonald, G. A. Stern and F. Wang. 2007. The delivery of mercury to the Beaufort Sea of the Arctic Ocean by the Mackenzie River. *Science of the Total Environment* 373: 178-195.
- Mason, R. P., A. L. Choi, W. F. Fitzgerald, C. R. Hammer-schmidt, C. H. Lamborg, A. L. Soerensen and E. M. Sunderland. 2012. Mercury biogeochemical cycling in the ocean and policy implications. *Environmental Research* 119: 101-117.
- Mason, R., M. Abbot, D. Bodaly, R. Bullock, C. Driscoll, D. Evers, S. Lindberg, M. Murray and E. Swain. 2005. Monitoring the environmental response to changes in mercury contamination from the atmosphere: a multi-media challenge. *Environmental Science and Technology* 39: 15A-22A.
- Minh, T. B., T. Kunisue, N. T. H. Yen, M. Watanabe, S. Tanabe, N. D. Hue and V. Qui. 2002. Persistent organochlorine residues and their bioaccumulation profiles in resident and migratory birds from North Vietnam. *Environmental Toxicology and Chemistry* 21: 2108-2118.
- Morrison, H. A. 2011. The Canadian clean air regulatory agenda mercury science program. *Ecotoxicology* 20: 1512-1519.
- Nam, D. H. and N. Basu. 2011. Rapid methods to detect organic mercury and total selenium in biological samples. *Chemistry Central Journal* 5: 1-5.
- North, M. R. 1994. Yellow-billed Loon (*Gavia adamsii*). No. 121 in *The Birds of North America* (A. Poole and F. Gill, Eds.). Academy of Natural Sciences, Philadelphia, Pennsylvania; American Ornithologists' Union, Washington, D.C.
- Pan, L., G. R. Carmichael, B. Adhikary, Y. Tang, D. Streets, J.-H. Woo, H. R. Friedl and L. F. Radke. 2008. A regional analysis of the fate and transport of mercury in East Asia and an assessment of major uncertainties. *Atmospheric Environment* 42: 1144-1159.
- Pirrone, N. and R. Mason. 2009. Mercury fate and transport in the global atmosphere: emissions, measurements and models. Springer, New York, New York.
- Power, M., G. M. Klein, K. R. R. A. Guiguer and M. K. H. Kwan. 2002. Mercury accumulation in the fish community of a sub-arctic lake in relation to trophic position and carbon sources. *Journal of Applied Ecology* 39: 819-830.
- R Development Core Team. 2012. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>, accessed 1 April 2013.
- Rigét, F., B. Braune, A. Bignert, S. Wilson, J. Aars, E. Born, M. Dam, R. Dietz, M. Evans, T. Evans, M. Gamberg, N. Gantner, N. Green, H. Gunnlaugsdóttir, K. Kannan, R. Letcher, D. Muir, P. Roach, C. Sonne, G. Stern and O. Wiig. 2011. Temporal trends of Hg in Arctic biota, an update. *Science of the Total Environment* 409: 3520-3526.
- SAS Institute, Inc. 2008. SAS/STAT v. 9.2 User's Guide. SAS Institute, Inc., Cary, North Carolina.
- Schmeltz, D., D. C. Evers, C. T. Driscoll, R. Artz, M. Cohen, D. Gay, R. Haeuber, D. P. Krabbenhoft, R. Mason, K. Morris and J. G. Wiener. 2011. MercNet: a national monitoring network to assess responses to changing mercury emissions in the United States. *Ecotoxicology* 20: 1713-1725.
- Schmutz, J. A., K. A. Trust and A. C. Matz. 2009. Red-throated Loons (*Gavia stellata*) breeding in Alaska, USA, are exposed to PCBs while on their Asian wintering grounds. *Environmental Pollution* 157: 2386-2393.
- Schmutz, J. A., K. G. Wright, C. R. DeSorbo, J. Fair, D. C. Evers, B. D. Uher-Koch and D. M. Mulcahy. 2014. Size and retention of breeding territories of Yellow-billed Loons (*Gavia adamsii*) in Alaska and Canada. *Waterbirds (Special Publication 1)* 37: 53-63.
- Schoch, N., M. J. Glennon, D. C. Evers, M. Duron, A. K. Jackson, C. T. Driscoll, J. W. Ozard and A. K. Sauer. 2014. The impact of mercury exposure on the Common Loon (*Gavia immer*) population in the Adirondack Park, New York, USA. *Waterbirds (Special Publication 1)* 37: 133-146.
- Schuur, E. A., J. G. Vogel, K. G. Crummer, H. Lee, J. O. Sickman and T. E. Osterkamp. 2009. The effect of permafrost thaw on old carbon release and net carbon exchange from tundra. *Nature* 459: 556-559.
- Sjölander, S. and G. Ågren. 1976. Reproductive behavior of the Yellow-billed Loon, *Gavia adamsii*. *Condor* 78: 454-463.

- St. Louis, V. L., M. J. Sharp, A. Steffen, A. May, J. Barker, J. L. Kirk, D. J. A. Kelly, S. E. Arnott, B. Keatley and J. P. Smol. 2005. Some sources and sinks of monomethyl and inorganic mercury on Ellesmere Island in the Canadian High Arctic. *Environmental Science and Technology* 39: 2686-2701.
- Stehn, R. A., W. W. Larned and R. M. Platte. 2013. Analysis of aerial survey indices monitoring waterbird populations of the Arctic Coastal Plain, Alaska, 1986-2012. Unpublished report, U.S. Department of the Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Anchorage, Alaska.
- Sunderland, E. M., D. P. Krabbenhoft, J. W. Moreau, S. A. Strode and W. M. Landing. 2009. Mercury sources, distribution, and bioavailability in the North Pacific Ocean: insights from data and models. *Global Biogeochemical Cycles* 23: GB2010.
- Swain E. B., D. R. Engstrom, M. E. Brigham, T. A. Henning and P. L. Brezonik. 1992. Increasing rates of atmospheric mercury deposition in Midcontinental North America. *Science* 257: 784-787.
- Swanson, H., N. Gantner, K. A. Kidd, D. C. G. Muir and J. D. Reist. 2011. Comparison of mercury concentrations in landlocked, resident, and sea-run fish (*Salvelinus* spp.) from Nunavut, Canada. *Environmental Toxicology and Chemistry* 30: 1459-1467.
- United Nations Environment Programme. 2013. Global mercury assessment 2013: sources, emissions, releases and environmental transport. United Nations Environment Programme, Chemicals Branch, Geneva, Switzerland.
- U.S. Environmental Protection Agency. 2007. Method 7473. <http://www.epa.gov/osw/hazard/test-methods/sw846/pdfs/7473.pdf>, accessed 2 December 2013.
- U.S. Fish and Wildlife Service. 2009. 12-month finding on a petition to list the Yellow-billed Loon as threatened or endangered. *Federal Register* 74: 12932-12968.
- Walker, D. A., M. K. Reynolds, F. J. A. Daniels, E. Einarsson, A. Elvebakk, W. A. Gould, A. E. Katenin, S. S. Kholod, D. J. Markon, E. S. Melnikov, N. G. Moskalenko, S. S. Talbot and B. A. Yurtsev. 2005. The circumpolar Arctic vegetation map. *Journal of Vegetation Science* 16: 267-282.
- Žydelis, R., J. Bellebaum, H. Österblom, M. Vetemaa, B. Schirmeister, A. Stipnice, M. Dagys, M. van Eerden and S. Garthe. 2009. Bycatch in gillnet fisheries – an overlooked threat to waterbird populations. *Biological Conservation* 142: 1269-1281.

