

A Case of Spontaneous Tool-Making by a Captive Capuchin Monkey

Authors: Soares Bortolini, Tiago, and Bicca-Marques, Júlio César

Source: Neotropical Primates, 14(2) : 74-76

Published By: Conservation International

URL: <https://doi.org/10.1896/044.014.0205>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

- Freese, C. H. e Oppenheimer, J. R. 1981. The capuchin monkeys, genus *Cebus*, p. 331–389.
- Em: A. F. Coimbra-Filho e R.A. Mittermeier (Ed.). Ecology and behavior of Neotropical primates. Rio de Janeiro, Academia Brasileira de Ciências, 496pp.
- Garber, P. A. 1989. The role of spatial memory in primate foraging patterns: *Saguinus mystax* and *Saguinus fuscicollis*. *Am. J. Primatol.* 19: 203–216.
- Hubener, F. e Laska, M. 1998. Assessing olfactory performance in an old world primate: *Macaca nemestrina*. *Physiol. Behav.* 64: 521–527.
- Hill, O. C. 1960. *Primate comparative anatomy and taxonomy. Vol. IV - Cebidae, Part A*. Edinburgh University Press.
- Laska, M. e Hudson, R. 1993. Discriminating parts from the whole: Determinants of odor mixture perception in squirrel monkeys, *Saimiri*. *Comp. Physiol.* 173: 249–256.
- Laska, M., Hudson, R. e Alicke, T. 1996. A study of long-term odor memory in squirrel monkeys (*Saimiri sciureus*). *J. Comp. Psychol.* 110: 125–130.
- Lorenzi, H. 1992. *Árvores Brasileiras*. Nova Odessa: Editora Plantarum. 384pp.
- Ribeiro, J. E. L. da S. 1999. *Flora da Reserva Ducke: Guia de Identificação das plantas Vasculares de Terra-firme na Amazônia Central*. Manaus: INPA. 816pp.
- Rimoli, J. 2001. Ecologia de macacos-pregos (*Cebus apella nigratus*, Goldfuss, 1809) na Estação Biológica de Caratinga, MG: Implicações para a Conservação de Fragmentos da Mata Atlântica. Tese de Doutorado, Centro de Ciências Biológicas, Universidade Federal do Pará.
- Robinson, J. G. e Janson, C. H. 1987. Capuchins, squirrel monkeys and Atelines: socioecological convergence with Old World Primates. Em: *Primate Societies*, Smuts, B.B. et al. (eds.). Chicago: Chicago Society Press.
- Sigg, H. e Stolba, A. 1981. Home Range and Daily March in a Hamadryas Baboon Troop. *Folia Primatol.* 36(1-2): 40–75.
- Silva Júnior, J. S. 2002. Taxonomy of capuchin monkeys, *Cebus* Erxleben, 1777. *Neotrop. Primates* 2(1): 29.
- Sussman, R. W. 2000. *Primate ecology and social structure: New World monkeys*. Needham Heights, Pearson Custom Publishing, 207pp.
- Terborgh, J. 1983. *Five new world primates: a study in comparative ecology*. Princeton University Press, New Jersey.
- Ueno, Y. (1994a). Olfactory discrimination of eight food flavors in the capuchin monkey (*Cebus apella*): Comparison between fruity and fishy odors. *Primates* 35: 301–310.
- Ueno, Y. (1994b). Olfactory discrimination of urine odors from five species by tufted capuchin (*Cebus apella*). *Primates* 35: 311–323.

A CASE OF SPONTANEOUS TOOL-MAKING BY A CAPTIVE CAPUCHIN MONKEY

Tiago Soares Bortolini
Júlio César Bicca-Marques

Introduction

Tool use, the use of a free object in the environment as a functional extension of one's own body (Beck, 1980), has been reported in invertebrates, fish, birds, and mammals, including primates (Alcock, 1989). However, tool use is not common or widespread in nonhuman primates. It has been observed in a small number of species including chimpanzees, bonobos, gorillas, orangutans, some macaques and baboons, and capuchin monkeys (van Schaik et al., 1999). Captive capuchin monkeys (*Cebus* spp.) were long reported to use tools in a variety of contexts (Visalberghi, 1990; Urbani, 1999). Recently, however, a growing body of evidence shows that semi-captive and free-ranging capuchins in several populations and species use tools, including the use of stones as hammers and anvils to crack and open nuts (Otoni and Mannu, 2001; Fragaszy et al., 2004; Moura and Lee, 2004; Waga et al., 2006). Tool-making is a cognitively complex process that involves an intentional modification of the tool for improving its efficiency (Beck, 1980). Reports of non-human primate tool-making have been restricted to the great apes (Boesch and Boesch, 1990; Fontaine et al., 1995; Tomasello and Call, 1997; Schick et al., 1999; van Schaik et al., 2003) with the exception of a few experimentally induced cases in captive capuchins (Westergaard and Suomi, 1994, 1995; Westergaard et al., 1995). Here we report a case of spontaneous tool-making by a captive capuchin monkey.

Methods

A group of capuchin monkeys (*Cebus* sp.) composed of an adult male, two adult females and three immature males living in an enriched enclosure 7.0 m long × 8.7 m wide × 2.9 m high at the Sapucaia do Sul Zoological Park, State of Rio Grande do Sul, Brazil, was opportunistically (*ad libitum*) observed and video-taped in January and February 2007. The enclosure included sand on the floor, trees, stones, and perches for the monkeys. For enrichment purposes, food was concealed inside ice cubes, PVC pipes, and bags.

Results

On 12 January 2007, an adult female (putative *Cebus nigratus*) was observed banging a twig with a piece of stone against a larger stone, licking/chewing and likely extracting something from it with her mouth. She was then observed probing an unseen structure (probably a hole in the enclosure's drinking fountain) with the modified twig (Fig. 1). This sequence of events occurred very rapidly. The latency



Figure 1. Adult female capuchin monkey (putative *Cebus nigritus*) using a hammer stone and an anvil stone to pound on a twig that was then inserted into a crevice (photos from video frames).



Figure 2. Adult female using a stone to crack open an ice cube with food inside.

between the end of banging and the start of probing was 3 to 4 seconds, during which time the female moved from the banging site to the probing site. After this observation, the group was monitored for 15 days and no additional cases of tool-making were observed. The capuchins, however, often used stones as hammers to crack nuts and other foods, including ice cubes containing food (Fig. 2).

Discussion

Although we do not know what happened immediately prior to this behavioral sequence and could not see whether the female acquired anything as a result of probing, the speed at which this sequence of events occurred is highly suggestive of a causal understanding during object manipulation and seems to qualify as a case of spontaneous tool-making. Future research will focus on confirming the capuchin monkeys' capability to make tools, an ability that would suggest less cognitive difference than is presently thought to exist between capuchins and the great apes (Visalberghi, 1990, 1997; Urbani and Garber, 2002). In addition to suggesting that capuchin monkeys understand cause and effect relationships during object manipulation, these findings strengthen the argument that the maintenance of captive animals in enriched environments is an

important strategy to allow the expression of the species' fullest behavioral repertoire. A previous study of the same group (200 hours during 2002–2003) in this enclosure, but with minimum enrichment (paved ground, a single swing and no feeding enrichment), failed to record any case of tool use (D.B. Montano, personal observation). Environmental enrichment serves an important function in improving capuchin monkeys' welfare by reducing boredom and eliciting tool use.

Acknowledgments

We thank the staff of the Núcleo de Zoologia of the Sapucaia do Sul Zoological Park, especially Renato Petry Leal and Marcelo Linck, for the permission to conduct this study. We also thank Paul A. Garber and Bernardo Urbani for reviewing an earlier version of this manuscript.

Tiago Soares Bortolini, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Avenida Bento Gonçalves 9500, Campus do Vale, Prédio 43323 Sala 115, Porto Alegre, RS 91501-970, Brazil, e-mail: <tbortolini@gmail.com>, and **Júlio César Bicca-Marques**, Laboratório de Primatologia, Faculdade de Biociências, Pontifícia Universidade Católica do Rio Grande do Sul, Avenida Ipiranga 6681 Prédio 12A, Porto Alegre, RS 90619-900, Brazil, e-mail: <jbicca@puccrs.br>.

References

- Alcock, J. 1989. *Animal Behavior*. 4th ed. Sinauer, Sunderland.
- Beck, B. B. 1980. *Animal Tool Behavior*. Garland, New York.
- Boesch, C. and Boesch, H. 1990. Tool use and tool making in wild chimpanzees. *Folia Primatol.* 54: 86–99.
- Fontaine, B., Moisson, P. Y. and Wickings, E. J. 1995. Observations of spontaneous tool making and tool use in a captive group of western lowland gorillas (*Gorilla gorilla gorilla*). *Folia Primatol.* 65: 219–223.
- Fragaszy, D., Izar, P., Visalberghi, E., Ottoni, E. B. and Oliveira, M. G. 2004. Wild capuchin monkeys (*Cebus*

- libidinosus*) use anvils and stone pounding tools. *Am. J. Primatol.* 64: 359–366.
- Moura, A. C. A. and Lee, P. 2004. Capuchin stone tool use in Caatinga dry forest. *Science* 306: 1909.
- Otonari, E. B. and Mannu, M. 2001. Semifree-ranging tufted capuchins (*Cebus apella*) spontaneously use tools to crack open nuts. *Int. J. Primatol.* 22: 347–358.
- Schick, K. D., Toth, N., Garufi, G., Savage-Rumbaugh, E. S., Rumbaugh, D. and Sevcik, R. 1999. Continuing investigations into the stone tool-making and tool-using capabilities of a bonobo (*Pan paniscus*). *J. Archaeol. Sci.* 26: 821–832.
- Tomasello, M. and Call, J. 1997. *Primate Cognition*. Oxford University Press, New York.
- Urbani, B. 1999. Spontaneous use of tools by wedge-capped capuchin monkeys (*Cebus olivaceus*). *Folia Primatol.* 70: 172–174.
- Urbani, B. and Garber, P. A. 2002. A stone in their hands... Are monkeys tool users? *Anthropologie* 40: 183–191.
- van Schaik, C. P., Deanes, R. O. and Merrill, M. Y. 1999. The conditions for tool use in primates: Implications for the evolution of material culture. *J. Hum. Evol.* 36: 719–741.
- van Schaik, C. P., Ancrenaz, M., Borgen, G., Galdikas, B., Knott, C. D., Singleton, I., Suzuki, A., Utami, S. S. and Merrill, M. 2003. Orangutan cultures and the evolution of material culture. *Science* 299: 102–105.
- Visalberghi, E. 1990. Tool use in *Cebus*. *Folia Primatol.* 54: 146–154.
- Visalberghi, E. 1997. Success and understanding in cognitive tasks: A comparison between *Cebus apella* and *Pan troglodytes*. *Int. J. Primatol.* 18: 811–830.
- Waga, I. C., Dacier, A. K., Pinha, P. S. and Tavares, M. C. H. 2006. Spontaneous tool use by wild capuchin monkeys (*Cebus libidinosus*) in the Cerrado. *Folia Primatol.* 77: 337–344.
- Westergaard, G. C. and Suomi, S. J. 1994. The use and modification of bone tools by capuchin monkeys. *Curr. Anthropol.* 35: 75–77.
- Westergaard, G. C. and Suomi, S. J. 1995. The manufacture and use of bamboo tools by monkeys: Possible implications for the development of material culture among east Asian hominids. *J. Archaeol. Sci.* 22: 677–681.
- Westergaard, G. C., Greene, J. A., Babitz, M. A. and Suomi, S. J. 1995. Pestle use and modification by tufted capuchins (*Cebus apella*). *Int. J. Primatol.* 16: 643–651.

POWER LINES AND HOWLER MONKEY CONSERVATION IN PORTO ALEGRE, RIO GRANDE DO SUL, BRAZIL

Luisa Xavier Lokschin
Rodrigo Camará Printes
Juliane Nunes Hallal Cabral
Gerson Buss

Introduction

Urban growth affects ecosystems in several ways, leaving them more vulnerable (Alberti and Marzluff, 2004). In Porto Alegre, the combined effects of human presence including deforestation, hunting and other indirect effects are reducing howler's area distribution with consequences still unknown (Lokschin *et al.*, 2005). Human density within a primates' geographical area should be considered by the World Conservation Union (IUCN) in the evaluation of species status (Harcourt and Parks, 2003). The southern brown howler monkey (*Alouatta guariba clamitans*, Cabrera 1940) is considered an endangered species in Rio Grande do Sul (Marques, 2003); in Brazil and globally it is considered near threatened (Machado *et al.*, 2005; Rylands *et al.*, 2006).

There are many species of Neotropical primates living close to urban areas, including: *Alouatta clamitans* (Buss, 1996), *Alouatta caraya* (Codenotti *et al.*, 2002), *Callicebus nigrifrons* (Oliveira *et al.*, 2003), *Saguinus leucopus* (Poveda and Sánchez-Palomino, 2004) and *Saguinus bicolor* (Vasconcelos *et al.*, 2005). Problems and threats linked to urbanization, such as danger from vehicles when crossing roads, predation by dogs and electric hazard, are already documented for *A. clamitans* (Printes, 1999; Alonso *et al.*, 2005), *C. jacchus* (Menezes, 2005) and *S. bicolor* (Vasconcelos *et al.*, 2005). Ecosystems close to urban areas are important for wildlife (Dickman, 1987) and measures must be taken to guarantee their existence. Howler monkeys (*A. g. clamitans*) utilize areas of forests close to urban developments and are suffering from contact with several electric hazards. Here we describe a way to mitigate the occurrence of such accidents around Porto Alegre.

Methods

Porto Alegre is the capital city of the state of Rio Grande do Sul, Brazil (Fig. 1), with a population of 1.4 million (IBGE, 2006). Approximately 10% of the municipal area is natural semi-deciduous seasonal forest, influenced by Atlantic rainforest (Brack *et al.*, 1998; Velez *et al.*, 1998). The southernmost area of the municipality (Fig. 2) is a rural landscape containing a number of small villages. The most important natural areas are also in this zone, which is also the most important area for howler monkeys (*Alouatta guariba clamitans*) (Romanowski *et al.*, 1998; Lokschin *et al.*, 2005). Lami Biological Reserve, the only biological