

Morphometric Data from a Wild Female Titi Monkey, *Callicebus cupreus*

Authors: Heymann, Eckhard W., Yamacita, Jenni G. Pérez, and Müller, Britta

Source: Neotropical Primates, 19(1) : 42-43

Published By: Conservation International

URL: <https://doi.org/10.1896/044.019.0110>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

- Link, A., Palma, A. C., Velez, A. and De Luna, A. G. 2006. Costs of twins in free-ranging white-bellied spider monkeys (*Ateles belzebuth*) at Tinigua National Park, Colombia. *Primates* 47:131–139.
- van Schaik, C. 2000. Infanticide by male primates: the sexual selection hypothesis revisited. In: *Infanticide by Males and its Implications*, C. van Schaik and C. H. Janson (eds.), pp. 27–60. Cambridge University Press, Cambridge.

MORPHOMETRIC DATA FROM A WILD FEMALE TITI MONKEY, *CALLICEBUS CUPREUS*

Eckhard W. Heymann
Jenni G. Pérez Yamacita
Britta Müller

Body size and the size of organs and other body structures are intimately related to the life history and ecology of a species (Temerin et al. 1984). Comparative analyses of data from different taxa can reveal allometric relationships and the adaptive value of different body or organ size (Martin 1990, Terborgh 1983). E.g., Terborgh (1992) showed the

Table 1. Morphometric measurements from the *Callicebus cupreus* female and comparative data from the literature

	This report	Hill 1960	Hershkovitz 1990	Bicca-Marques et al. 2002	Peres 1993	Ferrari & Lopes 1995
External measurements						
Body mass [g]	720		1106 (1000-1175)*	750, 900	860, 970*	880 [#] , 1020 [§]
Head-body length [mm]	285	325, 310	337 (270-410)	280, 310	302, 358 [‡]	
Tail length [mm]	475	440, 420	439 (405-470)	340, 440	412, 414*	
Hind foot length [mm]	89	100, 95	92 (85-100)		89, 91*	
Skull length [mm]	67.7	65 [†]	63.9 (60.0-66.8)			
Zygomatic breadth [mm]	42.2	41.25	39.0 (36.0-42.0)			
Orbital breadth [mm]	37.2					
Braincase width [mm]	39.8					
Postorbital restriction [mm]	31.4					
Across molars [mm]	19.9					
C ¹ - C ¹ [mm]	12.0		13.7 (12.9-15.0)			
C1 - C1 [mm]	8.7					
P ² - M ³ [mm]	15.1	15.4				
M ¹ (right) breadth [mm]	4.4					
M ² (right) breadth [mm]	4.1					
M ³ (right) breadth [mm]	3.3					
Mandibular height [mm]	35.4					
Orbita height [mm]	1.7					
Orbita breadth [mm]	1.4					
Internal organs						
Liver mass [g]	25.5					
Kidney mass [g], right	3.5					
left	4.8					
Adrenal length [mm], right	5					
left	7					
Spleen mass [g]	3.25					
Small intestine length [mm]	950					944 [#] , 1056 [§]
Caecum length [mm]	100					
Large intestine length [mm]	435					324 [#] , 521 [§]
Pluck (lung, heart, trachea, tongue) [g]	9.75					

* Data for male *C. cupreus* only; [#] *Callicebus caligatus*; [§] *Callicebus moloch*; [†] Hill (1960) provides a mean of two males and three females; [‡] Peres (1993) gives 716 and 770 mm, but this is likely to be head-body-tail length, values listed here are therefore the value given by Peres minus tail length

relationship between primate body size and dietary strategies. Ferrari et al. (1993) compared gut proportions of a specialized and an opportunistic gum feeder and showed the former to have a comparatively larger caecum, as an adaptation to the fermentation of gums. Morphometric data from skulls and skeletons can be obtained from museum material, but data on fresh body mass and on organ size or mass are generally not available from museum specimen. Given ethical implications and the increasing threat to wild primates, collecting wild primates for the purpose of obtaining organ size data is prohibitive. It is therefore imperative to exploit opportunities for taking morphometric data without collection, e.g. when a fresh carcass is found.

In this paper, we report morphometric data from a wild female red titi monkey, *Callicebus cupreus*, at the Estación Biológica Quebrada Blanco (EBQB) in north-eastern Peruvian Amazonia. This female was a member of one study group and found in a comatose state below the sleeping tree on early morning of 29 September 2002. It died a few hours later and was subjected to a field necropsy by the senior author, a trained veterinarian. For a detailed case report and the pathological findings see Müller et al. (2010). We measured body mass, head-body length, tail length and hind foot length on the fresh carcass with Pesola spring balances and a Vernor calliper, respectively. Skull length was measured before the braincase was opened for brain inspection and removal. After necropsy, skull and skeletal material were buried, to allow for decomposition of flesh, and recovered later. Unfortunately, part of the material was taken by scavengers. The remaining material was stored at EBQB and measured with a Mitutoyo CD-20DCX digital calliper by the first author in October 2012. Each variable was measured three times with and values were averaged.

Morphometric data collected before and during necropsy, and from the skull and skeletal material are provided in Table 1, together with data compiled from the literature. For many measurements taken here, actually no comparative data are available from the literature. Most skeletal and dental measurements are within or close to the range of values reported in the literature. The comparatively low body mass is likely due to the diseased condition. It should be noted that data for liver, spleen and adrenal, perhaps also for kidney may also represent pathological conditions (Müller et al. 2010).

Acknowledgements

The study during which these data were collected was supported by a grant from the Deutsche Forschungsgemeinschaft to EWH (DFG He 1870/13-1) and from the Deutsche Akademische Austauschdienst (DAAD) to BM, and carried out under permission from the Instituto Nacional de Recursos Naturales (INRENA) in Lima (Peru). We thank Ney Shahuano Tello for field assistance, and biology students Victor Raygada Guerra and Cristina Lopez Wong

from the Universidad Nacional de la Amazonía Peruana in Iquitos (Peru) for their help with the field necropsy.

Eckhard W. Heymann, Abteilung Verhaltensökologie & Soziobiologie, Deutsches Primatenzentrum, Kellnerweg 4, 37077 Göttingen, Germany, e-mail: <eheman@gwdg.de>; **Jenni G. Pérez Yamacita**, Facultad de Ciencias Biológicas, Universidad Nacional de la Amazonía Peruana, Iquitos, Perú; **Britta Müller**, Abteilung Verhaltensökologie & Soziobiologie, Deutsches Primatenzentrum, Kellnerweg 4, 37077 Göttingen, Germany, and Abteilung Infektionspathologie, Deutsches Primatenzentrum, Kellnerweg 4, 37077 Göttingen, Germany, and Bayerisches Landesamt für Gesundheit und Lebensmittelsicherheit, Veterinärstraße 2, 85764 Oberschleißheim, Germany.

References

- Bicca-Marques, J. C., Garber, P. A. and Azevedo-Lopes, M. A. O. 2002. Evidence of three resident adult male group members in a species of monogamous primates, the red titi monkey (*Callicebus cupreus*). *Mammalia* 66: 138–142.
- Ferrari, S. F. and Lopes, M. A. 1995. Comparison of gut proportions in four small bodied Amazonian cebids. *Am. J. Primatol.* 35: 139–142.
- Ferrari, S. F., Lopes, M. A. and Krause, E. A. K. 1993. Gut morphology of *Callithrix nigriceps* and *Saguinus labiatus* from Western Brazilian Amazonia. *Am. J. Phys. Anthropol.* 90: 487–493.
- Hershkovitz, P. 1990. Titis, New World monkeys of the genus *Callicebus* (Cebidae, Platyrrhini): a preliminary taxonomic review. *Fieldiana Zool.* 55: 1–109.
- Hill, W. C. O. 1960. *Primates. Comparative anatomy and taxonomy. IV. Cebidae Part A.* Edinburgh University Press, Edinburgh.
- Martin, R. D. 1990. *Primate origins and evolution.* Chapman and Hall, London.
- Müller, B., Mätz-Rensing, K., Pérez Yamacita, J. G. and Heymann, E. W. 2010. Pathological and parasitological findings in a wild red titi monkey, *Callicebus cupreus* (Pitheciidae, Platyrrhini). *Eur. J. Wildl. Res.* 56: 601–604.
- Peres, C.A. 1993. Notes on the primates of the Juruá river, western Brazilian Amazonia. *Folia Primatol.* 61:97–103.
- Temerin, L. A., Wheatley, B. P. and Rodman, P. S. 1984. Body size and foraging in primates. In: *Adaptations for foraging in nonhuman primates*, P. S. Rodman and J. G. H. (eds.), pp. 217–248. Columbia University Press, New York.
- Terborgh, J. 1983. *Five New World primates. A study in comparative ecology.* Princeton University Press, Princeton.
- Terborgh, J. 1992. *Diversity and the tropical rain forest.* Scientific American Library, New York.