

Report of a Black Spider Monkey (*Ateles chamek*) Swimming in a Large River in Central-Western Brazil

Author: Nunes, André Valle

Source: Neotropical Primates, 21(2) : 204-206

Published By: Conservation International

URL: <https://doi.org/10.1896/044.021.0210>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REPORT OF A BLACK SPIDER MONKEY (*ATELES CHAMEK*) SWIMMING IN A LARGE RIVER IN CENTRAL-WESTERN BRAZIL

André Valle Nunes

Rivers are considered to influence the current patterns of ecological and genetic variation of Amazonian species and communities (Gascon *et al.*, 2000). Rivers are impenetrable barriers to the dispersal of several vertebrate species, and, in some cases, they interrupt the expansion of species from their origin centers and gene flow between populations of sister species from opposite margins (Sick, 1967; Hershkovitz, 1977; Ayres and Clutton-Brock, 1992; Gascon *et al.*, 2000). Hence, rivers drive, inter and intraspecific differentiation patterns in the distribution of Amazonian primates (Ayres and Clutton-Brock, 1992).

There are few records of New World primate species swimming (Parnell and Buchanan-Smith, 2001). Some platyrrhines, such as *Cebus*, *Cacajao*, *Aotus*, and *Saimiri*, can inhabit or use swamps and seasonally flooded areas, but they do not swim between habitat patches (Socoloske and Kymberley, 2010; Bezerra *et al.*, 2010). The only reports of swimming refer to large primates, such as *Alouatta palliata*, *Cacajao melanocephalus*, as well as large species of Old World cercopithecoids: *Macaca radiata* (bonnet macaque), *Nasalis larvatus* (proboscis monkeys), *Papio anubis* (olive baboon), *Macaca fuscata* (japanese macaques), *Pan troglodytes* (chimpanzee), and *Pongo pygmaeus* (bornean orangutan) (Wata, 1981; Yeager, 1991; Forthman, 2000; Agormoorthy *et al.*, 2000; Gonzalez-Socoloske and Snarr, 2010; Bezerra *et al.*, 2010; Bender and Bender, 2013). Among the largest platyrrhines are the atelids, such as the black spider monkey (*Ateles chamek*), which may weight up to 10 kg and has two common characteristics of the subfamily Atelinae: immigration and movement through semibrachiation (Chapman and Chapman, 1989; Chapman and Chapman, 1990; Campbell *et al.* 2005). *Ateles chamek* occurs in lowlands from northeastern Peru, northern and central Bolivia in the Noel Kempf Mercado National Park, and western Brazil in the state of Mato Grosso on the left margins of the rivers Teles Pires and Tapajós (Wallace *et al.*, 1996; Iwanaga and Ferrari, 2002). There are reports of black spider monkeys using the ground to socialize, collect food, and cross open areas (Di Fiore, 2002; Campbell *et al.*, 2005). However, there is no information on swimming. Therefore, we report a rare case of swimming by a female *A. chamek* in an Amazonian River.

On November 28th, 2012, in the municipality of Comodoro (13°47'54"S, 60°27'53"W), in the Amazon of Mato Grosso, we observed a female *A. chamek* crossing the Guaporé River by swimming. Approximately at 16:30 h, the female began moving on the ground of the river's beach, which is located within the Noel Kempf Mercado




Figure 1. Location of the swimming of black spider monkey (*Ateles chamek*) in Guaporé River, border between Bolivia and Brazil in South America. Tropical and subtropical moist forests; Tropical and subtropical savannas; Floodplains; Tropical and subtropical dry forests; Mountain grasslands. Habitat types follow Olson *et al.* (2001).



Figure 2. Photography of the adult female black spider monkey (*Ateles chamek*) swimming towards the shore of the state of Mato Grosso, Brazil. Photo Credit: Antônio Linares



Figure 3. Locomotion of the adult female black spider monkey (*Ateles chamek*) on a submerged branch prior to climbing to tree crowns in the riparian forest of the Guaporé River in the state of Mato Grosso, Brazil.

National Park in the municipality of Santa Cruz, Bolivia (Fig. 1). Next, the specimen began to swim towards the opposite margin, located in the state of Mato Grosso, Brazil. The swimming activity lasted ca. 15 min, and the specimen crossed 38 meters from one margin to the other. The specimen had the body completely submerged, leaving only the head out of the water (Fig. 2), and moved its arms and legs. It was panting and was not scared by our boat; on the contrary, it even climbed onto the boat and walked around the boat's edge before immersing back into the water. Then, the specimen managed to climb onto a submerged branch and began a fast movement between tree crowns in the riparian forest of the Guaporé River, on the Brazilian side (Fig. 3).

The present report is consistent with the meta-analysis by Ayres and Clutton-Brock (1992), which assessed the relationship between the width of Amazonian rivers and the body weight of primates. This meta-analysis assumes that larger species are less affected by ecological barriers, which means that speciation and interspecific differences between primate communities in the Amazon may be correlated with the ability of the species to cross ecological barriers. Another factor that could have favored swimming by the female *A. chamek* is the morphodynamics of the Guaporé River. According to Souza-Filho *et al.* (1999) the Guaporé River has a fluvial meandering channel with the formation of beaches in the dry season. Rivers with sinuous conformation and sand banks on their margins allow animals to cross them, which lead to possibility of crossing of terrestrial vertebrates between opposite margins (Ayres and Clutton-Brock, 1992). In this scenario, the breaking of a physical barrier may favor gene flow between sister species of primates that evolved in opposite margins of Amazonian rivers and help us understand species distributions.

Acknowledgements

The present study received logistic support from Biocev and Environmental Consulting Services. We thank Leandro S. Moreira and Leandro Scoss for comments on the manuscript, and Antônio Meira Linares for providing us with the photos of *Ateles chamek*.

André Valle Nunes, Museu de Zoologia João Moojen, Departamento de Biologia Animal, Universidade Federal de Viçosa. Viçosa – MG, Brazil.

References

- Agoramoorthy, G., Smallegange, I., Spruit, I and Hsu, M. J. 2000. Swimming behavior among bonnet macaques in Tamil Nadu: A preliminary description of a new phenomenon in India. *Folia Primatol.* 71:152–153.
- Ayres, J. M. and Clutton-Brock, T. H. 1992. River boundaries and species range size in Amazonian. *Am. Natural.* 140(3):531–537.
- Bender, R. and Bender, N. 2013. Swimming and diving behavior in apes (*Pan troglodytes* and *Pongo*

- pygmaeus*) first documented report. *Am. J. Phys. Anthropol.* 152(1):156–162.
- Bezerra, B. M., Barnett, A. A., Souto, A. and Jones, G. In press. Ethogram and natural history of golden-backed uakaris (*Cacajao melanocephalus*). *Int. J. Primatol.* 32(1):46–68.
- Campbell, C. J., Aureli, F., Chapman, C. A., Ramos-Fernández, G., Matthews, K., Russo, S. E., Soares, S. and Vick, L. 2005. Terrestrial behavior of *Ateles* spp. *Int. J. Primatol.* 26(5):1039–1051.
- Chapman, C. A. and Chapman, L. J. 1989. Primate populations in north-western Costa Rica: Potential for recovery. *Primate Conserv.* 10:37–44.
- Chapman, C. A. and Chapman, L. J. 1990. Reproductive biology of captive and free ranging spider monkeys. *Zoo Biology.* 9:1–9.
- Di Fiore, A. 2002. Predator sensitive foraging in the ateline primates. In: *Predator Sensitive Foraging Among Primates*, L. Miller (ed.), pp. 242–267. Cambridge University Press, Cambridge.
- Forthman, D. 1999–2000. Note on water ‘play’ by an olive baboon (*Papio Anubis*) in Gilgil, Kenya. *Afr. Primates* 4(1&2):74.
- Gascon, C., Malcolm, J. R., Patton, J. L., Silva, M. N. F., Bogart, J. P., Loughheed, S. C., Peres, C. A., Neckel, S. and Bogart, P. T. 2000. Riverine barriers and the geographic distribution of Amazonian species. *P. Natl. Acad. Sci. (USA)* 97(25):13672–13677.
- Gonzalez-Socoloske, D. and Snarr, K. A. 2010. An incidente of swimming in a large river by a manthled howling monkey (*Alouatta palliata*) on the north coast of Honduras. *Neotrop. Primates.* 17(1):28–31.
- Hershkovitz, P. 1977. *Living New World monkeys (Platyrrhini) with an introduction to primates*. Vol. 1. University of Chicago Press, Chicago.
- Iwanaga, S. and Ferrari, S. F. 2002. Geographic distribution and abundance of woolly (*Lagothrix cana*) and spider (*Ateles chamek*) monkeys in southwestern Brazilian Amazonia. *Am. J. Primatol.* 56:57–64.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D’Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P. and Kassem, K. R. 2001. Terrestrial ecoregions of the world. A new map of life on Earth. *Bioscience.* 51:933–938.
- Parnell, J. P. and Buchanan-Smith, H. M. 2001. An unusual social display by gorillas. *Nature.* 142:294.
- Sick, H. 1967. Rios e enchentes na Amazônia como obstáculo para a avifauna. Manaus, *Atas Simpósio sobre a biota Amazônica*, 5:495–520.
- Souza-Filho, P. W. M., Quadros, M. L. E. S., Scandolara, J. E., Silva-Filho, E. P. and Reis, M. R. 1999. Compartimentação morfoestrutural e neotectônica do sistema fluvial Guaporé-Mamoré Alto Madeira, Rondônia, Brasil. *Revis. Brasil. Geociên.* 29(4):469–476.
- Wallace, R. B., Painter, R. L. E. and Taber, A. B. 1996. Primate diversity, habitat preferences, and population density estimates in Noel Kempff Mercado National Park, Santa Cruz Department, Bolivia. *Am. J. Phys. Anthropol.* 46(3):197–211.
- Wata, k. 1981. Habitat utilization by wintering Japanese macaques (*Macaca fuscata fuscata*) in the Shiga Heights. *Primates.* 22(3):330–348.
- Yeager, C. P. 1991. Possible antipredator behavior associated with river crossings by proboscis monkeys (*Nasalis larvatus*). *Am. J. Primatol.* 24(1):61–66.