



New Northwestern and Southwestern Range Limits of De Brazza's Monkey, Mbam et Djerem National Park, Cameroon, and Bateke Plateau, Gabon and Congo

Authors: Maisels, Fiona, Bout, Nicolas, Inkamba-Inkulu, Clement, Pearson, Liz, Aczel, Paul, et al.

Source: Primate Conservation, 22(1) : 107-110

Published By: Conservation International

URL: <https://doi.org/10.1896/052.022.0109>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

New Northwestern and Southwestern Range Limits of De Brazza's Monkey, Mbam et Djerem National Park, Cameroon, and Bateke Plateau, Gabon and Congo

Fiona Maisels¹, Nicolas Bout^{2,3}, Clement Inkamba-Inkulu⁴, Liz Pearson³, Paul Aczel³, Rufin Ambahe⁵, Edgar Ambassa⁵ and Roger Fotso⁵

¹Wildlife Conservation Society, Bronx, New York, USA

²Wildlife Conservation Society, Libreville, Gabon

³John Aspinall Foundation, Franceville/Mpassa, Gabon

⁴Wildlife Conservation Society, Brazzaville, Republic of Congo

⁵Wildlife Conservation Society, Yaoundé, Cameroon

Abstract: Recent surveys carried out in Cameroon in the Mbam Djerem National Park, in Gabon in the Bateke Plateau National Park, and the adjoining Bateke Plateau area in Congo, have recorded the presence of de Brazza's monkey (*Cercopithecus neglectus*) on both sides of the Djerem River in Cameroon, along the Mpassa and its tributaries in Gabon, along the Nambouli River in the Lefini Reserve in Congo, and up to the right bank of the Ogooué River on the Congo side of the border. These areas lie at the northern and southern edges of the Central African forest block, where rivers have relatively wide bands of riparian forest. As for the range extension of the talapoin (this volume), the species may simply have been overlooked by previous rapid wildlife surveys. It has an antipredator behavior that renders it relatively inconspicuous, tends to live in small family groups in the region, and lives in habitats that are difficult to survey on foot. However, unlike talapoin, it calls every morning along major watercourses, and can be heard for some distance. It is possible that the southern limit of this species in Gabon may be the Ogooué River. Future survey teams are encouraged to familiarize themselves with the long call of this species and to be aware that it can occur in gallery forests throughout the savannas of the Bateke Plateau and also in the area between the Mbam, the Djerem, and the Lom in Cameroon.

Résumé: Les recensements récents au Cameroun dans le parc national de Mbam et Djerem, au Gabon dans le parc national de Plateau Bateke, et dans les savanes Bateke avoisinante au Congo ont notée la présence du singe de Brazza (*Cercopithecus neglectus*) sur les deux rives du Djerem au Cameroun, le long de l'Mpassa et ses tributaires au Gabon, le long de la rivière Nambouli dans la Reserve de la Lefini au Congo, et jusqu'au rive droite de la rivière Ogooué sur la coté Congolaise de la frontière. Les deux zones se trouvent aux bords nord et sud du grand bloc forestier du bassin du Congo, et contiennent les rivières importantes, qui coulent dans les galeries assez larges de forêt ripicole. Comme pour les talapains (ce volume), l'espèce a été peut-être simplement ratée par les équipes de recensement dans la région auparavant. Il a un comportement anti-prédateur qui le rend très discret, une tendance à vivre par petits groupes familiaux, difficiles à repérer, de plus occupe un habitat plus difficilement accessible à un observateur à pied les habitats de la plupart des autres guenons. Néanmoins, et contrairement au comportement des talapains, chaque matin il pousse des cris très caractéristiques, audible sur des grandes distances le long des grandes rivières. Il est possible que la limite sud de cette espèce au Gabon soit la rivière Ogooué. Les équipes de recensement de la faune sont encouragées de se familiariser avec les vocalisations de cette espèce, et garder à l'esprit que le singe de Brazza peut être présent dans les forêts galeries dans toutes les savanes du plateau Batéké et aussi, au Cameroun, la zone entre les rivières Mbam, Djerem, et Lom.

Key Words: de Brazza's monkey, biogeography, distribution, Gabon, Cameroon, Congo

Introduction

De Brazza's monkey (*Cercopithecus neglectus*) is widely distributed in Central Africa, from the Atlantic coast of southern Cameroon, through Equatorial Guinea down to northern Gabon, and then eastward through the forest block and some of the fringing transitional forests to the Ruwenzoris (Kingdon 1997; IEA 1998). In West Central Africa it has been assumed that their distribution is limited to the north, as it is in many other species, by the Sanaga River. The southern limit is less clear. It was earlier thought that the species did not occur south of the Ivindo watershed (Malbrant and Maclatchy 1949) but later maps show them to half way between the Ivindo and the Ogooué (Kingdon 1997; IEA 1998). The published maps (Kingdon 1997; IEA 1998) suggest that they are absent from the Bateke Plateau and from the "Lake region" between the Komo and the Ogooué rivers (Fig. 1).

Observations

The Mbam et Djerem National Park was created in 2000 in central Cameroon. It lies across the Guinea-Congolia/

Sudania regional transition zone (White 1983) between the main forest block and the Guinean savannas to the north (Fig. 1). The Djerem River, which is, in fact, the upper reaches of the Sanaga, runs north-south through this park, and is lined by a broad band of riparian forest. The southeastern part of the park is lowland, tropical, semideciduous forest; the north-western part is wooded and bush savannah with gallery forests lining the watercourses. The transition between the forest and the savannah has a relatively high biodiversity because of the interlacing of the two main habitats. The forest has been slowly creeping north since the 1950s (evidenced by the old maps of the region and present conditions, which show the contrast between savannah in the past and young colonizing woodland today).

Another new national park, Bateke Plateau, was created in 2002, in southeastern Gabon. The vegetation is structurally similar to that of Mbam et Djerem: The main forest block of Gabon is at the extreme northwest of the park and there is a progression through forest-savannah mosaics to bush savannah and some extensive areas of grass savannah toward the east, at the Gabon-Republic of Congo international border. The Bateke Plateau in general is a savannah and gallery forest mosaic that extends far to the east, across the Republic of Congo, into western Democratic Republic of Congo. The Lefini Reserve is also in the Bateke Plateau, a little farther south than the Gabonese Bateke National Park (Fig. 1) and has the same vegetation: a mosaic of gallery forests and savannas. The Bateke Plateau National Park in Gabon is bisected north-south by the Mpassa River, which like the Djerem in Cameroon, runs through a wide corridor of riparian forest.

A series of wildlife and vegetation surveys were carried out in all three protected areas (Mbam et Djerem, Cameroon; Bateke National Park, Gabon; and the Lefini Reserve, Congo) between 2000–2005, partly as reconnaissance missions but mostly to provide the baseline for the park monitoring of wildlife and human impact. In addition, in the Congolese part of the Bateke Plateau just across the border from Gabon, a wildlife and socioeconomic survey was carried out in 2004 to evaluate the potential for an additional protected area that could form a transboundary park in the future.

De Brazza's monkeys were recorded nine times along the Djerem River and three times on various tributaries in Mbam et Djerem. In the Bateke Plateau (Gabon) it was recorded at least a dozen times: in the riparian forest along the Mpassa River and on eight different tributaries. In the Bateke Plateau (Congo) it was seen along two of the upper tributaries of the Congo (just across the frontier, south of the village of Edjangoulou: Fig. 1), several times on the Ogooué River, and twice in the Lefini Reserve along the Nambouli River.

We also asked local hunters about the monkey species present in the areas. In Mbam et Djerem de Brazza's monkeys are well known by their (onomatopoeic) local name, *founga*, in Gbaya (very similar to the Mboko name *pounga* from Odzala in Congo [F. Maisels pers. obs.] and to the Bakota name, *pounnga*, in Gabon [Malbrant and Maclatchy 1949]). By contrast, the Bateke hunters in Gabon ascribed the




Figure 1. Range of *Cercopithecus neglectus*, after Kingdon (1997) and IEA (1988), and the recent records from Mbam et Djerem National Park, Cameroon; the Lefini Reserve, Congo; and the Bateke Plateau area in Gabon and Congo.

characteristic long call of the species to *Cercopithecus cephus*. However, one Gabonese hunter knew the species and its calls, and noted that it was found also at Edjangoulou, about 60 km farther north on the plateau, corroborating the observations on the Congo side (Fig. 1). The Gabonese Teké name is *mfoue*; the Congolese Teké name is *nfua*.

Discussion

These records show that de Brazza's monkey occurs about 100 km farther north in Cameroon, about 140 km farther south in Gabon, and about 80 km farther west in Congo than was previously known. However to the east (Central African Republic and Republic of Congo), it occurs (or occurred) much farther north and south, respectively, so it is likely that this species was simply missed during surveys in the past. Downer (1998) carried out a wildlife survey and also questioned hunters in the Lefini Reserve in 1996 in Republic of Congo. Although the species was not recorded during the survey, the hunters said that de Brazza's monkeys occurred there, which was confirmed during the 2005 survey. The Bateke Plateau is heavily hunted (especially on the Congo side) so monkeys in the area will be very wary and in probably much reduced in numbers.

De Brazza's monkey is considerably more inconspicuous than most other guenons. It generally avoids living in polyspecific troops, and at least in Gabon, northern Congo, the Bateke Plateau and central Cameroon, lives in small family parties (Gautier-Hion 1988; N. Bout, C. Inkamba-Nkulu and F. Maisels pers. obs.). In areas where they are hunted, they freeze, being motionless sometimes for hours when disturbed, and hide their otherwise highly visible white beard and buttocks by crouching (Gautier-Hion and Gautier 1978). Like the talapoin, *Miopithecus*, and Allen's swamp monkey, *Allenopithecus nigroviridis*, their habitat is inundated forest along watercourses, which makes them less likely than terrestrial species to be seen by survey teams walking through forests and savannas. In fact, they are most often detected by their calls in the early morning, or from a canoe while traveling along rivers. To people unfamiliar with their vocalizations, the "boom" could be confused with that of *Mona* superspecies (*C. mona* in the northwest and *C. pogonias* in most other places west of the Congo River [Lernould 1988; Kingdon 1997; IEA 1998]). Recordings of *C. neglectus* calls can be found on the CD of primate vocalizations in Gautier-Hion *et al.* (1999), and it is important to note that (1) the two booms are longer than those of *C. mona*; (2) they are separated by a soft, audible sub-boom caused by the re-inflation of the vocal sac, and (3) that the "cough" following the booms is a series of single notes, unlike the rapid double cough of *C. mona*.

Although de Brazza's monkeys are not listed as threatened by the IUCN, they are nevertheless listed in Appendix II of CITES (Inskipp and Gillett 2005). At least in parts of the Bateke Plateau, it is clear that they are uncommon and very wary of humans; despite their antipredator behavior they are vulnerable to hunters if they are detected. We urge

wildlife survey teams to be aware that de Brazza's monkeys may be present along rivers in the savannas of the region, and to familiarize themselves with the vocalizations of this rather secretive species. In Gabon, the species almost certainly occurs up to the right bank of the Ogooué River in the Bateke Plateau area, and perhaps also in the very marshy lake region between the Komo and the Ogooué to the west of the country. Unless it has been hunted out, in Congo it is almost certainly present along watercourses across the whole of the Bateke Plateau between the Congo River and the Ogooué. In Cameroon, attention should be paid to the possibility of this species occurring along the major watercourses to the east and west of the Djerem: the Mbam and the Lom (Fig. 1).

Acknowledgments

We thank the Governments of Cameroon (MINFOF), Gabon (CNP and the DFC) and Republic of Congo (DFAP) for their efforts in wildlife conservation. For funding, we are grateful to the following: USAID (Congo Basin Forest Project); FEDEC (Foundation for Environment and Development in Cameroon); the British High Commission, Yaoundé: Global Opportunities Funds; the John Aspinall Foundation; WCS New York; WCS Cameroon; WCS Gabon; and WCS Congo. Thanks to all the staff of the projects at Mbam et Djerem, Bateke Plateau (both Gabon and Congo), Lefini, the JAF-Mpassa project, WCS Cameroon, WCS Congo, and WCS Gabon. Finally, enormous thanks to the trackers and porters from all sites.

Literature Cited

- Downer, A. 1998. A series of preliminary surveys: Ecological, demographic, socio-economic and their implications as regards the sustainability of the Lefini Game Reserve. Report, Wildlife Conservation Society (WCS) – Congo, Brazzaville.
- Gautier-Hion, A. 1988. Polyspecific associations among forest guenons: Ecological, behavioural and evolutionary aspects. In: *A Primate Radiation. Evolutionary Biology of the African Guenons*, A. Gautier-Hion, F. Bourlière, J. P. Gautier and J. Kingdon (eds.), pp.452–476. Cambridge University Press, Cambridge, UK.
- Gautier-Hion, A. and J. P. Gautier. 1978. Le singe de Brazza: Une stratégie originale. *Z. Tierpsychol.* 46: 84–104.
- Gautier-Hion, A., M. Colyn and J.-P. Gautier. 1999. *Histoire Naturelle des Primates d'Afrique Centrale*. ECOFAC, Libreville, Gabon.
- IEA. 1998. *Cercopithecus neglectus. African Mammals Databank—A Databank for the Conservation and Management of the African Mammals*. Vols. 1 and 2. Institute of Applied Ecology (IEA), European Commission Directorate, Brussels. Website: <<http://gorilla.bio.uniroma1.it/amd/amd072.html>>. Accessed 14 July 2006.
- Inskipp, T. and H. J. Gillett (eds.). 2005. *Checklist of CITES Species and Annotated CITES Appendices and*

- Reservations*. CITES Secretariat, Geneva, Switzerland and UNEP-WCMC, Cambridge, UK. 339pp. and CD-ROM.
- Kingdon, J. 1997. *The Kingdon Field Guide to African Mammals*. Academic Press, London.
- Lernould, J.-M. 1988. Classification and geographical distribution of guenons: A review. In: *A Primate Radiation: Evolutionary Biology of the African Guenons*, A. Gautier-Hion, F. Bourlière, J.-P. Gautier and J. Kingdon (eds.), pp.54–78. Cambridge University Press, Cambridge, UK.
- Malbrant R. and A. Maclatchy. 1949. *Faune de l'Equateur Africain Francais. Tome II. Mammifères*. Paul Lechevalier, Paris.
- White, F. 1983. *The Vegetation of Africa*. UNESCO, Paris.

Authors' addresses:

Fiona Maisels, Wildlife Conservation Society, 2300 Southern Boulevard, Bronx, New York, NY 10460-1099, USA. E-mail: <fmaisels@wcs.org>.

Nicolas Bout, WCS Gabon, BP 7847, Libreville, Gabon *and* John Aspinall Foundation, John Aspinall Foundation, BP 583, Franceville, Gabon.

Clement Inkamba-Inkulu, presently at University of East Anglia, Norwich NR4 7TJ, UK.

Liz Pearson and **Paul Aczel**, John Aspinall Foundation, BP 583, Franceville, Gabon.

Rufin Ambahe, **Edgar Ambassa** and **Roger Fotso**, Wildlife Conservation Society, BP 3055, Yaoundé, Cameroon.

Received for publication: 5 August 2006

Revised: 10 August 2006