

# Using the reference condition maintains the integrity of a bioassessment program in a changing climate

Author: Nichols, Susan J.

Source: Journal of the North American Benthological Society, 29(4): 1459-1471

Published By: Society for Freshwater Science

URL: https://doi.org/10.1899/09-165.1

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Using the reference condition maintains the integrity of a bioassessment program in a changing climate

#### Susan J. Nichols<sup>1</sup>

Institute for Applied Ecology and eWater Cooperative Research Centre, University of Canberra, Canberra, Australian Capital Territory, 2601 Australia

#### Wayne A. Robinson<sup>2</sup>

School of Environmental Sciences, Charles Sturt University, Thurgoona, New South Wales, 2641 Australia

#### Richard H. Norris<sup>3</sup>

Institute for Applied Ecology and eWater Cooperative Research Centre, University of Canberra, Canberra, Australian Capital Territory, 2601 Australia

Abstract. Climate change is gradual and long-term, consistently collected data are required to detect resulting biological responses and to separate such responses from local effects of human activities that monitoring programs usually are designed to assess. The reference-condition approach is commonly used in freshwater assessments that use predictive modeling, but a consistent reference condition is required to maintain the relevance and integrity of results over the long term. We investigated whether external influences, such as climate change, inhibited clear interpretation of bioassessment results in a study design using reference vs test sites. Macroinvertebrates were collected from 16 sites (11 sites affected by ski resorts and 5 reference sites) on 5 streams in 4 seasons each year from 1994 to 2008 within Kosciuszko National Park, Australia. We analyzed trends over 15 y to address questions regarding climate-change and macroinvertebrate bioindicators of stream condition (observed/expected [O/E] taxa; Stream Invertebrate Grade Number Average Level [SIGNAL] 2 scores; Simpson's Diversity; Ephemeroptera, Plecoptera, Trichoptera [EPT] richness ratio; and Oligochaeta abundance). Climate became slightly warmer and less humid (p < 0.0001), but no significant relationships between climate variables and bioindicators were evident. All bioindicators consistently distinguished between test and reference sites in all seasons. All bioindicators except for O/E taxa scores differed among streams (regardless of site type). O/E taxa are inherently adjusted for specific stream characteristics, and, thus, were robust to differences in stream type while remaining sensitive to reference and test site variation. Generally, reference and test sites did not respond differently to any gradual climate changes. Furthermore, the reference sites sampled through time remained in a condition equivalent to the previously defined reference condition and provided a valid comparison for current test sites of unknown condition. The bioindicators used here were insensitive to the small but significant changes in climate detected over the 15-y study. However, extreme climate-related events (such as severe drought and extensive bushfire) were detected by the chosen bioindicators at both reference and test sites. Ecological outcomes of climate change can be accounted for only by an appropriate study design that includes standardized sampling of fixed sites (both test and reference) over long periods.

**Key words:** long-term trends, reference condition, macroinvertebrates, bioassessment indicators, climate change.

<sup>1</sup> E-mail addresses: sue.nichols@canberra.edu.au

<sup>2</sup> wrobinson@csu.edu.au

<sup>3</sup> richard.norris@canberra.edu.au

Australian alpine rivers form only a small percentage of Australian running waters nationally (ISC 2004), but they are significant for their water purity, aesthetic appeal, and their volumetric contributions to downstream river systems (DECC 2006). Like headwater streams worldwide, streams in the Australian highlands are vulnerable to effects of increasing temperatures because of climate change (Brown et al. 2007, Durance and Ormerod 2007, Füreder 2007). The global average for surface air temperature has increased  $\sim 0.74 \pm 0.18^{\circ}$ C in the past 100 y (IPCC 2007), and average global increases of 0.7 to 2.5°C are predicted by 2050 and 1.4 to 5.8°C by 2100 (Hennessy et al. 2003, Root et al. 2003, Pickering et al. 2004). Climate-change scenarios for Australian alpine areas predict that average annual temperatures will rise by another 0.4 to 2.0°C by 2030 and 1.0 to 6.0°C by 2070 (Hennessy et al. 2003). Snow patches in the Australian Alps already have declined (Green and Pickering 2009), and 2050 projections indicate possible reductions of 96% in the area sustaining snow cover for >60 d/y (Hennessy et al. 2003). The expected reduction in the extent of the Australian alpine zone poses an uncertain future for biodiversity (Pickering and Armstrong 2003, Pickering et al. 2004, Taylor and Figgis 2007) especially because the climate appears to be changing faster than predicted (Steffen 2009). Meteorological data for the study period (1994-2008) in New South Wales shows a strong increasing trend in the state's mean annual air temperature and erratic, but generally declining, annual rainfall (Chessman 2009). Changes in temperature, evaporation, and precipitation will determine the consequent changes in river flows over space and time (Chiew 2006), which, in turn, are likely to affect stream macroinvertebrates as indicators of ecological condition.

Most of Australia's alpine and subalpine areas are within national parks, but they are not free from human development, which threatens their ecological integrity. The Kosciuszko National Park (KNP) is one of the largest conservation reserves in Australia and has the highest peak on the mainland, Mount Kosciuszko (2228 m asl). Development activities within the KNP are associated with hydro-electricity production and ski resorts (DECC 2006). The KNP management plan requires all proposed developments and activities to demonstrate that they can maintain or improve natural ecological, hydrologic, and geomorphologic conditions and processes of rivers and streams within the park (DECC 2006). To evaluate ecological condition, KNP management has undertaken standardized biological assessment of fixed sites (to ensure uniformity in data collection and to detect trends) quarterly since 1994 using AUStralian RIVers Assessment System (AUSRIVAS) biological assessment methods (Simpson and Norris 2000). The AUSRIVAS technique uses the referencecondition approach (Reynoldson et al. 1997, Bailey et al. 1998) and benthic macroinvertebrate data to assess stream condition. AUSRIVAS and most other biological assessment methods rely on ecological benchmarks to provide the context for assessments (Hawkins et al. 2010). The KNP stream assessments provide park managers with feedback regarding a stream's response to management actions, so managers can make informed decisions to protect the park. Thus, managers must have confidence in the conclusions drawn from the biological assessments.

The predictive modeling approach to bioassessment includes methods like AUSRIVAS (Simpson and Norris 2000), the River InVertebrate Prediction and Classification System (RIVPACS) (Wright et al. 1998, Moss et al. 1999), and similar methods (Hawkins et al. 2000, Feio et al. 2009). The approach relies on predictions based on macroinvertebrate data collected from numerous reference sites sampled over a restricted period. The predictive models provide the observed/expected (O/E) taxa scores to make sitespecific assessments of stream reaches of unknown condition. Macroinvertebrates are collected (observed) and compared to the assemblage predicted (expected), given the values of a characteristic set of the environmental predictor variables (Wright 1995, Simpson and Norris 2000). The performance of stream assessments (predicting biota) is critically linked to how well the freshwater environment is characterized (Hawkins et al. 2010). However, potential exists for the benchmarks that define the reference conditions to change through time in response to climate-related environmental change (Robinson et al. 2000). If environmental conditions change through time, the macroinvertebrate assemblages in streams (including those at reference sites) also could change (Mazor et al. 2009). In a changing environment, managers will need to be aware of climate-related effects on O/E biological indicators to continue to make valid assessments of stream condition. If predictive models are not updated and calibrated through time with resampled reference-site data, potential exists for test sites to be compared to an outdated reference condition.

Long-term biological data sets are particularly valuable for assessing ecological responses to environmental change (Jackson and Füreder 2006, Mazor et al. 2009). Most studies that use biological techniques to assess stream health have not sampled the same sites through different seasons or over many years (Jackson and Füreder 2006). We used a 15-year data set from the KNP where a subset of the reference sites originally used to create the predictive model and test sites potentially affected by ski-resort activities were sampled quarterly. Fifteen years might not be considered long-term in ecological studies, but it is an extensive data set in terms of bioassessment



FIG. 1. Locations of test and reference study sites in Kosciuszko National Park, Australia.

(Jackson and Füreder 2006). This data set provided an opportunity to test whether the reference condition has changed during the 15-y period of record.

The macroinvertebrate biological indicators of stream condition used in our study were AUSRIVAS O/E taxa, Stream Invertebrate Grade Number Average Level (SIGNAL) 2 scores (Chessman 2003), the ratio of Ephemeroptera, Plecoptera, and Trichoptera (EPT) families to all families in the sample, Simpson's Diversity Index, and Oligochaeta abundance. We analyzed trends in climate and macroinvertebrate indicators of stream condition over 15 y to address the following questions: 1) What are the changes in climate in KNP during the last 15 y? 2) What is the relationship between changes in climate variables and macroinvertebrate biological indicators in KNP over the study period? 3) Are there differences in macroinvertebrate bioindicators between test and reference sites, and which bioindicators are consistent between seasons, years, and streams?

#### Methods

#### Study area

The climate of KNP (Fig. 1) is characterized by mean temperatures that range from  $-6^{\circ}$ C in July to 21°C in January. The region experiences 4 distinct seasons. Average annual rainfall recorded at the Perisher Valley gauging station within KNP is 1828 mm. Most streams in the Australian Alps are chemically similar to streams in European and North American alpine regions, with high concentrations of

dissolved  $O_2$  and low concentrations of nutrients and dissolved salts. They often have torrential flows during snow melt (Tiller 1993), but they carry low sediment loads compared to many northern-hemisphere high-altitude streams (Good 1992). Above 1500 m, frosts and blizzards can occur at any time, and snow usually accumulates between June and September. The snow line generally extends down to 1200 m. Almost continual melting of snow under the snow pack occurs during winter, and the runoff can lead to substantial transport of materials to streams (Molles and Gosz 1980). However, no snow remains in the study area throughout the year.

Developments, such as the Snowy Mountains Hydro-electric Scheme, have affected the hydrologic, geomorphologic, and ecological condition of many park streams (DECC 2006). Ski resorts are another major type of development within the park that can damage streams via discharge of treated sewage effluent, development activities, parking lots and associated runoff of road-deicer and salt-laden sediment, and abstraction of water for domestic supply and artificial snow making (Cullen et al. 1992, Digance and Norris 2006). In January 2003, bushfires burned 71% of the park's 673,492 ha, including the areas surrounding many of the study sites (DECC 2008).

Sixteen sites (11 test and 5 reference) were sampled on 5 streams. Test sites were positioned strategically to assess ecological responses to resort activities (Fig. 1). The sites range in elevation from 1160 m asl on Sawpit Creek (site 162) to 1760 m asl on Spencer's Creek (105). Extensive snow cover often prevents sampling at these high-altitude sites in winter. The most upstream site on all streams (11, 105, 121, 128, and 160) are minimally disturbed sites. These 5 sites are a subset of the many reference sites originally sampled to collect data to build the KNP predictive model used in our study. We resampled these 5 reference sites to provide a baseline from which to identify any regional shift from reference condition as indicated by the macroinvertebrate assemblage.

#### Macroinvertebrate sampling and bioindicators

Macroinvertebrates were collected quarterly in February (summer), May (autumn), August (winter), and November (spring) (quarters 1-4, respectively) each year from autumn 1994 to autumn 2008. Samples were collected from riffle habitats at each site using standard AUSRIVAS sampling techniques (Parsons and Norris 1996, Nichols et al. 2006). Each collection was subsampled to collect ~200 animals/sample (Marchant 1989). All sites (test and reference) sampled during the study period were assessed by the *Thredbo* predictive model, which was developed specifically for biological assessment of the streams in the KNP region based on AUSRIVAS modeling approaches (Simpson and Norris 2000), which are similar to those described for RIVPACS (Wright 1995, Wright et al. 1998, Moss et al. 1999). The AUSRIVAS macroinvertebrate collections and analysis provided the observed/expected (O/E) taxa scores. To aid interpretation of model outputs, AUSRIVAS O/E taxa scores are allocated to different bands that represent levels of biological condition. Bandwidths are based on the distribution of O/E taxa scores for the reference sites in any particular model. Band A is centered on 1.0 and includes the middle 80% of the reference-site O/E taxa scores. For the Thredbo model, band A is between O/E taxa scores 0.87 and 1.14. Condition of test sites with an O/E taxa score within Band A is considered equivalent to reference condition. A test site with an O/E taxa score below the lower bound (<0.87 for the Thredbo model) is judged to be significantly impaired with fewer families than expected and is allocated to one of the lower bands according to its score (Coysh et al. 2000).

We calculated 4 other bioindicators for each macroinvertebrate collection: the SIGNAL score for the sample, which is scaled from 1 (presence of many tolerant taxa) to 10 (presence of many sensitive animals) (Chessman 2003), the ratio of EPT families to all families in the sample, Simpson's Diversity, and the number of Oligochaeta in the sample ( $\log_{10}[x]$ -transformed). We selected these bioindicators because

they represent a variety of commonly used bioindicators, including those based on richness, composition, abundance, and sensitivity.

#### Bureau of Meteorology data

The Australian Bureau of Meteorology supplied us with climate data up to 2008 for weather stations at Thredbo and Perisher ski resorts. Records at Perisher began in June 1976 and Thredbo in January 1969. We selected climate variables for their potential effects on flow and temperature, which in turn, can influence temporal variation (interannual and seasonal) in macroinvertebrate assemblages (Wade et al. 1989, Linke et al. 1999, Mazor et al. 2009). We used highest monthly temperature (°C), lowest monthly temperature (°C), mean 3-PM relative humidity (%), and number of days with rain (full descriptions of variables are available from: http://www.bom.gov. au/climate/cdo/about/definitions9and3.shtml). We converted all climate data used in the analysis to quarterly series with the discontinuous piecewise constant curve method, which fits the average value (of the monthly data) to each quarterly interval and applies the most recent value to missing data points (SAS, version 9.1.3; SAS Institute, Cary, North Carolina). The meteorology variables from Perisher and Thredbo were highly correlated (all correlation coefficients >0.9). Therefore, we used the average value for the 2 weather stations when relating climate data to macroinvertebrate data. Only data from 1994 onward were included in the analysis to maintain correspondence between the 2 data sets.

#### Statistical analysis

We expected seasonal differences in climate data and macroinvertebrate assemblages because sites were sampled quarterly (Bêche et al. 2006). Preliminary analysis confirmed that macroinvertebrate time series and climate variables both showed a significant seasonal effect. Therefore, we ran all macroinvertebrate analyses separately for each season. We seasonally adjusted the time-series data for each climate variable with the US Bureau of the Census X-11 Seasonal Adjustment program by the standard X-11 method. The X-11 census method uses a ratio to moving-average technique (Shiskin et al. 1967, Ladiray and Quenneville 2001). We estimated the 15-y trends for each series with simple linear regression, including analysis of residuals, on the seasonally adjusted data. The residual analyses included visual confirmation that the assumptions of normality, homoscedasticity, and independence of residuals were not violated for any series. Many authors use

the Mann–Kendall nonparametric approach to stream data of this type, but the parametric approach we used is as powerful as the Mann–Kendall approach when the data are not skewed (Onoz and Bayazit 2003) (confirmed by the residual analyses) and is more easily interpreted.

We determined relationships between quarterly biological indicators and climate variables with Spearman's rank correlation coefficient (Mazor et al. 2009). We were unable to model the potential autocorrelation between consecutive years in each season adequately because of small sample sizes. Therefore, we interpreted *p*-values conservatively and applied a Benjamini– Hochberg procedure (Benjamini and Hochberg 1995) to control the false discovery rate (FDR), rather than family-wise error rate, for multiple tests.

We analyzed the differences in bioindicators between test and reference sites among years and streams with mixed models, which we chose because they use maximum-likelihood estimates rather than least-squares estimates and are superior to traditional linear models when analyzing repeated-measures data with missing values (which was occasionally the case for macroinvertebrate data from winter). Preliminary models included season as a factor, but model complexity and interaction effects involving season and the other variables made interpretation difficult. Therefore, we fitted a separate model for each season. Models included stream (5 levels), year (15 levels), and site type (test or reference) as fixed factors with sites subjected to repeated measures. We included all interaction terms in the model and used residual analysis to assess the assumptions of normality and constant variances before interpreting the outputs for each model. We present all results, but the main tests of interest were the interactions between year and stream or site type because these effects would indicate the potential for confounding by climate-induced changes. We controlled the FDRs for the tests with the Benjamini-Hochberg method (Waite and Campbell 2006) and interpreted significant effects by comparing least squares means where they could be estimated.

#### Results

#### Climate trends and correlation with macroinvertebrate data

Highest monthly temperature, lowest monthly temperature, and number of days with rain have increased significantly at KNP weather stations since 1994. Mean 3-PM relative humidity has decreased significantly in the same period (Table 1). These changes were all highly significant (p < 0.0001) and, therefore, cannot be dismissed as chance effects or

TABLE 1. Annual changes between 1994 and 2008 and significance of regression (*p*) after seasonal adjustment for 5 climate-related variables in Kosciuszko National Park, Australia.

Climate variable	Mean annual change	р
Highest monthly temperature (°C)	0.004	< 0.0001
Lowest monthly temperature $(^{\circ}C)$	0.002	< 0.0001
Mean 3-PM relative	-0.011	< 0.0001
Number of days with rain	0.009	< 0.0001

Type I errors even though the magnitude of change is small. Some climate variables and macroinvertebrate bioindicators were correlated before correcting for false discovery (Table 2), but the relationships were inconsistent. Some climate variables were both positively and negatively correlated with bioindicator values (depending on the site) (Table 2). For example, EPT richness ratio and Oligochaeta abundance were strongly related to climate variables at 7 sites. None of these relationships were significant after Benjamini– Hochberg correction.

### Macroinvertebrate indicators and effects of site type, stream, and year

*Reference and test sites.*—Mean scores for all macroinvertebrate indicators differed significantly (p < 0.05) between test and reference sites in all seasons, except for EPT ratios in autumn (Table 3). Differences between test and reference sites remained significant after Benjamini–Hochberg correction (Fig. 2A–E, Table 3, 4). Mean scores for all bioindicators were lower at test than at reference sites, except for Oligochaeta abundance, which was generally greater at test sites. Mean O/E taxa scores for resampled reference sites were assessed as similar to reference condition (AUSRIVAS Band A) (Fig. 2A–C), except for the mean O/E taxa score in spring 2007, when it dipped below Band A (Fig. 3C).

Stream differences.—All bioindicator scores, except for O/E taxa, differed significantly among streams in all seasons (Table 3), and most bioindicator scores were stream-specific. EPT ratios were highest at Thredbo River and lowest at Sawpit Creek, except in spring (Fig. 2D). Oligochaeta abundance was lowest at Sawpit Creek (Fig. 2E) and always was higher at Perisher Creek than at Sawpit or Thredbo, but the difference depended on the year (Table 3). Differences between streams generally did not depend on whether sites were reference or test sites (Table 3),

				adapate (pro	min Jointit (nie				
Site	Stream	Site type	Variable	Season	EPT richness ratio	O/E taxa	Oligochaeta abundance	SIGNAL	Simpson's Diversity
106	Spencer's	Test	Mean 3-PM temperature	4				0.66	
107 122	Spencer's Perisher	Test Test		-1 m	-0.66 0.68				
106	Spencer's	Test	Mean 3-PM relative humiditv	5	0		0.82		
122	Perisher	Test		-					0.66
123	Perisher	Test		ი, -		-0.67			
124	Perisher	Doference		4 0	0.70	0 80			
121 121	r iper s Perisher	Reference		с <del>–</del>		-0.09			0.66
161	Sawpit	Test		С			-0.70		
123	Perisher	Test	Highest monthly temperature	Ц	-0.76				
178	Piner's	Reference		c	-0.72				
129	Piper's	Test		14	1		0.71		
160	Sawpit	Reference		2	-0.75				
12	Thredbo	Test		1				-0.66	
105	Spencer's	Reference	Lowest monthly temperature	7	-0.65		0.68		
ر ر ر	Donichon	T	Amindum	V		040			
124 124	r erisher Perisher	Test		4 0		07.0	0.71		-0.68
160	Sawpit	Reference		10	-0.79		1		
162	Sawpit	Test							0.74
12	Thredbo	Test		7			0.68		
13	Thredbo	Test		2			0.68		
14	Thredbo	Test		0	-0.74				
11	Thredbo	Reference Toot		ლ <del>-</del>				0.68	
122	l'erisner	l est	number of days with rain	-				0.00	
122	Perisher	Test		ŝ		!			0.69
12	Thredbo	Test		4,		0.67			
14	Thredbo	Test		4,		0.71			
161 161	Sawpit Sawpit	Test		4 -			0.72	0.71	
	J			1					

TABLE 2. Significant (p < 0.05) Spearman rank correlation coefficients between bioindicators based on macroinvertebrates collected from sites in Kosciuszko National Park and climate variables from nearby weather stations before Benjamini-Hochberg correction (after which no correlations remained significant).

Downloaded From: https://complete.bioone.org/journals/Journal-of-the-North-American-Benthological-Society on 30 Apr 2024 Terms of Use: https://complete.bioone.org/terms-of-use

TABLE 3. Probabilities associated with mixed-model tests for differences in values of bioindicators among site types (test vs reference), years (1994–2008), and streams for different seasons (1 = summer, 2 = autumn, 3 = winter, 4 = spring). Bold indicates *p*-values were significant after controlling the false discovery rate using Benjamini–Hochberg correction. O/E = Australian Rivers Assessment System observed/expected taxa scores, SIGNAL = Stream Invertebrate Grade Number Average Level 2 scores, EPT = Ephemeroptera, Plecoptera, Trichoptera.

	C	EPT richness		Oligochaeta	CICNIAL 2	Simpson's
Effect	Season	ratio	O/E taxa	abundance	SIGNAL 2	Diversity
Site type	1	0.00567	0.00439	0.00041	0.02008	0.00306
51	2	0.11990	0.00057	0.00509	0.02988	0.00104
	3	0.03636	0.00014	0.00147	0.00818	0.00743
	4	0.00024	0.00023	0.00011	0.03862	0.00166
Year	1	0.00821	0.00701	0.00001	0.00010	0.00865
	2	0.02739	0.86516	< 0.00001	0.00542	0.00310
	3	0.00038	0.00262	0.00580	0.01353	0.00001
	4	0.00175	0.00107	< 0.00001	0.00070	0.00094
Stream	1	0.00103	0.30925	0.00360	0.00846	0.00177
	2	0.00622	0.14630	0.00077	0.00126	0.00073
	3	0.00363	0.32761	0.00134	0.00014	0.00130
	4	0.00053	0.10647	0.00072	0.00032	0.00032
Year $\times$ type	1	0.14204	0.11799	0.18577	0.81467	0.67209
• •	2	0.12561	0.60210	0.51417	0.98002	0.16037
	3	0.46383	0.93406	0.20521	0.26057	0.00776
	4	0.71828	0.93714	0.14283	0.00664	0.85925
Year $ imes$ stream	1	0.19608	0.02963	0.01668	0.28800	0.07293
	2	0.76020	0.53155	0.01489	0.35797	0.01613
	3	0.38493	0.03174	0.00111	0.19793	0.04677
	4	0.32247	0.15691	< 0.00001	0.00173	0.01353
Stream $\times$ type	1	0.23692	0.00922	0.06203	0.10888	0.03902
	2	0.27715	0.20041	0.07419	0.05888	0.03515
	3	0.27809	0.05486	0.06377	0.01769	0.23001
	4	0.26479	0.02317	0.02581	0.00613	0.15732
Year $\times$ stream $\times$ type	1	0.35660	0.58063	0.04520	0.98639	0.95107
	2	0.60077	0.88371	0.63061	0.99587	0.67050
	3	0.69847	0.66235	0.79613	0.19858	0.00356
	4	0.68380	0.20456	0.46593	0.09942	0.12732

except that the magnitude of differences in SIGNAL scores in spring and Simpson's Diversity Index in winter depended on the stream. SIGNAL scores were lowest at Perisher Creek test sites (Fig. 2B). O/E taxa scores generally were higher at reference than at test sites, but Thredbo River test sites and Sawpit Creek test sites scored high in summer and spring.

Differences between years.—Bioindicator scores, except for O/E taxa in autumn, which appeared more stable through time compared with other seasons (Fig. 3B), differed significantly among years. However, except for SIGNAL scores in spring and Simpson's Diversity Index in winter, differences between years did not depend on whether sites were reference or test sites (Table 3). For all seasons, differences among years in Oligochaeta abundance were stream-specific, as were differences among years in Simpson's Diversity Index in autumn and SIGNAL in spring (Table 3).

A decreasing trend in O/E taxa scores was evident in summer and spring since 2005 (Fig. 3A, C). Mean O/E taxa scores were significantly lower in spring 2007 and 2008 than in some previous years (Fig. 3A–C), as was EPT richness ratio in winter and spring (Fig. 3L). In 2003, mean SIGNAL scores in all seasons were distinctly lower than in other years (Fig. 3D–F), as were Simpson's Diversity Index in summer and autumn (Fig. 3G–I) and O/E taxa scores at reference sites (Fig. 3A, B). Oligochaeta abundance was significantly higher in autumn 2003 than in other years (Fig. 3M, N).

One 3-way interaction (year  $\times$  stream  $\times$  site type) was significant for Simpson's Diversity Index in winter, but in general, few interactions were significant. Thus, mean differences between site types and among streams remained about the same across years (Table 3).

#### Discussion

#### Climate changes and correlations

Over the 15-y study period, the climate became slightly warmer and less humid, but with a slight



FIG. 2. Mean ( $\pm 1$  SE) values of Australian Rivers Assessment System observed/expected (O/E) taxa scores (A), Stream Invertebrate Grade Number Average Level (SIGNAL) 2 scores (B), Simpson's Diversity Index (C), ratio of Ephemeroptera, Plecoptera, Trichoptera (EPT) family

increase in the number of days with rain. The slight increase in the number of days with rain has not led to more rainfall and runoff for the region (BoM 2009). Our results indicated small but highly (statistically) significant changes in climate-related variables (Table 1), but the analysis did not conclusively establish a clear, consistent relationship between macroinvertebrate bioindicators and climate variables (Table 2). However, extreme climate-related events, such as the extensive bushfires of summer 2003, corresponded to changes in the bioindicator time series. Bushfires can release nutrients that are fixed in soils and plants with the result that nutrients are transported with sediments to nearby streams (Gresswell 1999). Such inputs to streams that are normally low in nutrients and sediment are likely to alter taxonomic composition. SIGNAL (Fig. 3D-F), Simpson's Diversity values at test and reference sites (Fig. 3G-I), and O/E taxa scores for reference sites (Fig. 3A–C), were lower after the fires, and Oligochaeta abundance increased in the summer and the following autumn (Fig. 3M–O).

Further severe climatic conditions occurred from May 2006 to December 2006 when most of Australia was strongly affected by El Niño. Large regions of southeastern Australia received the lowest rainfalls on record. These dry conditions, combined with little relief from the previous 2002–2003 El Niño, contributed to another extremely dry season in 2006–2007 that resulted in long-running bushfires in Victoria and New South Wales (http://www.bom.gov.au/ climate/enso/enlist/index.shtml). In the same period, poorer stream conditions were indicated by O/E taxa ratios (Fig. 3A–C), EPT richness ratios (Fig. 3J–L), and SIGNAL scores (Fig. 3D–F). Thus, bioindicators detected the effects of extreme climate-related events.

#### Reference condition

*←* 

Climate change can trigger major environmental changes in aquatic ecosystems, which, in turn, trigger shifts in assemblage composition and structure of benthic macroinvertebrates and the bioindicator values used to assess stream condition (Harper and

richness (D), and Oligochaeta abundance (E) by stream, site type (test vs reference), and season averaged for all years (1994–2008). For ease of interpretation, untransformed data were graphed. However, the analysis was based on least-square means. In panel A, the dotted line represents the lower limit of Band A (0.87–1.14), which represents the reference condition. Numbers on the *x*-axis indicate seasons (1 = summer, 2 = autumn, 3 = winter, 4 = spring).

TABLE 4. Mixed-model analysis of variance table for Australian Rivers Assessment System observed/expected (O/E) taxa scores in spring as an example to show structure and degrees of freedom. \* indicates a significant *p*-value after correction for false discovery.

Effect	Numerator df	Denominator df	F	Probability of $> F$	Significance
Site type	1	6	60.77	0.0002	*
Year	14	81	2.97	0.0011	*
Year $\times$ type	14	81	0.48	0.9371	
Stream	4	6	3.07	0.1065	
Stream $\times$ type	4	6	6.44	0.0232	
Year $\times$ stream	55	81	1.28	0.1569	
Year $\times$ stream $\times$ type	55	81	1.22	0.2046	



FIG. 3. Least square mean values of Australian Rivers Assessment System observed/expected (O/E) taxa scores (A, B, C), Stream Invertebrate Grade Number Average Level (SIGNAL) 2 scores (D, E, F), Simpson's Diversity Index (G, H, I), ratio of Ephemeroptera, Plecoptera, Trichoptera (EPT) family richness (J, K, L), and log<sub>10</sub>Oligochaeta abundance (M, N, O) in summer (A, D, G, J, M), autumn (B, E, H, K, N), and spring (C, F, I, L, O) at test and reference sites averaged across streams in Kosciuszko National Park, Australia, from 1994 to 2008. In panels A, B, and C, the dotted line represents the lower limit of AUSRIVAS Band A (0.87–1.14), which represents the reference condition.

Peckarsky 2006, Durance and Ormerod 2007). Understanding whether bioindicators are changing in response to climate change is particularly important where the reference-condition approach (Reynoldson et al. 1997, Bailey et al. 1998) is used to assess sites at risk from human activities. Also important are robust bioindicators that can detect effects resulting from management-generated changes designed to enhance the condition of degraded streams. All bioindicators differed significantly between test and reference sites in all seasons (Table 3). The bioindicator values showed that, on average, stream conditions at test sites were most degraded at Perisher and Piper's Creeks in winter and spring compared to other streams and seasons (Fig. 2A-E). The most likely causes of this pattern are the pressures of ski-resort use and the treated effluent discharged into streams (Cullen et al. 1992, Digance and Norris 2006).

The average values of each bioindicator differed among some years for some seasons regardless of testor reference-site status, but only O/E taxa scores at the resampled reference sites could be used to determine that the reference condition was still valid and had not changed consequent to climate-related changes. This determination was possible because a subset of the original reference sites used to define the reference condition of the predictive model had been sampled through time (Fig. 3A-C). This result suggests that, apart from extreme climate-related events such as drought and fire, the bioindicators tested here were not sufficiently sensitive to detect significant, but subtle, changes in climate during the 15-y study (but see Lawrence et al. 2010, Poff et al. 2010, Stamp et al. 2010 for more information on sensitive indicators). However, we think that clear shifts in these bioindicators are likely if the predicted changes to the frequency of extreme weather events (DECC 2008) and snow cover in the Australian Alps occur (96% reduction in the area covered by snow for >60 d by 2050).

#### Differences among streams

Macroinvertebrate communities are prone to temporal and spatial variation in natural and modified streams (Linke et al. 1999, Bêche and Resh 2007). Many factors, such as differences in substrate (Quinn and Hickey 1990), stream size and position in the catchment (Minshall et al. 1985), in-stream and riparian vegetation (Read and Barmuta 1999, Quinn et al. 2004), temperature (Jacobsen et al. 1997) and flow-related variables (Boulton et al. 1992, Scarsbrook 2002, Armitage 2006, Poff and Zimmerman 2010), contribute to variation in the distribution and abundance of benthic macroinvertebrates. All macroinvertebrate bioindicators tested, except O/E taxa scores, differed among streams in all seasons (Table 3). O/E taxa scores are inherently adjusted for individual site characteristics (Reynoldson et al. 1997, Simpson and Norris 2000). The O/E taxa value is derived by comparing the taxa observed at a site with those expected based on the taxonomic composition of macroinvertebrate assemblages from groups of reference sites (least disturbed by human activity). When making an assessment, the test site is allocated a probability of belonging to each group of reference sites based on the values of model-specific environmental characteristics (Wright 1995, Simpson and Norris 2000). The reference group that is most similar to the test site will contribute most to determining the expected assemblage at each test site. Thus, streams in KNP might be in similar condition overall, but have among-stream differences in macroinvertebrate assemblages that are related simply to the situation of the streams in the landscape and site-specific differences in macroinvertebrate habitat variables (Hawkins et al. 2010). Variation among streams in the EPT richness ratio and other richness-related indicators also can be caused by stressors that exclude sensitive taxa. The tolerant taxa that remain might differ by site or stream as a function of habitat-specific requirements (Bady et al. 2005, Pollard and Yuan 2006, Hawkins et al. 2010). Our results support the view that stressed systems may have macroinvertebrate assemblages that are unique to each stream.

## Concluding remark on bioindicators, study design, and climate change

The AUSRIVAS O/E taxa score (which uses the reference-condition approach and predictive modeling for bioassessment) was the only indicator in our study that was adjusted for specific characteristics of each stream, and, thus, was robust to stream differences while being sensitive to reference- and test-site variation. Furthermore, O/E taxa scores indicated that reference sites sampled through time have remained in a condition equivalent to the previously defined reference condition, and, thus, provide a valid comparison for current test sites of unknown condition. Other bioindicators (e.g., EPT richness ratio, SIGNAL, Simpson's Diversity Index) varied significantly among streams, a result indicating that care should be taken when making comparisons among sites in different streams based on indictors not adjusted for natural stream differences.

The effects of climate change can be seen in 2 main ways: 1) gradual changes in temperature, rainfall,

runoff, fogs, etc., and 2) changes in the frequency and severity of extreme events, such as drought and bushfires (Hughes 2003, Campbell 2008). The bioindicators used in our study were insensitive to gradual but significant climate changes that occurred over 15 y. Generally, reference and test sites did not respond differently to gradual climate changes and the differences between them were consistent over time. However, extreme climate-related events (such as severe drought and extensive bushfires) could be detected at both reference and test sites by the bioindicators. Appropriate study designs that include standardized sampling of fixed sites (both test and reference) over long periods are needed to enable detection of climate-change effects, regardless of whether they are gradual changes or extreme events.

#### Acknowledgements

We acknowledge the staff from the Freshwater Ecology Laboratory at the Institute for Applied Ecology, University of Canberra, for sampling and processing macroinvertebrate samples. This assessment program was undertaken for Kosciuszko Thredbo Pty Ltd., KNP management, and the Department of Environment, Climate Change, and Water.

#### Literature Cited

- ARMITAGE, P. D. 2006. Long-term faunal changes in a regulated and an unregulated stream — Cow Green thirty years on. River Research and Applications 22: 947–966.
- BADY, P., S. DOLÉDEC, C. FESL, S. GAYRAUD, M. BACCHI, AND F. SCHOLL. 2005. Use of invertebrate traits for the biomonitoring of European large rivers: the effects of sampling effort on genus richness and functional diversity. Freshwater Biology 50:159–173.
- BAILEY, R. C., M. G. KENNEDY, M. Z. DERVISH, AND R. M. TAYLOR. 1998. Biological assessment of freshwater ecosystems using a reference condition approach: comparing predicted and actual benthic invertebrate communities in Yukon streams. Freshwater Biology 39: 765–774.
- BÈCHE, L. A., E. P. MCELRAVY, AND V. H. RESH. 2006. Longterm seasonal variation in the biological traits of benthic-macroinvertebrates in two Mediterranean-climate streams in California, USA. Freshwater Biology 51: 56–75.
- BÈCHE, L. A., AND V. H. RESH. 2007. Short-term climatic trends affect the temporal variability of macroinvertebrates in California 'Mediterranean' streams. Freshwater Biology 52:2317–2339.
- BENJAMINI, Y., AND Y. HOCHBERG. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B: Statistical Methodology 57:289–300.

- BoM (BUREAU OF METEOROLOGY). 2009. Climate data. Commonwealth of Australia, Bureau of Meteorology, Darlinghurst, Australia. (Available from: http://www.bom. gov.au/climate/how/contacts.shtml)
- BOULTON, A. J., C. G. PETERSON, N. B. GRIMM, AND S. G. FISHER. 1992. Stability of an aquatic macroinvertebrate community in a multiyear hydrologic disturbance regime. Ecology 73:2192–2207.
- BROWN, L. E., D. M. HANNAH, AND A. M. MILNER. 2007. Vulnerability of alpine stream biodiversity to shrinking glaciers and snowpacks. Global Change Biology 13: 958–966.
- CAMPBELL, A. 2008. Managing Australian landscapes in a changing climate: a climate change primer for regional natural resource management bodies. Department of the Environment, Water, Heritage and the Arts, Canberra, Australia. (Available from: http://triplehelix. com.au/documents/ClimateChangeprimer\_002.pdf)
- CHESSMAN, B. C. 2003. New sensitivity grades for Australian river macroinvertebrates. Marine and Freshwater Research 54:95–103.
- CHESSMAN, B. C. 2009. Climatic changes and 13-year trends in stream macroinvertebrate assemblages in New South Wales, Australia. Global Change Biology 15:2791–2802.
- CHIEW, F. H. S. 2006. Estimation of rainfall elasticity of streamflow in Australia. Hydrological Sciences Journal – Journal Des Sciences Hydrologiques 51:613–625.
- COYSH, J. L., S. J. NICHOLS, J. C. SIMPSON, R. H. NORRIS, L. A. BARMUTA, B. C. CHESSMAN, AND P. BLACKMAN. 2000. AUStralian RIVer Assessment System (AUSRIVAS) National River Health Program predictive model manual. Cooperative Research Centre for Freshwater Ecology, University of Canberra, Canberra, Australia.
- CULLEN, P., R. H. NORRIS, AND L. THURTELL. 1992. Managing water quality in the Australian Alps. Pages 425–449 *in* P. Grenier and R. Good (editors). The Australian Alps: Revue de Geographie Alpine. University of Grenoble and Australian Alps Liaison Committee, France: Institut de Géographie Alpine, Université Joseph Fourier.
- DECC (DEPARTMENT OF ENVIRONMENT AND CLIMATE CHANGE). 2006. Kosciuszko National Park plan of management: 2007–2008 implementation report. Department of Environment and Climate Change, Parks and Wildlife Group, Jindabyne, Australia. (Available from: http:// www.environment.nsw.gov.au/resources/nature/08596. pdf)
- DECC (DEPARTMENT OF ENVIRONMENT AND CLIMATE CHANGE). 2008. Kosciuszko National Park fire management strategy 2008–2013. Department of Environment and Climate Change, Parks and Wildlife Group, Jindabyne, Australia. (Available from: http://www.environment. nsw.gov.au/firemanagement/KosciuszkoNpFMS.htm)
- DIGANCE, J., AND R. H. NORRIS. 2006. Environmental impacts of tourism in the Australian Alps: the Thredbo River Valley. Pacific Tourism Review 3:37–38.
- DURANCE, I., AND S. J. ORMEROD. 2007. Climate change effects on upland stream macroinvertebrates over a 25-year period. Global Change Biology 13:942–957.

- FEIO, M. J., R. H. NORRIS, M. A. S. GRAÇA, AND S. NICHOLS. 2009. Water quality assessment of Portuguese streams: regional or national predictive models? Ecological Indicators 9:791–806.
- FÜREDER, L. 2007. Life at the edge: habitat condition and bottom fauna of Alpine running waters. International Review of Hydrobiology 92:491–513.
- GOOD, R. B. 1992. Kosciusko heritage: the conservation significance of the Kosciusko National Park. National Parks and Wildlife Service of New South Wales, Hurstville, Australia. (Available from: http://www. shop.nsw.gov.au/pubdetails.jsp?publication=18)
- GREEN, K., AND C. M. PICKERING. 2009. The decline of snowpatches in the Snowy Mountains of Australia: importance of climate warming, variable snow, and wind. Arctic, Antarctic, and Alpine Research 41: 212–218.
- GRESSWELL, R. E. 1999. Fire and aquatic ecosystems in forested biomes of North America. Transactions of the American Fisheries Society 128:193–221.
- HARPER, M. P., AND B. L. PECKARSKY. 2006. Emergence cues of a mayfly in a high-altitude stream ecosystem: potential response to climate change. Ecological Applications 16: 612–621.
- HAWKINS, C. P., Y. CAO, AND B. ROPER. 2010. Method of predicting reference condition biota affects the performance and interpretation of ecological indices. Freshwater Biology 55:1066–1085.
- HAWKINS, C. P., R. H. NORRIS, J. N. HOGUE, AND J. W. FEMINELLA. 2000. Development and evaluation of predictive models for measuring the biological integrity of streams. Ecological Applications 10:1456–1477.
- HENNESSY, K., P. WHETTON, I. SMITH, J. BATHOLS, M. HUTCH-INSON, AND J. SHARPLES. 2003. The impact of climate change on snow conditions in mainland Australia. CSIRO Atmospheric Research, Aspendale, Australia. (Available from: http://www.arcc.vic.gov.au/ publicationsclimatechange.htm)
- HUGHES, L. 2003. Climate change and Australia: trends, projections and impacts. Austral Ecology 28:423–443.
- IPCC (INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE). 2007. Climate change 2007: the physical science basis. Contribution of working group I to the 4<sup>th</sup> assessment report of the Intergovernmental Panel on Climate Change S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller (editors). Cambridge University Press, Cambridge, UK.
- ISC (INDEPENDENT SCIENTIFIC COMMITTEE). 2004. Independent scientific committee: an assessment of Kosciuszko National Park values. New South Wales Parks and Wildlife Service, Sydney, Australia. (Available from: Canprint, PO Box 279, Fyshwick ACT 2906, Australia; Chapter 8 available online at http://www.environment. nsw.gov.au/resources/parks/ISCValuesOfKosciuszkoNPCh08. pdf)
- JACKSON, J. K., AND L. FÜREDER. 2006. Long-term studies of freshwater macroinvertebrates: a review of the frequency, duration and ecological significance. Freshwater Biology 51:591–603.

- JACOBSEN, D., R. SCHULTZ, AND A. ENCALADA. 1997. Structure and diversity of stream invertebrate assemblages: the influence of temperature with altitude and latitude. Freshwater Biology 38:247–261.
- LADIRAY, D., AND B. QUENNEVILLE. 2001. Seasonal adjustment with the X-11 Method. Springer-Verlag, New York.
- LAWRENCE, J. E., K. B. LUNDE, R. D. MAZOR, L. A. BÊCHE, E. P. MCELRAVY, AND V. H. RESH. 2010. Long-term benthic macroinvertebrate responses to climate change: implications for biological assessment in mediterraneanclimate streams. Journal of the North American Benthological Society 29:1424–1441.
- LINKE, S., R. C. BAILEY, AND J. SCHWINDT. 1999. Temporal variability of stream bioassessments using benthic macroinvertebrates. Freshwater Biology 42:575–584.
- MARCHANT, R. 1989. A subsampler for samples of benthic invertebrates. Bulletin of the Australian Society for Limnology 12:49–52.
- MAZOR, R. D., A. H. PURCELL, AND V. H. RESH. 2009. Longterm variability in bioassessments: a twenty-year study from two northern California streams. Environmental Management 43:1269–1286.
- MINSHALL, G. W., R. C. PETERSEN, AND C. F. NIMZ. 1985. Species richness in streams of different size from the same drainage basin. American Naturalist 125:16–38.
- MOLLES, M. C., AND J. R. GOSZ. 1980. Effects of a ski area on the water quality and invertebrates of a mountain stream. Water, Air, and Soil Pollution 14:187–205.
- Moss, D., J. F. WRIGHT, M. T. FURSE, AND R. T. CLARKE. 1999. A comparison of alternative techniques for prediction of the fauna of running-water sites in Great Britain. Freshwater Biology 41:167–181.
- NICHOLS, S. J., W. A. ROBINSON, AND R. H. NORRIS. 2006. Sample variability influences on the precision of predictive bioassessment. Hydrobiologia 572:215–233.
- ONOZ, B., AND M. BAYAZIT. 2003. The power of statistical tests for trend detection. Turkish Journal of Engineering and Environmental Sciences 27:247–251.
- PARSONS, M., AND R. H. NORRIS. 1996. The effect of habitatspecific sampling on biological assessment of water quality using a predictive model. Freshwater Biology 36: 419–434.
- PICKERING, C. M., AND T. ARMSTRONG. 2003. The potential impact of climate change on plant communities in the Kosciuszko alpine zone. Victorian Naturalist 120:15–24.
- PICKERING, C. M., R. GOOD, AND K. GREEN. 2004. Potential effects of global warming on the biota of the Australian Alps. Australian Greenhouse Office, Canberra, Australia. (Available from: http://books.google.com/ books?id=DrneQgAACAAJ&dq=Potential+effects+of+ global+warming+on+the+biota+of+the+Australian+Alps& hl=en&ei=D8ZsTLmUF4nZcbuelYkB&sa=X&oi=book\_ result&ct=result&resnum=1&ved=0CC0Q6AEwAA)
- POFF, N. L., M. I. PYNE, B. P. BLEDSOE, AND C. C. CUHACIYAN. 2010. Developing linkages between species traits and multi-scaled environmental variation to explore vulnerability of stream benthic communities to climate change. Journal of the North American Benthological Society 29: 1441–1458.

- POFF, N. L., AND J. K. H. ZIMMERMAN. 2010. Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows. Freshwater Biology 55:194–205.
- POLLARD, A. I., AND L. YUAN. 2006. Community response patterns: evaluating benthic invertebrate composition in metal-polluted streams. Ecological Applications 16: 645-655.
- QUINN, J. M., I. K. G. BOOTHROYD, AND B. J. SMITH. 2004. Riparian buffers mitigate effects of pine plantation logging on New Zealand streams 2. Invertebrate communities. Forest Ecology and Management 191: 129-146.
- QUINN, J. M., AND C. W. HICKEY. 1990. Magnitude of effects of substrate particle size, recent flooding, and catchment development on benthic invertebrates in 88 New Zealand rivers. New Zealand Journal of Marine and Freshwater Research 24:411-427.
- READ, M. G., AND L. A. BARMUTA. 1999. Comparisons of benthic communities adjacent to riparian native eucalypt and introduced willow vegetation. Freshwater Biology 42:359-374.
- REYNOLDSON, T. B., R. H. NORRIS, V. H. RESH, K. E. DAY, AND D. M. ROSENBERG. 1997. The reference condition: a comparison of multimetric and multivariate approaches to assess water-quality impairment using benthic macroinvertebrates. Journal of the North American Benthological Society 16:833-852.
- ROBINSON, C. T., G. W. MINSHALL, AND T. V. ROYER. 2000. Interannual patterns in macroinvertebrate communities of wilderness streams in Idaho, USA. Hydrobiologia 421: 187 - 198.
- ROOT, T. L., J. T. PRICE, K. R. HALL, S. H. SCHNEIDER, C. ROSENZWEIG, AND J. A. POUNDS. 2003. Fingerprints of global warming on wild animals and plants. Nature 421: 57-60.
- SCARSBROOK, M. R. 2002. Persistence and stability of lotic invertebrate communities in New Zealand. Freshwater Biology 47:417–431.
- SHISKIN, J., A. H. YOUNG, AND J. C. MUSGRAVE. 1967. The X-11 variant of the census method II seasonal adjustment program. Technical report 15. Bureau of the Census, US Department of Commerce, Washington, DC.
- SIMPSON, J. C., AND R. H. NORRIS. 2000. Biological assessment of river quality: development of AUSRIVAS models and

outputs. Pages 125-142 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). Assessing the biological quality of fresh waters: RIVPACS and other techniques. Freshwater Biological Association, Ambleside, UK.

- STAMP, J. D., A. HAMILTON, L. ZHENG, AND B. BIERWAGEN. 2010. Use of thermal preference metrics to examine state biomonitoring data for climate change effects. Journal of the North American Benthological Society 29:1410–1423.
- STEFFEN, W. 2009. Climate change 2009: faster change and more serious risks. Department of Climate Change, Canberra, Australia. (Available from: http://www. anu.edu.au/climatechange/content/news/report-climatechange-2009-faster-change-and-more-serious-risks/)
- TAYLOR, M., AND P. FIGGIS. 2007. Protected areas: buffering nature against climate change. World Wildlife Fund and International Union for Conservation of Nature World Commission on Protected Areas: 18-19 June 2007, Canberra. World Wildlife Fund, Sydney, Australia. (Available from: http://www.wwf.org.au/publications/ cc-report/)
- TILLER, D. 1993. A preliminary study of the effect of Victoria's alpine ski resorts on high mountain streams. Report no. SRS 90/010, State Government Victorian Environmental Protection Agency, Melbourne, Australia. (Available from: Victorian Environmental Protection Agency, GPO Box 4395 Melbourne, Victoria, Australia 3001 on request, call number: SF 628.509945 SCI2E)
- WADE, K. R., S. J. ORMEROD, AND A. S. GEE. 1989. Classification and ordination of macroinvertebrate assemblages to predict stream acidity in upland Wales. Hydrobiologia 171:59–78.
- WAITE, T. A., AND L. G. CAMPBELL. 2006. Controlling the false discovery rate and increasing statistical power in ecological studies. Ecoscience 13:439-442.
- WRIGHT, J. F. 1995. Development and use of a system for predicting the macroinvertebrate fauna in flowing waters. Australian Journal of Ecology 20:181-197.
- WRIGHT, J. F., M. T. FURSE, AND D. MOSS. 1998. River classification using invertebrates: RIVPACS applications. Aquatic Conservation: Marine and Freshwater Ecosystems 8:617-631.

Received: 30 November 2009 Accepted: 16 August 2010

2010]