

## **Characterization of a Highly Repeated DNA Family in Tapetinae Species (Mollusca Bivalvia: Veneridae)**

Authors: Passamonti, Marco, Mantovani, Barbara, and Scali, Valerio

Source: Zoological Science, 15(4) : 599-605

Published By: Zoological Society of Japan

URL: [https://doi.org/10.2108/0289-0003\(1998\)15\[599:COAHRD\]2.0.CO;2](https://doi.org/10.2108/0289-0003(1998)15[599:COAHRD]2.0.CO;2)

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Characterization of a Highly Repeated DNA Family in Tapetinae Species (Mollusca Bivalvia: Veneridae)

Marco Passamonti\*, Barbara Mantovani and Valerio Scali

*Dipartimento di Biologia Evoluzionistica Sperimentale, Università di Bologna,  
via Selmi 3; I-40126 Bologna, Italy*

**ABSTRACT**—A repetitive DNA family (*phBgll400*) was characterized in the clam species *Tapes philippinarum* (Veneridae Tapetinae). The tandemly repeated sequences are AT-rich and show a mainly pericentromeric localization, as most satellite DNAs do. Sequence analysis of *phBgll400* DNA family showed a high level of intraspecific homogeneity. Furthermore a 200 bp subunit motif within the 400 bp monomer was apparent as well as the existence of two main “open reading frames” along the 400 bp sequence.

In order to investigate the possible distribution of this DNA family among Veneridae, Southern blot analyses were performed on genomic DNAs of *Tapes decussatus*, *Venerupis aurea* and *Paphia undulata* (Tapetinae), *Callista chione* (Pitarinae), *Chamelea gallina* (Chioninae) and *Venus verrucosa* (Venerinae). The *phBgll400* family has been found in two additional Tapetinae, namely *V. aurea* and *P. undulata*, but not in *T. decussatus* or other analyzed species. This pattern of sat-DNA distribution supports the high level of differentiation of *T. decussatus* observed in the previous gene-allozyme analysis. All of these suggest a better allocation of *T. decussatus* to a genus different from that of *T. philippinarum*.

## INTRODUCTION

*Tapes philippinarum*, *Tapes decussatus*, *Venerupis aurea* and *Paphia undulata* (subfamily Tapetinae, Fischer, 1887) represent some of the economically most important bivalves. Their taxonomic position and phyletic relationships are questioned. On the basis of shell morphology, “*decussatus*” and “*philippinarum*” taxa have been invariably considered congeneric and ascribed to *Tapes* Von Muehlfeldt, 1811; “*aurea*” has been most often ascribed to *Venerupis* Lamarck, 1818 (Fischer-Piette and Metivier, 1971), but, recently, also to *Paphia* Roding, 1798 (Sabelli *et al.*, 1990). Finally, “*undulata*” has been always ascribed to *Paphia*. On the other hand, morphological data sharply contrast allozyme and mitochondrial DNA analyses (Borsa and Thiriot-Quévieux, 1990; Canapa *et al.*, 1996; Passamonti *et al.*, 1997). In particular, biochemical and mtDNA findings do not support the assignment of *T. decussatus* and *T. philippinarum* to the same genus.

A useful tool to a further understanding of the phyletic relationships within the genus *Tapes* and the Tapetinae subfamily, could be a molecular analysis of tandemly repeated DNA sequences in their genomes. Eukaryotic genomes embody large amounts of tandemly repeated, non-coding DNA sequences, so-called satellite DNA (John and Miklos, 1979). Sat-DNA is highly variable in complexity and copy number; the repeated units can range from 3, 4 to thousand base pairs

and may represent from a few units to more than 50% of the genomic complement and can be arranged into either short interspersed or long tandemly repeated units (Singer, 1982; Miklos, 1985).

The general significance of satellite DNA is rather controversial and several hypotheses have been developed to assign this fraction of the eukaryote genome a role. It has been involved in heterochromatin constitution (Gershenson, 1933, 1940; Pardue, 1975; Brutlag, 1980), chromosome pairing (Salser *et al.*, 1976; Fry and Salser, 1977), rearrangements (Hatch *et al.*, 1976), tridimensional organization of the interphasic nucleus (Hilliker *et al.*, 1980; Manuelidis, 1982), gene amplification events (Bostock and Clark, 1980) and chromosome - mitotic spindle interactions through a peculiar class of centromeric sat-DNA-binding proteins (CENP), that preferably interact with DNA curvatures caused by d(A·T)<sub>n≥5</sub> stretches (Masumoto *et al.*, 1989). Alternative hypotheses suggest that sat-DNA lacks any function except its own survival (Doolittle and Sapienza, 1980; Orgel and Crick, 1980). Since it often corresponds to a heterochromatic noncoding genomic fraction, some models have been produced to explain its maintenance and evolutionary processes, such as “molecular drive” and “concerted evolution” (Dover, 1982, 1986, 1989; Charlesworth *et al.*, 1994).

Although a peculiar sat-DNA family is usually restricted to a single or a few closely related species, sometimes sat-DNAs show remarkable similarities in taxonomically distant taxa; highly conserved satellite DNAs can be spread over a whole group of phylogenetically related organisms (Arnason

\* Corresponding author: Tel. +39-51-354168;  
FAX. +39-51-251208.

*et al.*, 1984; Cremisi *et al.*, 1988) or even be widely distributed among members of evolutionarily distant groups (Abad *et al.*, 1992). It should be noted, however, that a higher level of homogeneity has been commonly found in intraspecific sequence analyses than in inter-specific comparisons. Therefore this approach could help when traditional investigations of taxonomy appear to be inadequate or even give controversial results at the specific level (Bachmann *et al.*, 1993).

Therefore we started a genomic DNA screening of *T. philippinarum*, *T. decussatus*, *V. aurea* and *P. undulata*, all belonging to the Tapetinae subfamily. A sat-DNA family was first detected in *T. philippinarum* and its distribution in the remaining species was analyzed. Comparisons were afterwards made also with genomic DNAs of *Callista chione* (subfamily Pitarinae), *Chamelea gallina* (subfamily Chioninae) and *Venus verrucosa* (subfamily Venerinae).

## MATERIALS AND METHODS

Nine individuals of seven Veneridae species from different localities were analyzed: two *T. philippinarum* (*phSc* - Scardovari Lagoon, Po river estuary; *phGa* - Ganzirri Lagoon, Sicily) and *T. decussatus* (*deSf* - Sfax, Tunisia; *deCh* - Chioggia, Venice Lagoon); one each of *V. aurea* (*auCi* - Civitanova Marche, Middle Adriatic Sea), *P. undulata* (*unTh* - Thailand), *C. chione* (*chCh* - Chioggia, Venice Lagoon), *C. gallina* (*gaCh* - Chioggia, Venice Lagoon) and *V. verrucosa* (*veCh* - Chioggia, Venice Lagoon).

Genomic DNAs were isolated by homogenizing a single foot-muscle in 160 mM sucrose, 80 mM EDTA and 100 mM Tris-HCl, pH 8 buffer and incubated at 65°C for 1 hr after addition of 0.5% SDS and 1 µg/ml of Proteinase K (Boehringer). Homogenates were then extracted for 3-5 times with phenol and/or chloroform. SDS separation was performed using potassium acetate at the final concentration of 1.2 M for 30' on ice and centrifuging for 15' at 15,000 rpm. The DNA was precipitated by the addition of 2 volumes of ethanol, centrifuging for 15' at 12,000 rpm. The DNA was recovered in TE 1 × buffer (10 mM Tris-HCl pH 8 and 1 mM EDTA) and treated with 50 µg/ml of RNase A (Boehringer) for 10' at room temperature. The DNA was finally precipitated with 0.2 M sodium acetate and 2 volumes of ethanol. From 50 to 200 µg of genomic DNA were obtained through this procedure.

Genomic DNAs were digested with 5 units of specific endonucleases/µg of DNA for 7-8 hr, according to the manufacturer's instructions. The following endonucleases were used: *Acc*, *AluI*, *Apal*, *AvaI*, *BamHI*, *BclI*, *BglII*, *Dde*, *DraI*, *EcoRI*, *HaeIII*, *HindII*, *HindIII*, *HpaI*, *MspI*, *NdeI*, *NsiI*, *PstI*, *RsaI*, *SacI*, *Sall*, *SmaI*, *StuI*, *TaqI*. All restricted DNAs were separated on 1.2% agarose gels in 1 × TBE buffer (90 mM Tris-borate; 2 mM EDTA) and then transferred onto a Nylon Membrane (Boehringer) following the protocol of Southern (1975).

From the *BglII* digested genomic DNAs of *T. philippinarum* (Scardovari and Ganzirri), a band, at about 400 bp, was cut and eluted from a 1.2% agarose gel using the Genomed JETsorb Gel Extraction Kit (cat. n. 110150). Eluted DNAs were cloned into the *BamHI* sites of pGEM -7Zf(+) (Promega) or pUC19 (+) (Pharmacia) plasmid vectors.

*V. aurea* and *P. undulata* monomers, only observed in Southern blots, could not be cloned directly, owing to their extremely low quantity. Therefore, a PCR amplification was performed with 5'-AGATTCCGCATGGC-3' and 5'-GACGTGTTCCCTGGG-3' primers, designed on the already obtained *T. philippinarum* sequence; these primers amplified the second part of the 400 bp sequence, since no good couple of primers could be designed to amplify the whole monomer. Primers were designed using PCR PLAN software in the PC/GENE 6.6 software package (© 1991, IntelliGenetics, 700 East El

Camino Real, Mountain View, California). The obtained fragments were therefore cloned into a pCR-Script plasmid vector with the Stratagene pCR-Script™ Amp SK(+) Cloning Kit (Cat. n. 211189).


Labeling and Southern blot hybridizations were performed according to the DIG-DNA kit instructions by Boehringer.

The nucleotide sequences of the recombinant clones were determined using the AutoRead™ Sequencing Kit (Pharmacia) with T7 and SP6 universal primers and a ALF™9000 automated sequencer (Pharmacia). Acrylamide gels were casted using ReadyMix Gels, ALF™ grade (Pharmacia).

Sequence alignments, consensus sequence, mutation distribution and codon analysis were performed using the Clustal algorithm and the Sequence Navigator software (Applied Biosystems). Genetic distances and cluster analyses were performed using the MEGA program (ver. 1.01, © S. Kumar, K. Tamura and M. Nei, 1993); in particular was utilized the "Kimura's two parameters" mutational model (Kimura, 1980) in computing distances.

Giemsa and NOR karyotypes and *in situ* hybridizations were performed on chromosome preparations from the gill tissue of *T. philippinarum* from Scardovari. Living clams were treated overnight with 0.005% colchicine in marine water. The dissected gills were hypotonically shocked with 1% w/v sodium citrate solution and then fixed with Carnoy's liquid (3:1, ethanol: acetic acid). Slides were prepared using a slightly modified air-drying technique by Crozier (1968).

Some slides were directly Giemsa stained, whereas others were treated with a silver impregnation technique, according to Marescalchi and Scali (1990), to evidence the active nucleolar organizing regions (NOR).



**Fig. 1.** Southern blot hybridization (at 1% stringency) of the *BglII* restricted DNAs of Tapetinae *T. philippinarum* from Scardovari (1), *T. decussatus* from Sfax (2), *P. undulata* from Thailand (3) and *V. aurea* from Civitanova (4). Approximate base-pair lengths are indicated.

**Table 1.** Alignment of the *pH*BglII400 DNA family cloned monomers of *T. philippinarum* from Scardovari (*pphSc*) and Ganzirri (*pphGa*). A consensus sequence, with *Bgl*II-like site (underlined) is shown. Asterisks (\*) indicate gaps introduced to increase sequence similarity. Last digits of numerals are aligned with the corresponding nucleotide.


	10	20	30	40	50	60	70	80	90	100
cons	AGATCTCCGC	ATGGCTAGGT	cACTTACCAG	CGATTTGCTT	TTTCTGTATA	TAAATTTCCGG	gAAcAAACAT	CAAtTTTACG	AGCTTGCCGT	TCAAAAAGCCA
<i>pphSc4</i>	-----	A-----G-----	-----G-----	-----A-----	-----	-----A-----	-----A-----	-----A-----	-----A-----	-----
<i>pphSc10</i>	-----	-----G-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphSc11</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphGa3</i>	-----	A-----G-----	-----*	-----	-----	-----	-----*	-----C-----	-----A-----	-----G-----
<i>pphGa4</i>	-----	-----G-----	-----C-----	-----	-----	-----	-----	-----C-----	-----A-----	-----
cons	TTTGACCAAT	TCCAATGCCA	ATTCGATTT	GTACAACTAG	AGAGGTCTT	CTTTGCACAT	GAAAAGGCCT	aAAGCCCTAA	CCCAAGTAAT	ATGTCCAGCA
<i>pphSc4</i>	-----	-----	-----	-----	T-----A-----	-----	-----	-----	-----	-----
<i>pphSc10</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphSc11</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphGa3</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphGa4</i>	-----	-----	-----	-----	-----C-----	-----G-----	-----	-----	-----A-----T-----	-----T-----
cons	GATTTCCGCA	TGGCTTGGTC	ATTTTTGCA	CGGGCATTT	GITTCCTTCT	CTATGTAAT	TTCGGGAAGG	AACATCATAT	TCACGAGCCT	ACCATTCAA
<i>pphSc4</i>	-----	-----C-----	-----	-----	*	G-----	-----A-----	-----	-----	-----
<i>pphSc10</i>	-----	-----	-----	-----	*	-----	-----	-----	-----	-----
<i>pphSc11</i>	-----	-----	-----	-----	G-----	-----	-----	-----	-----	-----
<i>pphGa3</i>	-----	-----	-----	-----	*	-----	-----	-----	-----	-----
<i>pphGa4</i>	A-----	-----T-----	-----	-----C-----	*	-----	-----A-----	-----	-----C-----	-----
cons	AACCAATTGA	TCAATTCCGT	TGCAAAATTG	ATTTGTACA	ACTTGAGGGG	GTCrKCTTTG	CACATgaita	AaGGCTTAAA	GCCCTAAACC	AAGGAACACG
<i>pphSc4</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphSc10</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphSc11</i>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>pphGa3</i>	-----	-----	-----GA-----	-----	-----	-----	-----	-----	-----	-----
<i>pphGa4</i>	-----	-----	-----	-----	TT-----	-----	-----*	-----	-----	-----G-----





**Table 5.** “Kimura two parameters” genetic distances between PCR-obtained 200 bp sequences of the *phBgII400* family in three Tapetinae species (*T. philippinarum*, *ph*; *V. aurea*, *au*; *P. undulata*, *un*).

	<i>pphSc</i> PCR1	<i>pphSc</i> PCR12	<i>pauCi</i> PCR5	<i>pauCi</i> PCR26	<i>punTh</i> PCR1	<i>punTh</i> PCR17
<i>pphSc</i> PCR1	–					
<i>pphSc</i> PCR12	0.000	–				
<i>pauCi</i> PCR5	0.010	0.010	–			
<i>pauCi</i> PCR26	0.036	0.036	0.036	–		
<i>punTh</i> PCR1	0.031	0.031	0.030	0.042	–	
<i>punTh</i> PCR17	0.026	0.026	0.025	0.042	0.031	–



**Fig. 2.** FISH labeled *in situ* hybridization of a *T. philippinarum* metaphase with a digoxigenin-labeled monomer obtained from the *pphSc11* recombinant clone.

tion of *T. decussatus* from the other Tapetinae taxa, which form a sound cluster. As a consequence, these data sets suggest that the two *Tapes* species could be separated to different genera.

While *V. aurea* is found only in the Mediterranean basin, *T. philippinarum* and *P. undulata* are Indo-Pacific clams in origin, their evolutionary divergence dating back, at least, to the closure of Tethys Ocean. Even if at present *T. philippinarum* and *V. aurea* have become artificially sympatric, the hypothesis of a recent “horizontal” transfer of these sequences appears very unlikely, since neither natural hybridization has been observed, nor artificial hybrids have been obtained. The assumption seems to be inconsistent with the observation that intra-specific sequence divergence in the 405 bp monomers of *T. philippinarum* is somewhat higher than the inter-specific one, as obtained from the PCR sequences. It is to be noted, however, that this comparison is heavily biased by the non-random working of the PCR-cloning procedure: likely, the observed distances are an outcome of the primer design performed on *T. philippinarum*. As a consequence, PCR amplified sequences of *V. aurea* and *P. undulata* would have artificially increased similarity to those of *T. philippinarum*.

Very little is known about satellite DNA in the Mollusca;

in the Bivalvia, only available for *Mytilus edulis* (Ruiz-Lara *et al.*, 1992). No definite hypotheses on the function of these sequences can be put forward. The high inter- and intra-specific sequence similarity of the *phBgII400* family may be attributed to the homogenization phenomena of the “concerted evolution”; these proceed through genomic turnover mechanisms (such as unequal crossing over, slippage replication and rolling circle amplification) and bisexual reproduction (Dover, 1982, 1986, 1989; Charlesworth *et al.*, 1994). Even the existence of some sort of selective constraints, such as satellite DNA-centromeric protein (CENP) interactions or hypothetical ORF transcription, could play a role in the maintenance of a high similarity of the *phBgII400* monomers.

## ACKNOWLEDGMENTS

We greatly appreciated the skilful assistance of Dr. Ettore Randi (Istituto Nazionale per la Fauna Selvatica, Ozzano Emilia, Bologna) in the PCR approach. This work was supported by M.U.R.S.T. 40% and 60% funds.

## REFERENCES

- Abad JP, Carmena M, Barrs S, Saunders RDC, Glover DM, Ludena P, Sentis C, Tyler-Smith C, Villasante A (1992) Dodecasatellite: a conserved G-C rich satellite from the centromeric heterochromatin of *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 89: 4663–4667
- Arnason U, Hoglund M, Widegren B (1984) Conservation of highly repeated DNA in cetaceans. *Chromosoma* 89: 238–242
- Bachmann L, Schibel JM, Raab M, Sperlich D (1993) Satellite DNA as a taxonomic marker. *Bioc Syst Ecol* 21: 3–11
- Borsa PE, Thiriou-Quievreux C (1990) Karyological and allozymatic characterization of *Ruditapes philippinarum*, *R. aureus* and *R. decussatus* (Bivalvia, Veneridae). *Aquaculture* 90: 209–227
- Bostock CJ, Clark EM (1980) Satellite DNA in large marker chromosomes of methotrexate-resistant mouse cells. *Cell* 19: 709–715
- Brutlag DL (1980) Molecular arrangement and evolution of heterochromatic DNA. *An Rev Genet* 14: 121–144
- Canapa A, Marota I, Rollo F, Olmo E (1996) Phylogenetic analysis of Veneridae (Bivalvia): comparison of molecular and palaeontological data. *J Mol Evol* 43: 517–522
- Charlesworth B, Sniegowski PE, Stephan W (1994) The evolutionary dynamics of repetitive DNA in eukaryotes. *Nature* 37: 215–220
- Cremisi F, Vignali R, Batistoni R, Barsacchi G (1988) Heterochromatic DNA in *Triturus* (Amphibia, Urodela). II. A centromeric satellite DNA. *Chromosoma* 97: 204–211

- Crozier RH (1968) An acetic acid dissociation, air-drying technique for insect chromosomes, with aceto-lactic orcein staining. *Stain Techn* 43: 171–173
- Doolittle FW, Sapienza C (1980) Selfish genes, the phenotype paradigm and genome evolution. *Nature* 299: 601–603
- Dover GA (1982) Molecular drive: a cohesive mode of species evolution. *Nature* 299: 111–117
- Dover GA (1986) Molecular drive in multigene families: how biological novelties arise, spread and are assimilated. *Trends Genet* 2: 159–165
- Dover GA (1989) Slips, strins and species. *Monitor* 4: 100–102
- Fischer-Piette E, Metivier B (1971) Révision des Tapetinae (Mollusques Bivalves). *Mem Mus Hist Natur Paris* 71: 1–106
- Fry K, Salser W (1977) Nucleotide sequences of HS- $\alpha$  DNA from kangaroo rat *Dipodomys ordii* and characterization of similar sequences in other rodents. *Cell* 12: 1069–1084
- Gershenson S (1933) Studies on the genetically inert region of the X-chromosome of *Drosophila*. I. Behavior of an X-chromosome deficient for a part of its insert region. *J Genet* 28: 297–313
- Gershenson S (1940) The nature of the so-called genetically inert parts of chromosomes. *Vid Akad Nauk USSR* 1940: 3–116
- Hatch FT, Bodner AJ, Mazrimas JA, Moore DH (1976) Satellite DNA and cytogenetic evolution. DNA quantity, satellite DNA and karyotypic variations in kangaroo rats (Genus *Dipodomys*). *Chromosoma* 58: 155–168
- Hilliker AJ, Appels R, Schalet A (1980) The genetic analysis of *D. melanogaster* heterochromatin. *Cell* 21: 607–619
- John BE, Miklos GLG (1979) Functional aspects of satellite DNA and heterochromatin. *Int Rev Cytol* 58: 1–114
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16: 111–120
- Maluszynska J, Heslop-Harrison JS (1993) Physical mapping of rDNA loci in *Brassica* species. *Genome* 36: 774–781
- Manuelidis L (1982) Repeated DNA sequences and nuclear structure. In "Genome Evolution" Ed by GA Dover and Flavell RB, Academic Press, London, UK, pp 263–285
- Marescalchi O, Scali V (1990) Cytogenetic studies on *Bacillus grandii grandii* and *Bacillus grandii benazzii*: karyotype description, constitutive heterochromatin and nucleolus organizer regions. *Genetica* 82: 117–124
- Masumoto H, Sugimoto K, Okazaki T (1989) Alphoid satellite DNA is tightly associated with centromere antigens in human chromosomes throughout the cell cycle. *Exp Cell Res* 181: 181–196
- Miklos GLG (1985) Localised highly repetitive DNA sequences in vertebrate and invertebrate genomes. In "Molecular Evolutionary Genetics" Ed by MacIntyre RJ, Plenum, New York, pp 241–321
- Orgel LE, Crick FHC (1980) Selfish DNA: the ultimate parasite. *Nature* 284: 604–607
- Pardue ML (1975) Repeated DNA sequences in the chromosomes of higher organisms. *Genetics* 79: 159–170
- Passamonti M, Mantovani B, Scali V (1997) Allozymic characterization and genetic relationships among four species of Tapetinae (Bivalvia Veneridae). *It J Zool* 64: 117–124
- Ruiz-Lara S, Prats E, Sainz J, Cornudella L (1992) Cloning and characterization of a highly conserved satellite DNA from the mollusc *Mytilus edulis*. *Gene* 117: 237–242
- Sabelli B, Giannuzzi-Savelli R, Bedulli D (1990) Catalogo Annotato dei Molluschi Marini del Mediterraneo. Libreria Naturalistica Bolognese, Bologna
- Salser W, Bowen S, Browne D, El Adli F, Fedoroff N, Fry K, Heindell H, Paddock G, Poon R, Wallace B, Whitcome P (1976) Investigation of the organization of mammalian chromosomes at the DNA sequence level. *Fed Proc* 35: 23–35
- Singer MF (1982) Highly repeated sequences in mammalian genomes. *Int Rev Cytol* 67: 67–112
- Southern EM (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 98: 503–517

(Received November 19, 1997 / Accepted April 15, 1998)