

Loss of Photoperiodic Control of Juvenile-Hormone Signaling Pathway Underlying the Evolution of Obligate Parthenogenesis in the Pea Aphid

Authors: Ishikawa, Asano, Gotoh, Hiroki, Ogawa, Kota, Kanbe, Takashi, Akimoto, Shin-ichi, et al.

Source: Zoological Science, 42(2): 186-195

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zs240075

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Loss of Photoperiodic Control of Juvenile-Hormone Signaling Pathway Underlying the Evolution of Obligate Parthenogenesis in the Pea Aphid

Asano Ishikawa^{1,2*}, Hiroki Gotoh^{1,3}, Kota Ogawa^{1,4,5}, Takashi Kanbe⁶, Shin-ichi Akimoto⁶, and Toru Miura^{1,7}

 ¹Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan
 ²Department of Biological Sciences, Graduate School of Sciences, The University of Tokyo, Kashiwa, Chiba 277-8562, Japan
 ³Department of Science, Graduate School of Integrated Science and Technology, Shizuoka University, Shizuoka, Shizuoka 422-8529, Japan
 ⁴Biosystematics Laboratory, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka, Fukuoka 819-0395, Japan
 ⁵Insect Science and Creative Entomology Center, Kyushu University, Fukuoka, Fukuoka 819-0395, Japan
 ⁶Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan
 ⁷Misaki Marine Biological Station, School of Science, The University of Tokyo, Miura, Tokyo 238-0225, Japan

Phenotypic plasticity, the ability of organisms to change their phenotype depending on external stimuli, enables their survival in fluctuating environments. An extreme example is polyphenism, in which a single genotype produces discrete phenotypes in response to external cues. However, under persistent environmental conditions, natural selection would favor reduced plasticity. This study focused on the loss of reproductive polyphenism and revealed the underlying mechanism in the pea aphid Acyrthosiphon pisum. Although most populations exhibit reproductive polyphenism, known as cyclical parthenogenesis, with a seasonal shift between parthenogenesis and sexual reproduction, some exhibit obligate parthenogenesis. To investigate the potential role of changes in the environmental sensitivity of the juvenile hormone (JH) pathway during this evolutionary shift, we analyzed the expression of genes involved in JH synthesis and degradation. We found that five of seven JH-related genes exhibited photoperiodic responses in one cyclicalparthenogenetic strain, whereas none of them responded to photoperiod in the two obligateparthenogenetic strains. Notably, CYP15A and JHEH genes, which are involved in the final step of JH synthesis and in the initiation of JH degradation, respectively, showed strong photoperiodic responses in the cyclical-parthenogenetic strain but showed no responses in the obligateparthenogenetic strains. Acetone treatment induces male production in obligate-parthenogenetic strains, suggesting that the developmental pathway for male production remains functional in these strains. These results suggest that the loss of the photoperiodic response in both JH synthesis and degradation pathways is a key mechanism underlying the elimination of the sexual phase, resulting in the loss of reproductive polyphenism in aphids.

Key words: phenotypic plasticity, cyclical parthenogenesis, obligate parthenogenesis, loss of polyphenism, pea aphid

INTRODUCTION

In phenotypic plasticity, organisms can modify their phenotype by altering their developmental processes to adapt to their surrounding environment (West-Eberhard, 2003). An extreme example is polyphenism, such as seasonal color morphs in butterflies, defensive morphs in Daphnia, and caste differentiation in social insects, in which organisms produce multiple discrete phenotypes, even from a single genotype, in response to different environmental conditions. The ability to change phenotypes in response to external stimuli enables organisms to attain multiple optimal states in fluctuating environments. However, under persistent envi-

^{*} Corresponding author. E-mail: ishikawa@k.u-tokyo.ac.jp doi:10.2108/zs240075

ronmental conditions, natural selection favors genetic changes that produce a constant phenotype. Theoretical and empirical studies have suggested that both the gain and loss of phenotypic plasticity play substantial roles in phenotypic diversification and speciation (Pigliucci, 2001; West-Eberhard, 2003; Aubret and Shine, 2009; Lande, 2009; Pfennig et al., 2010; Schwander and Leimar, 2011). However, few examples have been reported regarding the proximate mechanisms underlying a reduction in plasticity, especially in natural populations.

Notably, aphids (Insecta, Hemiptera) exhibit life-cycle polyphenism, including seasonal reproductive polyphenism and wing polyphenism, although phenotypic responses vary even within a single species (Moran, 1992; Dixon, 1997; Simon et al., 2002; Ogawa and Miura, 2014; Shuo et al., 2020). Most aphid species exhibit reproductive polyphenism, also known as cyclical parthenogenesis, which involves switching between parthenogenesis and sexual reproduction depending on the season (Dixon, 1997). During the spring and summer, aphids reproduce via viviparous parthenogenesis. In the fall, parthenogenetic females produce sexual morphs, males and oviparous females, that mate and produce overwintering eggs. In most aphids, short-day photoperiods and low temperatures are the major environmental cues that trigger the production of sexual morphs (Moran, 1992; Simon et al., 2002). Because all aphid species exhibit cyclical parthenogenesis, this strategy appears to have been acquired in the early stages of aphid evolution. However, approximately 30–40% of aphid species possess populations that reproduce exclusively by obligate parthenogenesis (Fig. 1A) (Moran, 1992; Dixon, 1997; Simon et al., 2002). These populations are thought to be insensitive to photoperiodic changes, which results in the loss of the sexual phase (Lees, 1959, 1963; Simon et al., 2010).

In many insects, physiological mechanisms, particularly those of the endocrine system, control phenotypic changes in response to environmental cues (Hartfelder and Emlen, 2012). Especially, juvenile hormone (JH) plays a key role in regulating development, metamorphosis, reproduction, diapause, and polyphenism in insects (Nijhout, 1994, 1999, 2003; Zera and Denno, 1997; Tanaka, 2001; Miura, 2005; Hartfelder and Emlen, 2012). In aphid cyclical parthenogenesis, JH has been suggested to trigger the production of sexual morphs in response to short photoperiods (Hales and Mittler, 1983; Corbitt and Hardie, 1985; Ishikawa et al., 2012). Therefore, we hypothesized that modifications in the photoperiodic response of the JH signaling pathway might be a key mechanism underlying the loss of the sexual phase. The functional JH of aphids is considered to be juvenile hormone III (JHIII) (Hardie et al., 1985; Westerlund and Hoffmann,



Fig. 1. The cyclical-parthenogenetic (CP) and the obligate-parthenogenetic (OP) populations of the pea aphid. P and S females indicate parthenogenetic and sexual females, respectively. **(A)** Annual life cycles of the CP and OP populations. The CP populations change their reproductive mode from parthenogenesis to sexual reproduction depending on the seasonal changes (upper). On the other hand, the OP populations have lost the sexual phase in their life cycle (lower). **(B)** The sampling points and reproductive schedules of the CP, OP1, and OP2 strains in Japan. The experimental categories are shown with the schedules. G1 and G2 indicate the first generation and second generation, respectively, after we started rearing the aphids under long-day or short-day conditions (see Fig. 3).

2004; Chen et al., 2007; Schwartzberg et al., 2008; Ishikawa et al., 2012, 2013). However, recent studies have revealed that many hemipteran species use JHIII skipped bisepoxide (JHSB₃) as a functional JH (Kotaki et al., 2009, 2012; Ando et al., 2020; Matsumoto et al., 2020; Villalobos-Sambucaro et al., 2020; Kodama et al., 2023). In aphids (Aphididae), both JHIII and JHSB3 were detected in the hemolymph of Myzus persicae and Rhopalosiphum maidis (Yi et al., 2023), whereas only JHSB3 was detected in Lipaphis erysimi (Yi et al., 2023). Thus, the types of chemical compounds that function as JH in aphids remain unclear. However, the synthesis pathways of JH III and JHSB3 and the genes encoding the enzymes that catalyze these reactions are largely conserved (Tsang et al., 2020; Mano and Goto, 2022). Therefore, we investigated the photoperiodic responses of the JH pathway by examining the expression profiles of the genes involved in JH synthesis and degradation.

Here, we compared the photoperiodic responses of genes involved in JH synthesis and degradation between cyclical-parthenogenetic (CP) and obligate-parthenogenetic (OP) strains of the pea aphid, Acyrthosiphon pisum (Fig. 1A). Intraspecific variations in reproductive polyphenism have been reported in pea aphids (Kanbe and Akimoto, 2009). The OP strains are distributed in central Japan, whereas the CP strains are distributed in northern Japan. Our previous research revealed that in the CP strain, aphids reared under short-day conditions showed enhanced expression of JH-degradation genes, such as JH esterase (JHE) and JH epoxide hydrolase (JHEH). This upregulation of the JH degradation pathway likely triggers the induction of sexual morphs (Ishikawa et al., 2012). In addition to the genes related to JH degradation, we analyzed the genes involved in JH biosynthetic pathways, such as JH acid methyltransferase (JHAMT), JH epoxidase (CYP15A1), and short-chain dehydrogenase (SDR) (see Fig. 2) (Shinoda and Itoyama, 2003; Mayoral et al., 2009; Daimon and Shinoda, 2013). Furthermore, we conducted an artificial male induction in obligate-parthenogenetic strains to test whether they retained the developmental cascade for producing sexual individuals that occurs downstream of the JH signaling pathway.

MATERIALS AND METHODS

Insects

To compare differences in photoperiodic responses between cyclical-parthenogenetic and obligate-parthenogenetic populations, one CP strain (ApL) and two OP strains, Hac06VsXI30 and Sait06Vs4 (hereafter called OP1 and OP2, respectively), of the pea aphid, Acyrthosiphon pisum, were used in this study. The CP strain was established from a cyclical-parthenogenetic population collected from Sapporo, Hokkaido, Japan (Ishikawa et al., 2012). The OP1 and OP2 strains were established from an obligate-parthenogenetic population collected from Hachinohe City, Aomori, Japan, in November 2006 and an obligate-parthenogenetic population collected from Saitama City, Saitama, Japan, in May 2006 (Fig. 1B). Based on microsatellite analysis, the OP1 and OP2 strains had MLG120 and MLG122 genotypes, respectively, which are common and widely distributed over the years in obligate-parthenogenetic populations (Kanbe and Akimoto, 2009). Thus, both strains were predicted not to produce males or sexual females under short-day conditions. Microsatellite analysis was performed as previously described (Kanbe and Akimoto, 2009).

Stock cultures of parthenogenetic aphids were maintained in

tubes (diameter: 2.5 cm, height: 10 cm) in which a bean seedling (*Vicia faba*) was placed on wet vermiculite at 20°C under long-day conditions (16L8D) (Wilkinson and Ishikawa, 2000). Only wingless parthenogenetic females were used for all experiments because winged parthenogenetic females did not produce males, even under short-day conditions (8L16D, 15° C) (Ishikawa et al., 2012). Since the effect of rearing conditions lasts over two or three generations (Mackay and Wellington, 1977), aphids were kept on a 3-cm vetch seedling over three generations under long-day conditions (16L8D, 15° C) as the pre-induction to prevent to production of winged parthenogenetic females (Ishikawa et al., 2008; Ogawa and Miura, 2014).

Reproductive schedule of obligate-parthenogenetic strains under short-day conditions

To confirm the obligate parthenogenesis of the OP1 and OP2 strains, the aphids were exposed to short-day conditions (8L16D) and low temperature (15°C), in the same manner as that used to induce the production of males and sexual females in the CP strain (Ishikawa et al., 2012). Briefly, first-instar nymphs produced by single wingless mothers of the OP1 and OP2 strains were isolated and reared on seedlings under short-day conditions. When the aphids reached the adult stage and began reproducing, they were transferred daily to a new seedling in another container. The number and morphs (parthenogenetic females, males, and sexual females) of the progeny produced each day were recorded. Because the morphs could not be identified in newborn nymphs, progenies were reared until they reached fifth-instar adults for accurate identification. In the adult stage, parthenogenetic females and oviparous sexual females were distinguished by the appearance of ovaries: parthenogenetic females have parthenogenetic ovaries with many embryos, whereas sexual females possess gametic ovaries containing only haploid eggs (Blackman, 1987). Males were discriminated based on their wings, small abdomens, and male gonads. To confirm asexuality over generations, we maintained the strains under short-day conditions for several generations.

Identification of orthologs for genes involved in JH synthesis and degradation

Although the ortholog searches were conducted in our previous studies (Ishikawa et al., 2012), the genome assembly of A. pisum in AphidBase was updated to version 2.1. Therefore, Basic Local Alignment Search Tool (BLAST) search was conducted for Acyrthosiphon pisum annotation ncbi 2.1 (https://bipaa.genouest. org/is/aphidbase/) to identify the pea aphid orthologs of SDR, JHAMT, CYP15A1, JHEH, and JHE (Fig. 2, Table 1) (Li et al., 2019). For SDR, we used the protein sequence of Aedes aegypti, AAEL017302, as the query sequence (Mayoral et al., 2009). For JHAMT, we used the protein sequence of Bombyx mori (AB113578) as the query. For the CYP15A1 gene, Daimon and Shinoda (2013) identified one ortholog, XP_001952620; therefore, we used it as a query. For the JHEH gene, we used the protein sequence of Manduca sexta, AAC47018, as a query (Wojtasek and Prestwich, 1996). For JHE, we use the protein sequence of Nilaparvata lugens, EU380769, as the query (Kamita and Hammock, 2010). Genes with more than 40% similarity were used as putative orthologs. To confirm whether the ortholog encodes a functional enzyme, we surveyed the critical residues known to characterize these enzymes in each ortholog.

Experimental designs for gene expression analysis

To test the hypothesis that the modification of the photoperiodic response of genes related to JH synthesis and degradation underlies the loss of reproductive polyphenism, we analyzed the expression of genes involved in JH synthesis and degradation in response to photoperiodic changes in the CP, OP1, and OP2 strains. As reported previously, CP strains produced sexual individuals (males and sexual females) under short-day conditions (Figs. 1B, 3A) (Ishikawa et al., 2012). After the first-instar nymphs were isolated at 15°C under short-day conditions (8L16D), the parthenogenetic females (generation 1, G1) produced only parthenogenetic progeny until 8 days after the first larviposition, and then started to produce male progeny in the late period from 10 days after the first larviposition. The parthenogenetic females (generation 2: G2) produced on the second day after the first larviposition by G1 females preferentially produced sexual females in the early period and then started to produce both males and parthenogenetic females in the late period. Under long-day conditions (16L8D), all parthenogenetic females (but no sexual individuals). Based on the reproductive schedule of the CP strain, gene expression was examined on the



Fig. 2. Abbreviated pathways of JH biosynthesis and degradation. Genes denoted by grey boxes are analyzed in this study.

first (day 1) and eighth (day 8) day after the first larviposition under long- and short-day conditions, resulting in the following four categories: LD-Day 1, LD-Day 8, SD-Day 1, and SD-Day 8. In the obligate-parthenogenetic OP1 and OP2 strains, all parthenogenetic females on LD-Day 1, LD-Day 8, SD-Day 1, and SD-Day 8 produced future parthenogenetic female embryos (Figs. 1B; 3B, C).

RNA extraction and quantification of gene expression involved in JH synthesis and degradation

Total RNA was extracted from whole bodies of five individuals stored at -80°C using the SV Total RNA Isolation System (Promega, Madison, WI, USA). Three biological replicates (five individuals per replicate) were prepared for each category. One microgram of total RNA from each sample was reverse-transcribed using a High Capacity cDNA Reverse Transcription Kit according to the manufacturer's instructions (Applied Biosystems, Foster City, CA, USA). To compare the photoperiodic response of gene expression involved in JH synthesis and degradation pathways among the three strains, we performed real-time gPCR for SDR1, SDR2, SDR3, SDR4, JHAMT, CYP15A1, and JHEH genes using the SYBR Green I chemistry system and sequence detection system ABI PRISM 7500 (Applied Biosystems), as described previously (Ishikawa et al., 2012). Data acquisition and analyses were performed using ABI Prism 7500 software v2.0.1 (Applied Biosystems). The baselines and thresholds for Ct were set automatically. The relative standard curve method was used for gRT-PCR quantification, as described in User Bulletin 2 for the ABI Prism 7700 Sequence Detection System (Applied Biosystems). Primers for both target and endogenous control transcripts were designed using Primer Express software (Applied Biosystems). EF1a (elongation factor 1 alpha) was used as an endogenous control, as described previously, and was determined using geNorm and NormFinder software for comparison with glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and beta-actin genes (Ishikawa et al., 2012). For JHAMT and JHEH, we used the same primer set as in our previous study (Ishikawa et al., 2012). The primer sequences are listed in Supplementary Table S1. Two-way ANOVA was performed using R. 3.0.3 software (R Core Team, 2014).

Acetone treatment

To investigate whether the obligate-parthenogenetic strains possess a developmental cascade to produce sexual individuals, we artificially induced male production in the OP strains. Although a previous study reported that treatment with precocene dissolved in acetone induces male production in the green peach aphid *Myzus persicae* (Hales and Mittler, 1983), our preliminary experiment indicated that acetone itself induced male production. Therefore, we artificially induced males by acetone treatment in the OP1

Gene symbol	Function	ACYPI ID	Reference nucleotide sequence	Reference protein sequence	
SDR1	JH synthesis		NM_001160394	NP_001153866	
SDR2	JH synthesis	ACYPI009545	XM_008179948, NM_001293482	XP_008178170, NP_001280411	
SDR3	JH synthesis	ACYPI001301	XM_008187359	XP_008185581	
SDR4	JH synthesis	ACYPI003207	NM_001162459	NP_001155931	
JHAMT1	JH synthesis	ACYPI007696	NM_001162779	NP_001156251	
JHAMT2	JH synthesis	ACYPI001588	XM_001943694	XP_001943729	
JHAMT3	JH synthesis		XM_003244251	XP_003244299	
JHAMT4	JH synthesis		XM_016802746	XP_016658235	
CYP15A1	JH synthesis	ACYPI003882	XM_001952585	XP_001952620	
JHEH	JH degradation		XM_003241505	XP_003241553	

Table 1. List of juvenile hormone-related genes analyzed in this study.

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 24 May 2025 Terms of Use: https://complete.bioone.org/terms-of-use



Days after first larviposition

Fig. 3. Reproductive schedules of the CP (**A**), OP1 (**B**), and OP2 (**C**) strains under short-day conditions. Mean number and standard errors of parthenogenetic females (triangles and pink dashed line), males (circles and solid light blue line), and oviparous sexual females (squares and dash-dotted dark blue line) produced by aphids in the first generation (G1) or second generation (G2) under short-days (CP [G1]: n = 8, CP [G2]: n = 12, OP1: n = 10, OP2: n = 10). Upper panels of the CP strain are modified from Ishikawa et al. (2012).

and OP2 strains. Approximately 0.5 μ L of acetone was applied to the abdomen of wingless parthenogenetic females just after the final ecdysis. As a control, we used 0.5 μ L of ethanol and pentane. The aphids were then placed individually on a seedling at 20°C under long-day conditions (16L8D) and were transferred every 2–3 days to a new seedling to confirm the production of males or sexual females in the next generations. Fisher's exact test was conducted using R. 3.0.3 software (R Core Team, 2014).

RESULTS

Confirmation of obligate parthenogenesis in OP1 and OP2 strains

Under short-day conditions, all females of both the OP1 and OP2 strains produced only parthenogenetic females (Fig. 3, OP1: n = 10, OP2: n = 10). In addition, during the subsequent generations under short-day conditions, they also reproduced exclusively by parthenogenesis (data not shown).

Homologs of JH-synthesis/degradation genes in the pea aphid

After updating AphidBase, the genome database for *A. pisum*, we re-surveyed and identified the genes involved in JH synthesis and degradation by comparing amino acid sequences, before investigating the photoperiodic regulation of these genes (see Supplementary Table S2). For *SDR*, we found 16 putative orthologs in the AphidBase (ver.2.1). We excluded XP_016656002 from the NCBI database

because of standard genome annotation processing. Furthermore, we narrowed these putative orthologs down to four potential orthologs using the classical SDR motif, hNhxG, on the fourth alpha helix, which was suggested to be part of the active site (see Supplementary Figure S1) (Persson et al., 2003). For JHAMT, we found 10 putative orthologs in the AphidBase (ver.2.1). Two orthologs, XP_016663097 and XP_008188739, were excluded from the NCBI database. Three of the orthologs, NP_001156251, XP_016663096, and XP_016663095, possessed identical protein sequences and an AphidBase ID (ACYPI007696). NP_001156251 and XP_001943729 have been identified in a previous study (Ishikawa et al., 2012). NP_001156251, XP_001943729, XP_003244299, and XP_016658235 were also identified in a recent study (Smykal and Dolezel, 2023). Because XP_016659614 and XP_001950668 showed relatively lower similarity to the query protein sequence compared to the other orthologs, we excluded them from the following analysis. For the CYP15A1 gene, we found an ortholog, XP_016660351, in addition to XP_001952620, identified by Daimon and Shinoda (2013), which we used as a query sequence. XP_016660351 was then removed from the NCBI database. For the JHEH gene, we identified one JHEH ortholog, XP_003241553. For the JHE gene, we found no ortholog with over 40% similarity to the JHE gene in Nilaparvata lugens. Although our previous study identified two orthologs of the JHE gene, both were excluded from the



Fig. 4. Expression profiles of JH-synthesis/degradation genes under the long- and short-day conditions. Y-axes show relative expression levels, which are specific to each gene. Averages and standard errors of three biological replicates are indicated. The result of statistical analysis is shown in Table 2. The panel of *JHAMT* and *JHEH* expression levels in the CP strain are modified from Ishikawa et al. (2012).

NCBI database. Overall, we identified four orthologs of the *SDR* gene, four orthologs of *JHAMT* genes, one ortholog of the *CYP15A1* gene, and one ortholog of the *JHEH* gene (Table 1, Fig. 2).

Photoperiodic response of JH-synthesis/degradation gene expressions in cyclical-parthenogenetic and obligate-parthenogenetic strains

To investigate the mechanisms underlying the loss of reproductive polyphenism, we quantified the photoperiodic response of gene expression that acts in the direct steps of JH synthesis. Four of six genes related to JH synthesis (CYP15A1, SDR2, SDR3, and SDR4) showed significant photoperiodic responses in the CP strain (Fig. 4, Table 2). CYP15A1 exhibited a pronounced decrease in expression under short-day conditions (two-way ANOVA, F = 31.57, P < 0.0001). In contrast, JHAMT did not show a photoperiodic response (two-way ANOVA, F = 1.842, P > 0.05) (Fig. 4, Table 2) (Ishikawa et al., 2012). Additionally, CYP15A1 expression changed between days 1 and 8 after larviposition (two-way ANOVA: F = 3.009, P < 0.05), indicating an interaction between the photoperiod and age (two-way ANOVA: F = 2.412, P < 0.05). SDR2 and SDR4 displayed higher expression levels under short-day conditions than under long-day conditions (two-way ANOVA, F = 1.715, P <0.05; F = 0.549, P < 0.05, respectively). Conversely, SDR3 expression was lower under short-day conditions (two-way ANOVA, F = 4.998, P < 0.05). SDR2 also showed increased expression on day 8 compared to that on day 1 after larviposition.

In contrast to the CP strain, neither OP1 nor OP2 displayed significant photoperiodic responses in any of the JH synthesis genes (Fig. 4, Table 2). However, in both strains, some genes showed different expression levels between days 1 and 8 after larviposition. In the OP1 strain, the *SDR1* gene showed higher expression on day 1 (two-way ANOVA: F = 1.217, P < 0.05). In the OP2 strain, *SDR2*, *SDR3*, and *CYP15A1* showed higher expression levels on day 8 (two-way ANOVA: F = 0.287, P < 0.05; F = 0.455, P < 0.05; and F = 0.101, P < 0.05, respectively).

We also investigated the response of the JH degradation gene *JHEH* to photoperiodic changes in the OP1 and OP2 strains. Our previous study on the CP strain showed that *JHEH* gene expression significantly increased under short-day conditions (two-way ANOVA: F = 1.339, P < 0.05) (Fig. 4, Table 2) (Ishikawa et al., 2012). However, in the OP1 and OP2 strains, *JHEH* expression did not differ significantly between day-length conditions (two-way ANOVA: F = 0.074, P > 0.05; F = 0.166, P > 0.05, respectively) (Fig. 4, Table 2). Our gene expression analysis revealed that obligateparthenogenetic strains did not show photoperiodic responses in JH-related genes, suggesting a potential link to the loss of the sexual phase.

Male production from obligate-parthenogenetic strains induced by acetone treatment

To investigate whether obligate-parthenogenetic strains retain the downstream developmental pathway for male production, we artificially induced male production by acetone treatment in the OP1 strain. After the application of

Cono	Strain	Sexual reproduction	<i>F-value</i>			P-value		
Gene	Sliain		Photoperiod	Age	Photoperiod x age	Photoperiod	Age	Photoperiod x age
	CP	+	0.019	0.234	0.161	0.5035	0.0902	0.1496
SDR1	OP1	-	0.038	1.217	0.002	0.43634	**0.00266	0.96325
	OP2	-	0.071	0.473	0.197	0.3606	0.0696	0.1573
	CP	+	1.715	1.004	0.174	***0.000115	***0.000654	0.060744
SDR2	OP1	-	0.441	0.592	0.008	0.0896	0.0517	0.7491
	OP2	-	0.287	1.651	0.089	0.2459	*0.0167	0.5807
	CP	+	4.998	0.313	0.393	*0.0242	0.4848	0.5137
SDR3	OP1	-	0.482	1.875	0.683	0.376	0.231	0.434
	OP2	_	0.455	3.449	0.077	0.4172	*0.0269	0.8273
	CP	+	0.549	0.414	0.025	*0.0468	0.1248	0.5954
SDR4	OP1	-	0.011	0.015	0.044	0.668	0.688	0.548
	OP2	_	0.019	0.006	0.244	0.665	0.884	0.149
	CP	+	1.842	0.349	0.128	0.0956	0.5179	0.6727
JHAMT	OP1	-	0.483	1.364	0.014	0.335	0.133	0.775
	OP2	_	1.403	0.027	0.349	0.12	0.843	0.498
	CP	+	31.57	3.022	2.412	***7.73e-06	*0.0102	*0.0162
CYP15A1	OP1	-	0.647	0.199	2.426	0.2428	0.49	*0.0238
	OP2	_	0.101	3.993	0.039	0.51476	**0.00533	0.60484
	CP	+	1.339	0	0.138	**0.00104	0.91741	0.15437
JHEH	OP1	-	0.074	0.866	0.044	0.4913	0.0985	0.6136
	OP2	-	0.166	0.002	0.116	0.412	0.852	0.372

Table 2. Two-way ANOVA results to detect the photoperiodic and age effects on genes involved in JH synthesis and degradation.

*: P < 0.05, **: P < 0.01, ***: P < 0.001

acetone, 16.7% of the aphids (n = 18) began to produce males (Fig. 5). Male producers produced one to three males (mean ± standard error: 11.1 ± 3.8%) at 9–10 days after the treatment. No males were observed in the control groups treated with ethanol (n = 21) or pentane (n = 27) (Fisher's exact test: P < 0.05). Acetone treatment also induced male production in 3% of the aphids in the OP2 strain (n = 33).

DISCUSSION

In the present study, we investigated the endocrine mechanisms underlying the loss of reproductive polyphen-



Fig. 5. Percentage of parthenogenetic female (P-female) producers (stripes with pink) and male producers (circles with light blue) induced by acetone, ethanol, and pentane treatments in the OP1 strain.



Fig. 6. Photoperiodic response of JH pathway in the CP and OP strains. The CP strain exhibits upregulation of the JH-synthesis pathway (*CYP15A* expression) and downregulation of the JH-degradation pathway (*JHEH* expression) in response to short-day length, which may be responsible for the decrease of JH titer, leading to the change of reproductive mode from parthenogenesis (P) to sexual reproduction (S). However, the OP strains do not show photoperiodic responses of the JH-synthesis/degradation gene expressions, resulting in obligate parthenogenesis. This suggests that modification of the photoperiodic sensitivity of the JH signaling pathway would underlie the multiple loss of reproductive polyphenism in the aphid life cycle.

ism in natural aphid populations. We found that five out of seven genes involved in the biosynthesis and degradation of JH showed photoperiodic responses in the CP strain. In contrast, none of these genes exhibited photoperiodic responses in the OP1 and OP2 strains. Notably, CYP15A, which encodes an enzyme for the final step of JH synthesis, showed a distinct response, with high expression under long-day conditions and decreased expression under short days in the CP strain. However, this response was completely absent in the OP1 and OP2 strains. Similarly, JHEH, which is responsible for initiating JH degradation, exhibited higher expression under short-day conditions in the CP strain; however, this response was not observed in the OP1 and OP2 strains. These results strongly suggest that the loss of the photoperiodic response in both the JH synthesis and degradation pathways may be a key mechanism underlying the elimination of the sexual phase in the aphid life cycle (Fig. 6). Aphid reproductive strategies are heavily influenced by the winter climate (Simon et al., 2010). Cyclical-parthenogenetic populations pay a two-fold cost of sex and require longer developmental arrest (Maynard-Smith, 1978), whereas they can produce cold-resistant overwintering eggs (Simon et al., 2010). In contrast, obligate-parthenogenetic populations can potentially reproduce year-round through parthenogenesis and exhibit shorter developmental periods. However, parthenogenetic females lack resistance to low temperatures (Simon et al., 2010). Therefore, reproductive polyphenism, which switches between parthenogenesis and sexual reproduction, is advantageous in areas with harsh seasonal variations. However, obligate-parthenogenetic populations potentially driven by the loss of the photoperiodic response in JH signaling may be favored in environments with milder winter temperatures (Dedryver et al., 2008).

Interestingly, we also demonstrated that topical application of acetone can induce the production of sexual males in OP strains. This result indicates that the OP1 and OP2 strains retained the developmental cascade to produce sexual males. Hormones act as molecular switches that alter gene expression patterns and initiate developmental events (Nijhout, 1999). In aphid reproductive polyphenism, JH has been suggested to integrate information on photoperiod and temperature, leading to the production of sexual morphs (Ishikawa et al., 2012). Thus, modifying the JH signaling pathway might be an effective way to eliminate the entire developmental process of the sexual phase from the aphid's annual life cycle. A previous study on Manduca sexta also showed that artificial selection to decrease larval color polyphenism resulted in the loss of sensitivity of JH titers to heat shock (Suzuki and Nijhout, 2006). Those previous findings taken together with the results of the present study indicate that changes in environmental sensitivity of hormone biosynthesis and degradation may play an important role in the evolutionary reduction of plasticity in both natural and artificially selected populations.

However, the specific mutations underlying the loss of photoperiodic responses in these genes remain unknown. Recent genomic studies have revealed that a single genomic region on the X chromosome controls the loss of ability to produce sexual females in response to a short photoperiod in pea aphids (Nouhaud et al., 2014; Huguet et al., 2024). In addition, several photoreceptors, circadian clock genes, and insulin-like peptides, such as ILP4, have been suggested to control the reproductive mode change from parthenogenesis to sexual reproduction (Goto, 2022; Colizzi et al., 2024). These genomic regions and genes are strong candidates for upstream factors that cause the loss of photoperiodic response in JH-related genes. Recently developed molecular tools, such as genome editing tools like CRISPR/cas9, can potentially enable functional assays for candidate genes or sequences that may underlie the modification of the photoperiodic control of the JH pathway. These techniques will also help to reveal the synthetic and degradation pathways of JH and identify functional JH in pea aphids. Further studies are required to fully understand the evolutionary processes that lead to the loss of reproductive polyphenism.

ACKNOWLEDGMENTS

We thank Miura laboratory members for their discussions and technical assistance, especially Hitoshi Miyakawa for his helpful comments on the manuscript. This project was supported by a DC1 Fellowship (grant no. 08J01695) from Japan Society for the Promotion of Science (JSPS) to AI, and a Grant-in-Aid for Challenging Exploratory Research (grant no. 20657004) from JSPS to TM.

COMPETING INTERESTS

We declare we have no competing interests.

AUTHOR CONTRIBUTIONS

Al designed the study, and performed experiments and data analysis. TK provided the aphid strains. Al wrote the first draft of the manuscript with HG, KO, TK, SA, and TM making significant contributions to editing. All authors agree to be held accountable for the content therein and approve the final version of the manuscript.

SUPPLEMENTARY MATERIALS

Supplementary materials for this article are available online (URL: https://doi.org/10.2108/zs240075)

Supplementary Figure S1. Alignment of putative orthologs of SDR.

Supplementary Table S1. List of primer sequences for qRT-PCR.

Supplementary Table S2. Putative orthologs of JH synthesis and degradation genes.

REFERENCES

- Ando Y, Matsumoto K, Misaki K, Mano G, Shiga S, Numata H, et al. (2020) Juvenile hormone III skipped bisepoxide, not its stereoisomers, as a juvenile hormone of the bean bug *Riptortus pedestris*. Gen Comp Endocrinol 289: 113394
- Aubret F, Shine R (2009) Genetic assimilation and the postcolonization erosion of phenotypic plasticity in island tiger snakes. Curr Biol 19: 1932–1936
- Blackman R (1987) Reproduction, cytogenetics and development.
 In "Aphids, Their Biology, Natural Enemies and Control, Vol 2A"
 Ed by AK Minks, P Harrewijn, Elsevier, Amsterdam, pp 163– 195

- Chen Z, Linse KD, Taub-Montemayor TE, Rankin MA (2007) Comparison of radioimmunoassay and liquid chromatography tandem mass spectrometry for determination of juvenile hormone titers. Insect Biochem Mol Biol 37: 799–807
- Colizzi FS, Martínez-Torres D, Helfrich-Förster C (2024). The circadian and photoperiodic clock of the pea aphid. J Comp Physiol A 210: 627–639
- Corbitt TS, Hardie J (1985) Juvenile hormone effects on polymorphism in the pea aphid, *Acyrthosiphon pisum*. Entomol Exp Appl 38: 131–135
- Daimon T, Shinoda T (2013) Function, diversity, and application of insect juvenile hormone epoxidases (CYP15). Biotechnol Appl Biochem 60: 82–91
- Dedryver CA, Le Gallic JF, Haack L, Halkett F, Outreman Y, Simon JC (2008) Seasonal and annual genotypic variation and the effect of climate on population genetic structure of the cereal aphid *Sitobion avenae* in northern France. Bull Entomol Res 98: 159–168
- Dixon AFG (1997) Aphid Ecology: An Optimization Approach. Chapman & Hall, London
- Goto SG (2022) Photoperiodic time measurement, photoreception, and circadian clocks in insect photoperiodism. Appl Entomol Zool 57: 193–212
- Hales DF, Mittler TE (1983) Precocene causes male determination in the aphid *Myzus persicae*. J Insect Physiol 29: 819–823
- Hardie J, Baker FC, Jamieson GC, Lees AD, Schooley DA (1985) The identification of an aphid juvenile hormone, and its titre in relation to photoperiod. Physiol Entomol 10: 297–302
- Hartfelder K, Emlen DJ (2012) Endocrine control of insect polyphenism. In "Insect Endocrinology" Ed by LI Gilbert, Academic Press, San Diego, pp 464–522
- Huguet MD, Robin S, Hudaverdian S, Tanguy S, Leterme-Prunier N, Cloteau R, et al. (2024) Transcriptomic basis of sex loss in the pea aphid. BMC Genomics 25: 1–17
- Ishikawa A, Hongo S, Miura T (2008) Morphological and histological examination of polyphenic wing formation in the pea aphid *Acyrthosiphon pisum* (Hemiptera, Hexapoda). Zoomorphology 127: 121–133
- Ishikawa A, Ogawa K, Gotoh H, Walsh TK, Tagu D, Brisson JA, et al. (2012) Juvenile hormone titre and related gene expression during the change of reproductive modes in the pea aphid. Insect Mol Biol 21: 49–60
- Ishikawa A, Gotoh H, Abe T, Miura T (2013) Juvenile hormone titer and wing-morph differentiation in the vetch aphid *Megoura crassicauda*. J Insect Physiol 59: 444–449
- Kamita SG, Hammock BD (2010) Juvenile hormone esterase: biochemistry and structure. J Pestic Sci 35: 265–274
- Kanbe T, Akimoto SI (2009) Allelic and genotypic diversity in longterm asexual populations of the pea aphid, *Acyrthosiphon pisum* in comparison with sexual populations. Mol Ecol 18: 801–816
- Kodama A, Matsumoto K, Shinada T, Goto SG (2023) Juvenile hormone identification in the cabbage bug *Eurydema rugosa*. Bull Entomol Res 113: 293–298
- Kotaki T, Shinada T, Kaihara K, Ohfune Y, Numata H (2009) Structure determination of a new juvenile hormone from a heteropteran insect. Org Lett 11: 5234–5237
- Kotaki T, Shinada T, Numata H (2012) Structure determination of a natural juvenile hormone isolated from a heteropteran insect. Psyche (Camb Mass) 2012: 924256
- Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. J Evol Biol 22: 1435–1446
- Lees AD (1959) The role of photoperiod and temperature in the determination of parthenogenetic and sexual forms in the aphid *Megoura viciae* Buckton—I: The influence of these factors on apterous virginoparae and their progeny. J Insect

Physiol 3: 92-117

- Lees AD (1963) The role of photoperiod and temperature in the determination of parthenogenetic and sexual forms in the aphid Megoura viciae Buckton-III. Further properties of the maternal switching mechanism in apterous aphids. J Insect Physiol 9: 153-164
- Li Y, Park H, Smith TE, Moran NA (2019) Gene family evolution in the pea aphid based on chromosome-level genome assembly. Mol Biol Evol 36: 2143-2156
- Mackay PA, Wellington WG (1977) Maternal age as a source of variation in the ability of an aphid to produce dispersing forms. Res Popul Ecol 18: 195-209
- Mano G, Goto SG (2022) Photoperiod controls insulin and juvenile hormone signaling pathways via the circadian clock in the bean bug Riptortus pedestris (Hemiptera: Alydidae). Appl Entomol Zool 57: 363-377
- Matsumoto K, Yasuno Y, Yasuda K, Hayashi T, Goto SG, Shinada T (2020) Structure determination of juvenile hormone from chagas disease vectors, Rhodnius prolixus and Triatoma infestans. Chem Lett 49: 538-541
- Maynard-Smith J (1978) The Evolution of Sex. Cambridge University Press, London
- Mayoral JG, Nouzova M, Navare A, Noriega FG (2009) NADP+dependent farnesol dehydrogenase, a corpora allata enzyme involved in juvenile hormone synthesis. Proc Natl Acad Sci U S A 106: 21091-21096
- Miura T (2005) Developmental regulation of caste-specific characters in social-insect polyphenism. Evol Dev 7: 122-129
- Moran N (1992) The evolution of aphid life cycles. Annu Rev Entomol 37: 321-348
- Nijhout HF (1994) Developmental perspectives on evolution of butterfly mimicry. Bioscience 44: 148-157
- Nijhout HF (1999) Control mechanisms of polyphenic development in insects: in polyphenic development, environmental factors alter some aspects of development in an orderly and predictable way. Bioscience 49: 181-192
- Nijhout HF (2003) Development and evolution of adaptive polyphenisms. Evol Dev 5: 9-18
- Nouhaud P, Peccoud J, Mahéo F, Mieuzet L, Jaquiéry J, Simon JC (2014) Genomic regions repeatedly involved in divergence among plant-specialized pea aphid biotypes. J Evol Biol 27: 2013-2020
- Ogawa K, Miura T (2014) Aphid polyphenisms: Trans-generational developmental regulation through viviparity. Front Physiol 5: 1
- Persson B, Kallberg Y, Oppermann U, Jörnvall H (2003) Coenzymebased functional assignments of short-chain dehydrogenases/ reductases (SDRs). Chem Biol Interact 143-144: 271-278
- Pfennig DW, Wund MA, Snell-Rood EC, Cruickshank T, Schlichting CD, Moczek AP (2010) Phenotypic plasticity's impacts on diversification and speciation. Trends Ecol Evol 25: 459-467
- Pigliucci M (2001) Phenotypic Plasticity: Beyond Nature and Nurture. Johns Hopkins University Press, Baltimore
- R Core Team (2014) R: A language and environment for statistical computing

Schwander T, Leimar O (2011) Genes as leaders and followers in

evolution. Trends Ecol Evol 26: 143-151

- Schwartzberg EG, Kunert G, Westerlund SA, Hoffmann KH, Weisser WW (2008) Juvenile hormone titres and winged offspring production do not correlate in the pea aphid, Acyrthosiphon pisum. J Insect Physiol 54: 1332-1336
- Shinoda T, Itoyama K (2003) Juvenile hormone acid methyltransferase: A key regulatory enzyme for insect metamorphosis. Proc Natl Acad Sci U S A 100: 11986-11991
- Shuo Y, Wang WX, Shen J (2020) Reproductive polyphenism and its advantages in aphids: Switching between sexual and asexual reproduction. J Integr Agric 19: 1447-1457
- Simon JC, Rispe C, Sunnucks P (2002) Ecology and evolution of sex in aphids. Trends Ecol Evol 17: 34-39
- Simon JC, Stoeckel S, Tagu D (2010) Evolutionary and functional insights into reproductive strategies of aphids. C R Biol 333: 488-496
- Smykal V, Dolezel D (2023) Evolution of proteins involved in the final steps of juvenile hormone synthesis. J Insect Physiol 145: 104487
- Suzuki Y, Nijhout HF (2006) Evolution of a polyphenism by genetic accommodation. Science 311: 650-652
- Tanaka S (2001) Endocrine mechanisms controlling body-color polymorphism in locusts. Arch Insect Biochem Physiol 47: 139-149
- Tsang SSK, Law STS, Li C, Qu Z, Bendena WG, Tobe SS, et al. (2020) Diversity of insect sesquiterpenoid regulation. Front Genet 11: 1027
- Villalobos-Sambucaro MJ, Nouzova M, Ramirez CE, Eugenia Alzugaray M, Fernandez-Lima F, Ronderos JR, et al. (2020) The juvenile hormone described in Rhodnius prolixus by Wigglesworth is juvenile hormone III skipped bisepoxide. Sci Rep 10: 3091
- West-Eberhard MJ (2003) Developmental Plasticity and Evolution. Oxford University Press, Oxford
- Westerlund SA, Hoffmann KH (2004) Rapid quantification of juvenile hormones and their metabolites in insect haemolymph by liquid chromatography-mass spectrometry (LC-MS). Anal Bioanal Chem 379: 540-543
- Wilkinson TL, Ishikawa H (2000) Injection of essential amino acids substitutes for bacterial supply in aposymbiotic pea aphids (Acyrthosiphon pisum). Entomol Exp Appl 94: 85-91
- Wojtasek H, Prestwich GD (1996) An insect juvenile hormone-specific epoxide hydrolase is related to vertebrate microsomal epoxide hydrolases. Biochem Biophys Res Commun 220: 323-329
- Yi G, Ba R, Luo J, Zou L, Huang M, Li Y, et al. (2023) Simultaneous detection and distribution of five juvenile hormones in 58 insect species and the absolute configuration in 32 insect species. J Agric Food Chem 71: 7878–7890
- Zera AJ, Denno RF (1997) Physiology and ecology of dispersal polymorphism in insects. Annu Rev Entomol 42: 207-230
- (Received August 16, 2024 / Accepted October 15, 2024 / Published online February 11, 2025)