

Regulation of the IGF System by Glucocorticoids

Authors: Dell, Ghislaine, Ward, Andrew, Shokrai, Arman, Madej, Andrej, and Engström, Wilhelm

Source: Zoological Science, 16(3) : 377-385

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.16.377>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

[REVIEW]

Regulation of the IGF System by Glucocorticoids

Ghislaine Dell¹, Andrew Ward¹, Arman Shokrai², Andrej Madej³ and Wilhelm Engström^{2*}

¹*Developmental Biology Programme, School of Biology and Biochemistry,
University of Bath, BATH, UK,*

²*Department of Pathology, Faculty of Veterinary Medicine, Swedish University of Agricultural Sciences,
PO Box 7028, S-750 07 Uppsala, Sweden and*

³*Department of Physiology, Faculty of Veterinary Medicine, Swedish University of Agricultural Sciences,
PO Box 7045, S-750 07 Uppsala, Sweden*

Introduction

Insulin-like growth factors (IGFs) play a pivotal role in promoting embryonic and fetal growth, and display a wider range of developmental and tissue specific expression than any other growth factors. Glucocorticoids exert a variety of anabolic and catabolic effects and are involved in the organisms response to stress. During development, there appears to be a co-ordinated regulation of glucocorticoid biosynthesis and IGF gene expression in the embryo. In particular, attention has been focussed on the role of the glucocorticoid surge just before birth in regulating IGF levels in the embryo. IGFs and glucocorticoids have been known to have corroborating effects on cell proliferation, with glucocorticoids affecting IGFs at a transcriptional level.

The purpose of this review, then, is to collate the current knowledge of how glucocorticoids can effect regulation of IGFs, both directly and via regulation of the IGF receptors and binding proteins, which themselves regulate IGFs and their effects. We show that this regulation is complex and bi-directional, occurring in different tissue types to up- or downregulate levels of IGF or IGF-binding protein transcription or protein levels, intricately linking the glucocorticoids to the insulin like growth factors in the control of cell growth and proliferation.

The insulin-like growth factors and their receptors

The insulin like growth factors IGF I and IGF II are single chain polypeptides with an approximate molecular weight of 7 kDa (Daughaday and Rotwein, 1989). The mature peptides consist of four distinct domains in IGF I as well as in IGF II-A, B, C and D. The A and B chains show strong homology with preproinsulin. Both IGF I and IGF II are produced as pre-peptides that contain a signal peptide as well as a trailer peptide. In IGF I, there are different signal and trailer peptides that combine to yield different precursor molecules. Hence multistep posttranslational processing is required to obtain

identical end-products.

IGFs were discovered on the basis of their ability to stimulate cartilage sulphation and to replace the sulphation activity of growth hormone both in *in vivo* and *in vitro* test systems (Salmon and Daughaday, 1957). The biological significance of this finding was rapidly expanded beyond the study of cartilage sulphation to include stimulation of DNA replication, proteoglycan synthesis, glycosamin synthesis, protein synthesis and accumulation, motility and cell survival (see Jones and Clemmons, 1995 for review). Purification and subsequent amino acid sequence determination revealed the existence of two separate molecules that contain 70 and 67 amino acids, respectively. Due to their high degree of homology with insulin they were denominated IGF I and IGF II (Rinderknecht and Humbel, 1978a, 1978b).

IGFs bind to, and act via three different membrane receptors: the type I and type II IGF receptors and the insulin receptor. The affinities and kinetic properties differ among each of the ligand-receptor interactions. The type I IGF receptor has the highest affinity for IGF I and the type II receptor the highest affinity for IGF II. The insulin receptor binds both IGF I and IGF II with low affinity (Nissey and Kiess, 1991; Steele-Perkins *et al.*, 1988; Werner *et al.*, 1991). The type I IGF receptor resembles the insulin receptor. It is a heterodimeric transmembrane protein that consists of two alpha and two beta subunits. Ligand binding induces tyrosine specific autophosphorylation of the receptor as well as of cytoplasmic substrate proteins which is followed by a multifaceted biological response. The type I IGF receptor mediated a vast variety of biological effects exerted by the IGFs (de Meyts *et al.*, 1994; Jones and Clemmons, 1995) and recent mouse genetic experiments show that it is the major mitogenic signalling receptor for both IGF I and IGF II in fetal development (Baker *et al.*, 1993). The functional relationship between IGF II and the insulin receptor was unclear for some time, in particular since tumour hypoglycaemia increased the rate of IGF II gene transcription (Schofield *et al.*, 1991). A recent study shed some light on this issue by demonstrating that the insulin receptor

* Corresponding author: Tel. +46-18-67 11 94;
FAX. +46-18-67 35 32.

can mediate the mitogenic messages of IGF II but not of IGF I (Morrione *et al.*, 1997). Furthermore, IGF II mitogenic signalling via the insulin receptor is known to be necessary for full growth of the placenta. In contrast, the type II IGF receptor is a monomeric protein which consists of a major extracellular portion with fifteen repeats of a cysteine rich sequence, as well as a single hydrophobic transmembrane helix and a minor cytoplasmic part. The type II receptor in mammals is also the mannose-6-phosphate receptor (Morgan *et al.*, 1987) and binds IGF II and Mannose-6-phosphate at distinct binding sites on the receptor protein (Braulke *et al.*, 1988). Binding of either of the ligands does not induce a phosphorylation response from this receptor, rather its role appears to be to participate in endocytosis as well as sorting of lysosomal enzymes. The type II receptor is also involved in membrane trafficking through rapid cycling between cytosolic membrane compartments and the plasma membrane. It also induces a redistribution of receptors (Braulke and Mieskes, 1992) as well as modulating insulin exocytosis under physiological conditions (Zhang *et al.*, 1997). In keeping with this cell biology, mouse genetic experiments indicate that the type II receptor acts primarily as a scavenger for IGF II. Loss or inactivation of the type II receptor gene results in a general overgrowth that is ameliorated in the absence of the IGF II ligand (Filson *et al.*, 1993; Wang *et al.*, 1994).

The insulin like growth factors IGF I and IGF II display a wider range of developmental and tissue specific expression than any other known growth factors (Schofield, 1992). It is generally implied that they play a pivotal role in promoting embryonic and fetal growth. Although the IGFs were originally believed to act as a classical hormone, mediating the action of growth hormone, they are now known to act in a paracrine or autocrine fashion. In the fetus and adult, both IGFs are mainly synthesised in the liver. However nearly all embryonic tissues express the IGF II gene (Hydahl *et al.*, 1986; Scott *et al.*, 1985). During development many fetal tissues express one or both IGFs with expression detectable from early post implantation and onwards. It is noteworthy that type I IGF receptors are expressed either by the IGF expressing cells or by adjacent cells, which forms a prerequisite for paracrine or autocrine loops.

A large variety of normal and neoplastic cells cultured *in vitro* express the IGF II gene. The level of expression can be influenced by a variety of culture conditions including the serum concentration. In addition to the bona fide 67 amino acid IGF II protein, there are examples of high molecular weight variants produced by cells cultured *in vitro* (Gowan *et al.*, 1987; Granerus *et al.*, 1993; Schofield *et al.*, 1990) that show a different affinity to the IGF receptors (Schofield *et al.*, 1994). The biological implications of these variant IGF II molecules are unclear, but it has been suggested that competition for the type I receptor might modulate the amplitude of the biological response (Schofield *et al.*, 1994). IGF II exerts a wide range of biological activities in cells in culture. It can promote cell proliferation by acting on the chromosome cell cycle (ie DNA-replication and mitosis) as well as on the cell growth

cycle (cellular enlargement) (Dafgård *et al.*, 1986; Zetterberg *et al.*, 1982, 1984). IGF II can induce differentiation *in vitro*, an effect which has been characterised in detail in myoblasts (Florini *et al.*, 1991). IGF II can counteract apoptosis in some cell systems and thereby enhance survival (Granerus *et al.*, 1995; Granerus and Engström, 1996), whereas in other cell lines there appears to be an apoptosis inducing effect by IGF II (Granerus *et al.*, 1998). IGF II release also induces a functional modulation of certain cell types. It stimulates hormone synthesis and secretion in ovarian granulosa and theca cells (Giudice, 1992). It also binds to the type I receptor and thereby potentiates the release of histamine from basophils in response to immunoglobulin E (Hirai *et al.*, 1993). Finally, it has been shown that IGF II can stimulate motility in cultured rhabdomyosarcoma cells (Minitti *et al.*, 1992).

Overexpression of IGFs in transgenic mice has resulted in altered growth properties. Increased expression of an IGF I transgene leads to increased bodyweight and a limited overgrowth. Different tissues responded differently and growth disturbances and tumour formation were sometimes observed (Bol *et al.*, 1997; Coleman *et al.*, 1995; Matthews *et al.*, 1988; Reiss *et al.*, 1996). In several experimental situations, prolonged IGF II expression from transgenes using tissue-restricted regulatory elements has led to organ overgrowth and/or tumour formation (Bates *et al.*, 1995; Rogler *et al.*, 1994; Rossetti *et al.*, 1996; van Buul-Offers *et al.*, 1995; Ward *et al.*, 1994). More generalised IGF II overexpression has been achieved by introducing additional copies of the IGF II gene into embryonic stem cells, which were then used to generate chimaeric mice. An alternative approach to study how increased levels of IGF II can affect overall growth properties was to assay double mutant mice carrying a deletion around the H19 region as well as a targeted IGF type 2 receptor allele. Such mice have extremely high levels of IGF II and display most of the clinical features of the Wiedemann-Beckwith syndrome, as well as skeletal defects and a cleft palate: features of Simpson-Golabi-Behmel syndrome (Eggenschwiler *et al.*, 1997). In both these models of more general overgrowth, the affected animals die perinatally, thus making it impossible to assess their susceptibility to neoplasms. The development of transgenic technology has also rapidly made it possible to examine the effects of growth factor deficiency *in vivo*. When a disrupted IGF II gene was introduced into the mouse germ line, the prenatal growth rate decreased and the body weight at term only reached 60% of the normal birth weight. However, the growth rate post partum appeared to be normal (deChiara *et al.*, 1990). Likewise, knockout-mice carrying null mutations for the IGF I gene lead to a significantly decreased birthweight, but with otherwise normal body proportions. Unlike the IGF II deficient mice, these transgenic animals had a decreased postnatal growth rate and a high degree of neonatal lethality (Baker *et al.*, 1993; Liu *et al.*, 1993).

The glucocorticoids and their receptors

Glucocorticoid hormones are biologically active steroid compounds that bear a close structural resemblance to one

another. Their biosynthesis is regulated by the hypothalamo-pituitary-adrenal axis which includes a number of intrinsic feedback mechanisms. In brief, a small peptide-corticotropin releasing factor (CRF) is secreted by the hypothalamus and induces the synthesis of adrenocorticotrophic hormone (ACTH) from the pituitary gland. ACTH subsequently stimulates the production and release of cortisol and other steroids from the adrenal cortex. Increased levels of cortisol also act as a feedback mechanism, whereby CRF and ACTH are inhibited.

Glucocorticoids induce glucose biosynthesis as well as exerting a variety of anabolic and catabolic effects. High levels of glucocorticoids suppress inflammation and the host immune response. This capacity has made them useful in pharmacological treatment of autoimmune diseases and suppression of transplant rejection. The physiological relevance of these profound effects on the modulation of the immune system are debatable since only pharmacological doses of glucocorticoids tend to yield them.

Glucocorticoids are involved in the organism's response to stress. The role of glucocorticoids in this context is to ensure that the brain and other critical tissues are adequately supplied with glucose. Glucocorticoids hence do not counteract the stress itself but rather abrogate the body's harmful response to stress.

The glucocorticoids are by nature hydrophobic and therefore transported in the bloodstream complexed with transcortin or corticosteroid binding proteins (CBG). Once released from its binding protein, the glucocorticoid enters the cell by diffusion or active transport. In the cytoplasm, glucocorticoids bind reversibly to a specific glucocorticoid receptor (see Wright *et al.*, 1993 for a review). This receptor is maintained in an inactive state in a multiprotein complex, consisting of one receptor molecule and several heat shock proteins including hsp90, hsp70 and hsp56. Upon binding of the glucocorticoid, a dimer of hsp90 proteins is released from the complex whereby the glucocorticoid receptor acquires an increased affinity for DNA. The activated receptor dimerises before binding to specific sequences in the DNA. The glucocorticoid can alter the expression of target genes through at least three different mechanisms; (i) recruitment of the general transcriptional machinery (ii) modulation of transcription factor action, independent of DNA binding, through direct protein-protein interactions, and (iii) modulation of chromatin structure to allow the assembly of other gene regulatory proteins and/or general transcription machinery on the DNA. (McEwan *et al.*, 1997)

The specificity of glucocorticoid action is determined by well conserved DNA sequences which are bound by zinc finger domains within the activated receptor. The target DNA sequences are referred to as glucocorticoid response elements (GREs). A specific recognition site for the glucocorticoid receptor was first identified in the Mouse Mammary Tumour Virus long terminal repeats (MMTV-LTR). Subsequently, elements which mediate the action of glucocorticoids were found in the vicinity of a number of genes. By comparing the sequences a 15 base pair consensus motif was identified. It contains two partially palindromic hexamers (Zilliaccus *et al.*,

1994) with a 3 base pair spacer in between. The conserved palindromic sequences as well as the spacer, which could consist of any three bases are critical for receptor mediated action *in vivo*. Activated glucocorticoid receptors, like other steroid receptors, bind to GREs and affect transcription. Therefore the hormonal response must be determined by the presence or absence of particular receptors and/or receptor specific requirements for additional factors to achieve transcriptional modulation within a given cell.

Considerably less is known about how DNA binding represses transcription than about how it activates it (see Dahlman-Wright, 1991 for review). Studies of a variety of genes that are downregulated by glucocorticoids suggest that the activated glucocorticoid receptor interferes with the binding or activity of other transcription factors.

A consensus sequence for receptor binding sites which mediate negative regulation—ATYACnnTnTGATCn—was proposed by Beato and co-workers (Beato, 1989). The biological significance of this element remains to be clarified.

One area that has attracted much attention has been the potential interplay between the signal transduction pathways mediating responses by the glucocorticoid receptor and the AP-1 family of transcription factors. It was shown that collagenase promoter activity is inhibited by glucocorticoids via the AP-1 site with a stringent requirement of glucocorticoid receptor. C-jun, which is one component of the AP-1 complex, inhibits the transactivation of the glucocorticoid receptor (Jonat *et al.*, 1990; Schule *et al.*, 1990; Yang *et al.*, 1990). These authors also demonstrated a direct interaction between c-jun and the glucocorticoid receptor where they were found to inhibit each other's binding. C-fos, another component of the AP-1 complex, was found to transrepress the glucocorticoid receptor in a similar fashion (Lucibello *et al.*, 1990). Subsequently a distinct modulating domain was identified in glucocorticoid receptor monomers that repress the activity of the transcription factor AP-1 (Heck *et al.*, 1994).

The effects of glucocorticoids on IGF transcription

The combined effects of insulin like growth factors and glucocorticoid hormones on cell proliferation has been known for some time (Concover *et al.*, 1983). However, a direct effect of glucocorticoid administration on the transcriptional activity of the IGF II gene was first documented by Beck *et al.* (1987, 1988) and Levinovitz and Norstedt (1989). In both studies neonatal rats were injected with the cortisone analogue dexamethasone and the resultant effects on IGF II transcription in the liver were monitored. The reduction in IGF II transcription levels was dramatic which led the authors to suggest a general down-regulatory action of glucocorticoid hormones on IGF II transcription. Similar results have since been reported in a variety of biological systems. Administration of the naturally occurring active steroid hormone corticosterone resulted in a rapid and significant decrease in hepatic mRNA levels in neonatal rats (Kitriki *et al.*, 1992). Further support for this notion was given in a study in which a patient with non-islet cell tumour hypoglycaemia was treated

with prednisolone following which IGF II production was suppressed (Baxter, 1996). In other animals, profound effects of glucocorticoid administration on the IGF II gene has been demonstrated. In sheep, infusion of cortisol as well as ACTH led to a decrease in IGF II expression in fetal adrenal glands (Lu *et al.*, 1994). Moreover, it has recently been shown that infusion of cortisol to fetal sheep during late gestation also results in a significant down-regulation of hepatic IGF II mRNA abundance (Forhead *et al.*, 1998). This effect was shown to be mediated by a specific suppression of the ovine P4 promoter (Li *et al.*, 1998). Pregnant minks that were treated with polychlorinated biphenyls increased their endogenous production of glucocorticoids which resulted in a decrease of IGF II transcription in the maternal liver (Bäcklin *et al.*, 1998). It has been speculated whether the temporary release of cortisol which normally occurs soon after birth in fact is responsible for their normal post-natal decline in IGF II transcription (Dalle *et al.*, 1985). The studies involving glucocorticoid administration during pregnancy suggested that this control point could be brought forward to an earlier developmental stage. Downregulation of gene transcription by glucocorticoids, has also been observed in the IGF I gene. Dexamethasone was shown to down regulate IGF I mRNA levels in rat neuronal and glial cells *in vitro* (Adamo *et al.*, 1988). Cortisol decreased the transcriptional activity of the IGF I gene in human osteoblast cells *in vitro* (Swolin *et al.*, 1996).

The generality of the concept that glucocorticoids suppress IGF gene expression has been challenged several times over the last decade. For instance it was shown that dexamethasone treatment of pregnant rats resulted in a limited but significant increase in IGF II mRNA levels in fetal livers (Price *et al.*, 1992). There seemed to be a certain amount of organ specificity, since lung tissues taken from the same animals appeared to contain similar levels of IGF II transcript whether or not their mothers had received glucocorticoid treatment (Price *et al.*, 1992). In a series of adult male volunteers, dexamethasone did not affect serum levels of IGF II, but this result does not exclude local differences in transcriptional activity between different organs (Miell *et al.*, 1994). Interestingly, several cell types grown *in vitro* increase their production of IGF II mRNA in response to glucocorticoid addition to the culture medium including pheochromocytoma cells (Liu *et al.*, 1994) and mouse myogenic cells (Yoshiko *et al.*, 1998). When pregnant sows were injected with either dexamethasone or hydrocortisone their first trimester fetuses contained increased quantities of IGF II mRNA driven from the fetal promoters (Madej *et al.*, 1996; unpublished).

The up-regulation of IGF II expression in response to glucocorticoids is difficult to square with the absence of obvious GREs in some of the IGF II promoter regions. However, it has recently been shown that an isolated IGF II P3 promoter construct can be activated by glucocorticoids in the absence of a GRE. The augmentation of this stimulatory effect by other flanking enhancer elements pointed at the possible existence of indirect mechanisms for glucocorticoid induced gene activation. In principle, this is testable in cell culture systems where

simultaneous addition of hormone and protein synthesis inhibitors would ameliorate any indirect effects on IGF expression. It should be noted, however, that there seems to exist a coordinated regulation of glucocorticoid biosynthesis and IGF gene expression, at least in the embryo (Yuan *et al.*, 1996). Also, insulin-like growth factors appear to exert an effect on steroid production, thereby providing evidence for further complexity in steroid-IGF interplay (Mesiano *et al.*, 1997)

IGF binding proteins and glucocorticoids

IGF binding proteins are secreted proteins which comprise a major regulatory component of the IGF signalling pathway (Cohick and Clemmons, 1993). Six IGFBPs have so far been characterised; all bind IGFs with high affinity and are capable of modifying the biological actions of IGFs within tissues as well as transporting the IGFs between body compartments. They are separate gene products, but much of the amino acid sequence and multiple cystine bridges are conserved among the different IGFBPs (Drop *et al.*, 1992; Shimasaki and Ling, 1991). Each IGFBP has a distinct tissue-specific and developmentally regulated expression pattern (Cheung *et al.*, 1994; Lindenberg-Kortleve *et al.*, 1997; Shimasaki and Ling, 1991), as well as differing affinities for the IGFs. This has led to the belief that each IGFBP has a specialised role in each tissue, modulating the activity of IGF I and II in a negative (complexing with IGFs to limit availability of free ligand for interaction with its receptor) or a positive (facilitating interaction with the receptor) manner. For example, IGFBP-6 binds IGF II with high affinity and prevents IGF II-mediated effects (Gabbitas and Canalis, 1997), whereas IGFBP-3, which is held to be the major carrier protein for IGFs, can either inhibit or potentiate actions of IGFs depending on the target cell type (Clemmons, 1992). It should also be noted that at least some of the IGFBPs may also exert effects that are independent of IGFs.

We have already reviewed the direct effects that glucocorticoids have on IGFs. As the IGFBPs clearly have an important part to play in regulating IGF activity, we should also consider whether glucocorticoids can regulate IGFs indirectly by regulating IGFBPs. This is indeed the case, especially so in tissues where IGFs are developmentally important including bone, muscle, and liver. As IGFBPs can either inhibit or potentiate IGF actions, it should be no surprise that they can be regulated in different directions by the same compound in different cell types. The interaction between glucocorticoids (mostly dexamethasone) and IGFBPs is better documented for some members of the family than others.

A. Glucocorticoids and IGFBP-1

The observation by Price *et al.* (1992) that treatment of pregnant rats with dexamethasone led to an 8-fold increase in IGFBP-1 mRNA expression in foetal livers and a smaller increase in foetal lungs was corroborated by *in vitro* experiments using primary rat hepatocytes (Miura *et al.*, 1992; Robertson *et al.*, 1994), human (Suwanichkul *et al.*, 1994) and rat hepatoma cells (Goswami *et al.*, 1994; Orłowski *et al.*,

1989; Suh *et al.*, 1996). In rat foetal osteoblasts also demonstrated a rise in IGFBP-1 mRNA and protein after dexamethasone treatment, suggesting that IGFBP-1 can modulate local IGF actions on bone formation in response to changes in glucocorticoid concentration. Similar regulation has been observed in foetal osteoblasts (Concover *et al.*, 1995). Also, cortisol levels in cord blood and IGFBP-1 levels were found to be directly correlated in newborn infants (Concover *et al.*, 1996). Further work has been done to elucidate the mechanism of the regulation. In cells, dexamethasone regulated IGFBP-1 at the level of transcription by interacting with a GRE at -91/-77 relative to the start site of transcription of the IGFBP-1 promoter (Suh *et al.*, 1996), although another putative GRE was identified further upstream (99). This GRE is essential for dexamethasone-stimulated activity; its deletion reduced transcription to basal levels. However, this GRE is of low affinity and for maximal basal and glucocorticoid-stimulated activity, an insulin response element, HNF-1 site and an AP-2 site are also necessary (Goswami *et al.*, 1994; Robertson *et al.*, 1994; Suh *et al.*, 1996, 1997; Suwanichkul *et al.*, 1994). As IGFBP-1 is unique among IGFBPs in its rapid regulation to metabolic and hormonal changes (Suh *et al.*, 1997), it is not surprising to find that its regulation is influenced by insulin and glucocorticoids.

B. Glucocorticoids and IGFBP-2

Much less is known about the regulation of IGFBP-2 by glucocorticoids. However, an important interaction between the two was shown by Mouhediene *et al.* (1996). Glucocorticoids are thought to be involved in lung maturation; IGFs are known to be important in development and differentiation. Incubation of rat type 2 stem cells of the alveolar epithelium with dexamethasone showed a marked increase, again at the transcriptional level, of IGFBP-2 mRNA and protein. In a key *in vivo* study (Price *et al.*, 1992), IGFBP-2 was also found to be increased in foetal lung. An *in vitro* study (Mouhediene *et al.*, 1996) showed that an increase in IGFBP-2 correlated with a decrease in DNA synthesis. On further investigation of the regulation, a putative GRE was found in the IGFBP-2 promoter. It is not known how IGFBP-2 affects IGF II in this system, although the up-regulation of IGFBP-2 was accompanied by an increase in IGF II and IGF2R. It is plausible that glucocorticoid directs the lung towards differentiation rather than growth through IGF II. IGFBP-2 expression and mRNA levels were also found to be upregulated in pancreatic cell lines after treatment with dexamethasone (Katz *et al.*, 1997); this could also indicate a role in differentiation control for this IGF binding protein.

C. Glucocorticoids and IGFBP-3

The majority of total IGFs circulate as a complex with ubiquitously expressed IGFBP-3; formation of this complex alters IGF distribution and clearance, and modifies IGF bioactivity (Villafuerte *et al.*, 1995). IGFBP-3 levels are also modulated by a dynamic balance between soluble and membrane-bound protein (McCusker *et al.*, 1990). Bearing this in

mind, it is not surprising that glucocorticoids have been reported to up- and downregulate this protein in different tissues. Hepatic IGFBP-3, which probably represents the majority of circulating IGFBP-3, is downregulated after exposure of primary rat hepatocytes (co-cultured parenchymal and non-parenchymal cells) (Villafuerte *et al.*, 1995). This regulation was dose-dependent using physiological (10⁻¹⁰–10⁻⁸M) and pharmacological (10⁻⁶M) levels of dexamethasone. No GRE was found in the IGFBP-3 promoter, thus an indirect regulation was proposed. Dexamethasone was also found to decrease IGFBP-3 in human osteoblast cultures (Chevalley *et al.*, 1996). In bone, IGFBP-3 enhances IGF actions and this was thus proposed as a mechanism by which glucocorticoids could inhibit bone formation by inhibiting IGF anabolic activity.

In dermal papilla explants, however, dexamethasone was reported to increase IGFBP-3 levels eightfold (Hembree *et al.*, 1996). As IGFs are involved in hair follicle elongation, and glucocorticoids have been reported to suppress hair growth, it was proposed that in this system IGFBP-3 is acting to inhibit the mitogenic effects of IGFs, presumably by sequestering free IGF. In contrast to the results of Villafuerte *et al.* (1995), an earlier *in vivo* study showed an upregulation of IGFBP-3 at the protein and mRNA levels in rats treated with dexamethasone (Luo and Murphy, 1990), pointing at a mechanism for dexamethasone-induced growth retardation. Both serum and hepatic levels were affected. This difference may purely be due to the difference between an *in vivo* and an *in vitro* system, and does indeed point to an indirect effect of glucocorticoids on IGFBP-3.

D. Glucocorticoids and IGFBP-4

A different and rather attractive mechanism has been proposed for glucocorticoid regulation of IGFBP-4, which has been postulated to be important in the nervous system (Cheung *et al.*, 1994). When dexamethasone was applied to cultures of rat neuronal cells (Cheung *et al.*, 1994), abundance of native IGFBP-4 protein dropped to 10% of control levels. Unusually, this was not accompanied by a change in mRNA levels but an increase in a breakdown product of IGFBP-4 was observed. Dexamethasone had induced a protease which specifically cleaves IGFBP-4, thus reducing native levels and increasing levels of a smaller form with lower affinity for IGFs. It is difficult to speculate on how this would affect IGF actions; IGFBP-4 has been shown *in vitro* to inhibit IGFs, presumably because its affinity for IGFs is higher than that of the IGF1 receptor (Orlowski *et al.*, 1989). Thus, proteolysis of IGFBP-4 could permit IGF action by allowing access of IGF to its receptor (Cheung *et al.*, 1994). Dexamethasone has also been shown to inhibit basal IGFBP-4 secretion in bovine and human fibroblasts (Concover *et al.*, 1995), but here an effect on mRNA levels was also observed, indicating a different mechanism at work. In cultured pancreatic cells, however, dexamethasone stimulated secretion of IGFBP-4 (Katz *et al.*, 1997). This would appear to provide further evidence for a complex cell-specific regulation of IGFBPs by glucocorticoids.

E. Glucocorticoids and IGFBP-5

Again, not much is known about this IGFBP regarding its physiological role. It is thought to enhance IGF actions, and its expression was found to be decreased after treatment of human osteoblasts with dexamethasone (Chevally *et al.*, 1996). This, together with the regulation already described of IGFBP-3 and IGFBP-4 in this system, suggests a role for IGFbps in controlling IGF activity in bone formation.

F. Glucocorticoids and IGFBP-6

IGFBP-6 binds IGF II with high affinity (at least 20 times higher) than IGF I and prevents IGF II-mediated effects on myoblasts and osteoblasts (Bach *et al.*, 1993), indicating a role for this IGFBP in control of muscle and bone differentiation in fetal growth (Gabbitas and Canalis, 1997). Glucocorticoids induce the expression of IGFBP-6 in cultured foetal rat osteoblasts (Gabbitas and Canalis, 1996): cortisol induced a time- and dose-dependent increase in IGFBP-6 mRNA and protein, at the transcriptional level, indicating a possible mechanism for the inhibitory effects of glucocorticoids on bone formation. In contrast, though, dexamethasone decreased IGFBP-6 protein and mRNA levels in rat PC12 pheochromocytoma cells, also at the level of transcription (Bach *et al.*, 1993). This cell-specific regulation, coupled with the fact that glucocorticoids induce differentiation of PC12 cells to a chromaffin rather than a neural phenotype, suggests that IGFBP-6 and the IGF system may be involved in chromaffin differentiation of these cells.

Glucocorticoids and IGF receptors

Considering the importance of the IGF receptors in the IGF signalling system, surprisingly little is known about their regulation by glucocorticoids. The *in vivo* study conducted by Price *et al.* (1992) in rats showed that exposure of fetuses to dexamethasone resulted in increased IGF1R mRNA levels in liver and lung (the two tissues examined), suggesting that decreased receptor availability does not contribute to dexamethasone-induced growth retardation. Two interesting effects of glucocorticoids on IGF receptors have been documented. Firstly, in rat fetal osteoblasts, cortisol did not change IGF1R mRNA levels but did time- and dose-dependently transcriptionally decrease IGF2R mRNA and protein levels (Rydzziel and Canalis, 1995). IGF2R is thought mainly to be a sink for IGFs, lowering free IGF levels, so this result, which would have the effect of raising free IGF levels, is intriguing. Secondly, two contrasting but similar effects of dexamethasone have been observed. Dexamethasone blocked the IGF I-induced increase in IGF binding in rat chondrocytes, without having any effects on basal IGF1R levels (Jux *et al.*, 1998). In porcine ovary granulosa cells, however, dexamethasone prevented the normal decrease in IGF1R levels induced by IGF I (Urban *et al.*, 1994), effectively raising IGF1R levels but with no change in mRNA or protein levels. These would appear to be indirect effects, dexamethasone blocking the effects of IGF I and thus affecting the receptor, and it would be interesting to ascertain the mechanism behind this.

Acknowledgements

The authors wish to acknowledge the financial support of Cancerfonden, Barncancerfonden and the Cancer Research Campaign of Great Britain

REFERENCES

- Adamo M, Werner H, Farnsworth W, Roberts CT, Raizada M, LeRoith D (1988) Dexamethasone reduces steady state insulin like growth factor I mRNA levels in rat neuronal and glial cells in primary culture. *Endocrinol* 123: 2565–2570
- Bach LA, Leeding KS, Leng SI (1993) Regulation of IGF binding protein 6 by dexamethasone and IGF:s in PC12 rat pheochromocytoma cells. *J Endocrinol* 155: 225–232
- Bäcklin BM, Gessbo Å, Forsberg M, Shokrai A, Rozell B, Engström W (1998) Expression of the insulin like growth factor II gene in polychlorinated biphenyl exposed female mink (*Mustela vison*) and their fetuses. *J Clin Pathol* in press
- Baker J, Liu JP, Robertson EJ, Efstratiadis A (1993) Role of insulin like growth factors in embryonic and postnatal growth. *Cell* 75: 72–82
- Bates P, Fisher R, Ward A, Richardson L, Hill DJ, Graham CF (1995) Mammary cancer in transgenic mice expressing insulin like growth factor II. *Br J Cancer* 72: 1189–1193
- Baxter RC (1996) The role of insulin like growth factors and their binding proteins in tumour hypoglycaemia. *Horm Res* 46: 195–201
- Beato M (1989) Gene regulation by steroid hormones. *Cell* 56: 335–344
- Beck F, Samani NJ, Penschow JD, Thorley B, Tregear CW, Coghlan JP (1987) Histochemical localisation of IGF I and II mRNA in the developing rat embryo. *Development* 101: 175–184.
- Beck F, Samani NJ, Senoir P, Byrne S, Morgan K, Gebhard R, Brammar WJ (1988) Control of IGF II mRNA levels by glucocorticoids in the neonatal rat. *J Mol Endocrinol* 1: R5–R8
- Bol DK, Kiguchi K, Gimenez-Conti I, Rupp T, deGiovanni J (1997) Overexpression of insulin like growth factor I induces hyperplasia, dermal abnormalities and spontaneous tumour formation in transgenic mice. *Oncogene* 14: 1725–1734
- Braulke T, Causin C, Waheed A, Junghans U, Hasilik A, Maly P, Humbel RE, von Figura K (1988) Mannose 6 phosphate-insulin like growth factor II receptor; distinct binding sites for mannose 6 phosphate and insulin like growth factor II. *Biochem Biophys Res Commun* 150: 1287–1293
- Braulke T, Mieskes G (1992) Role of protein phosphatases in insulin like growth factor II (IGF II) stimulated mannose 6 phosphate-IGF II receptor redistribution. *J Biol Chem* 267: 17347–17353
- Cheung PT, Wu J, Banach W, Chernauek SD (1994) Glucocorticoid regulation of an insulin like growth factor binding protein 4 protease produced by a rat neuronal cell line. *Endocrinol* 135: 1328–1335
- Chevalley T, Strong DD, Mohan S, Baylink D, Linkhart TA (1996) Evidence for a role for insulin like growth factor binding protein in glucocorticoid inhibition of normal human osteoblast like cell proliferation. *Eur J Endocrinol* 134: 591–601
- Clemmons DR (1992) IGF binding proteins. Regulation of cellular actions. *Growth Regul* 2: 80–87
- Cohick WS, Clemmons DR (1993) The insulin like growth factors. *Ann Rev Physiol* 55: 131–153
- Coleman ME, deMayo F, Yin KC, Lee HM, Geske R, Montgomery C, Schwartz RJ (1995) Myogenic vector expression of insulin like growth factor I stimulates muscle cell differentiation and myofiber hypertrophy in transgenic mice. *J Biol Chem* 270: 12109–12116
- Concover C, Dollar LA, Hintz RL, Rosenfeld RG (1983) Insulin like growth factor I/somatomedin C and glucocorticoids synergistically regulate mitosis in competent human fibroblasts. *J Cell Physiol* 116: 191–197

- Concover CA, Hintz RL, Johnson BF (1995) Differential effects of glucocorticoids on insulin like growth factor I action on cultured human fibroblasts. *J Cell Physiol* 163:615–622
- Concover CA, Lee PD, Riggs BL, Powell DR (1996) Insulin like growth factor binding protein 1 expression in cultured human bone cells. Regulation by insulin and glucocorticoids. *Endocrinol* 137: 3295–3301
- Dafgård E, Engström W, Larsson O, Zetterberg A (1986) The effects of factors released from tumour transformed cells on DNA synthesis, mitosis and cellular enlargement in 3T3 fibroblasts. *J Cell Physiol* 132: 295–302
- Dahlman-Wright K (1991) DNA binding by the glucocorticoid receptor. A structural and functional analysis. PhD thesis Karolinska Institutet.
- Dalle M, Pradier P, Delost P (1985) The regulation of glucocorticoid secretion during the perinatal period. *Reprod Nutr Dev* 25: 977–991
- Daughaday WH, Rotwein P (1989) Insulin like growth factors I and II. Peptide messenger RNA and gene structures, serum and tissue concentrations. *Endocrine Rev*. 10: 69–91
- deChiara TM, Efstratiadis A, Robertson EJ (1990) A growth deficiency phenotype in heterozygous mice carrying an insulin like growth factor II gene disrupted by gene targeting. *Nature* 345: 78–80
- de Meyts P, Wallach B, Christoffersen CT, Urso B, Gronskov K, Latus LJ, Yakushiji F, Ilondo MM, Shymko RM (1994) The insulin like growth factor I receptor. Structure, ligand binding mechanisms and signal transduction. *Horm Res* 42: 152–169
- Drop SL, Schuller AG, Lindebergh-Kortleve DJ, Groffen C, Brinkman A, Zwarthoff EC (1992) Structural aspects of the IGFBP family. *Growth Regul* 2: 69–79
- Eggenchwiler J, Ludwig T, Fisher P, Leighton PA, Tilghman SM, Efstradiatis A (1997) Mouse mutant embryos expressing IGF II exhibit phenotypic features of the Beck with Wiedemann and Simpson Golabi Behmel Syndromes. *Genes Dev* 11: 3128–3142
- Filson AJ, Louvi A, Efstradiatis A, Robertson EJ (1993) Rescue of the T-associated maternal effect in mice carrying null mutations in *Igf-2* and *Igf-2r*, two reciprocally imprinted genes. *Development* 118: 731–736
- Florini JR, Magri KA, Ewton DZ, James PL, Grindstaff K, Rotwein P (1991) Spontaneous differentiation of skeletal myoblasts is dependent upon autocrine secretion of insulinlike growth factor II. *J Biol Chem* 266: 15917–15923
- Forhead AJ, Li J, Gilmour RS, Fowden AL (1998) Control of hepatic insulin like growth factor II gene expression by thyroid hormones in fetal sheep near term. *Am J Physiol* 275: E149–E156
- Gabbitas B, Canalis E (1996) Cortisol enhances the transcription of insulin like growth factor binding protein 6 in cultured osteoblasts. *Endocrinol* 137: 1687–1692
- Gabbitas B Canalis E (1997) Growth factor regulation of insulin like growth factor binding protein 6 expression in osteoblasts. *J Cell Biochem* 66: 77–86
- Giudice LC (1992) Insulin like growth factors and ovarian follicular development. *Endocrine Rev* 13: 641–649
- Goswami R, Lacson R, Yang E, Sam R, Unterman (1994) Functional analysis of glucocorticoid and insulin response sequences in the rat insulin like growth factor binding protein 1 promoter. *Endocrinol* 134: 736–743
- Gowan L, Hapton B, Hill DJ, Schlueter RJ, Perdue J (1987) Purification and characterisation of a unique high molecular weight form of insulin like growth factor II. *Endocrinol* 121: 449–458
- Granerus M, Pettersson E, Gustavsson L, Lake M, Tally M, Schofield PN, Engström W (1993) Growth Factors in early embryogenesis. *Reprod Dom Anim*. 28: 176–182
- Granerus M, Bierke P, Zumkeller W, Smith J, Engström W, Schofield PN (1995) Insulin like Growth Factor II prevents apoptosis in a human teratoma derived cell line. *J Clin Pathol* 48: M153–M157
- Granerus M, Engström W (1996) Growth Factors and Apoptosis. *Cell proliferation* 29: 309–314
- Granerus M, Johannisson A, Ekblom P, Engström W (1998) Manuscript in preparation
- Heck S, Kullmann M, Gast A, Ponta H, Rahmsdorf HJ, Herrlich P, Cato ACB (1994) A distinct modulating domain in glucocorticoid receptor monomers in the repression of activity and activity of the transcription factor AP-1. *EMBO J* 13: 4087–4095
- Hembree JR, Harman CS, Nevins TD, Eckert RL (1996) Regulation of human dermal papilla cell production of insulin like growth factor binding protein by retinoic acid, glucocorticoids and insulin like growth factor I. *J Cell Physiol* 167: 556–561
- Hirai K, Miyamasi M, Yamaguchi M, Nakajima K, Ohtoshi T, Koshino T, Takaishi T, Morita Y, Ito K (1993) Modulation of human basophil histamin release by insulin like growth factors. *J Immunol* 150: 1503–1508
- Hyldahl L, Engstrom W, Schofield PN (1986) Stimulatory effects of insulin like growth factors on DNA-replication in the human embryonic cornea. *J Embryol Exptl Morphol* 98: 71–83
- Jonat C, Rahmsdorf HJ, Park KK, Cato ACB, Gebel S, Ponta H, Herrlich P (1990) Antitumor promotion and antinflammation. Down modulation of AP-1 activity by glucocorticoid hormone. *Cell* 62: 1189–1204
- Jones JL, Clemmons DR (1995) Insulin like growth factors and their binding proteins. *Biological actions*. *Endocrine Rev* 16: 3–34
- Jux C, Leiber K, Hugel U, Blum W, Ohlsson C, Klaus G, Mehls O (1998) Dexamethasone impairs growth hormone stimulated growth by suppressing local insulin like growth factor I production and expression of GH and IGF I receptors in cultured rat chondrocytes. *Endocrinol* 139: 3296–3305
- Katz LE, Bhala A, Camron E, Nunn SE, Hintz RL, Cohen P (1997) IGF II, IGF binding proteins and IGF receptors in pancreatic beta cell lines. *J Endocrinol* 152: 455–464
- Kitriki E, Philippidis H, Stylianopoulou F (1992) Hormonal control of insulin like growth factor II gene expression in the rat liver. *J Mol Endocrinol* 9: 131–136
- Levinovitz A, Norstedt G (1989) Developmental and steroid hormonal regulation of insulin like growth factor II expression. *Mol Endocrinol* 3: 797–804
- Li J, Saunders JC, Fowden AL, Dauncey MJ, Gilmour RS (1998) Transcriptional regulation of insulin like growth factor II gene expression by cortisol in fetal sheep during late gestation. *J Biol Chem* 273: 10586–10593
- Lindenberg-Kortleve DJ, Rosato RR, van Neck JW, Nauta J, von Kleffens M, Groffen C, Zwarthoff EC, Drop SL (1997) Gene expression of the insulin like growth factor system during mouse kidney development. *Mol Cell Endocrinol* 132: 81–91
- Liu JP, Baker J, Perkins AS, Robertson EJ, Efstradiatis AJ (1993) Mice carrying null mutations of the genes encoding like growth factor I and type I receptor. *Cell* 75: 59–72
- Liu J, Kahri AL, Heikkila P, Blum WF, Voutilainen R (1994) Glucocorticoids increase insulin like growth factor II mRNA accumulation in cultured humanphaeochromocytoma cells. *J Endocrinol* 142: 29–35
- Lu F, Han VK, Milne WK, Fraser M, Carter AM, Berdusco ET, Challis JR (1994) Regulation of insulin like growth factor II gene expression in the ovine fetal adrenal gland by adrenocorticotrophic hormone and cortisol. *Endocrinol* 134: 2628–2635
- Lucibello FC, Slater WEP, Jooss KU, Beato M, Muller R (1990) Mutual transrepression of fos and the glucocorticoid receptor. Involvement of a functional domain in fos which is absent in fos B. *EMBO J* 9: 2827–2834
- Luo JM, Murphy LJ (1990) Regulation of insulin like growth factor binding protein 3 expression by dexamethasone. *Mol Cell Endocrinol* 74: 213–219
- Madej A, Einarsson S, Romanowicz K, Forsberg M, Tsuma VT, Engström W, Barcikowski B. (1996). Simulated stress and the biochemical, endocrine and reproductive consequences in the

- pig. *Reprod Dom Animals* 31: 565–569
- Matthews LS, Hammer RE, Behringer RR, d'Ercole J, Bell GI, Brinster RL, Palmiter RD (1988) Growth enhancement of transgenic mice expressing human insulin like growth factor I. *Endocrinol* 123: 2827–2833
- McCusker RH, Camacho-Hubner C, Bayne ML, Cascieri MA, Clemmons DR (1990) Insulin like growth factor binding to human fibroblast and glioblastoma cells: The modulating effect of cell released IGF binding proteins. *J Cell Physiol* 144: 244–253
- McEwan IJ, Wright AP, Gustafsson JÅ (1997) Mechanism of gene expression by the glucocorticoid receptor. Role of protein-protein interactions. *Bioessays* 19: 153–160
- Mesiano S, Katz SL, Lee JY, Jaffe RB (1996) Insulin like growth factors augment steroid production and expression of steroidogenic enzymes in human fetal adrenal cortical cells. Implications for adrenal androgen regulation. *J Clin Endocrinol Metab* 82: 1390–1396
- Miell JP, Buchanan CR, Norman MR, Maheshvari HG, Blum WF (1994) The evolution of changes in immunoreactive serum insulin like growth factors, IGF binding proteins, circulatory growth hormone (GH) and GH-binding proteins as a result of short term dexamethasone treatment. *J Endocrinol* 142: 547–554
- Minititi CP, Kohn EC, Grubb JH, Sly WS, Oh Y, Muller HL, Rosenfeld RG, Helman LJ (1992) The insulin like growth factor II (IGF II)-mannose 6 phosphate receptor mediates IGF II induced motility in human rhabdomyosarcoma cells. *J Biol Chem* 267: 9000–9004
- Miura Y, Higashi Y, Kato H, Takahashi S, Nagushi T (1992) Effects of dexamethasone on the production of insulin like growth factor binding proteins in primary cultures of rat hepatocytes. *Biosci Biotechnol Biochem* 56: 1396–1400
- Morgan DO, Edman JC, Standing DN, Fried VA, Smith MC, Roth RA, Rutter WJ (1987) Insulin like Growth Factor II receptor as a multifunctional binding protein. *Nature* 329: 301–307
- Morrione A, Valentinis B, Xu S, Yumet G, Louvi A, Efstratiadis A, Baserga R (1997) Insulin like growth factor II stimulates cell proliferation through the insulin receptor. *Proc Natl Acad Sci USA* 94: 3777–3782
- Mouhedianne OB, Cazals V, Kuto E, le Bouc Y, Clement A (1996) Glucocorticoid induced growth arrest of lung alveolar epithelial cells is associated with increased production of insulin like growth factor binding protein 2. *Endocrinol* 137: 287–295
- Nissley SP, Kiess M (1991) Binding of IGF II and lysozymal enzymes to the IGF II/ Mannose 6 phosphate receptor. In "Modern Concepts of Insulin like Growth Factors". Ed by EM Spencer, Elsevier, New York, pp 419–430
- Orlowski CC, Chernašek SD, Åkeson R (1989) Actions of insulin like growth factor 1 on the B104 neuronal cell line: Effects on cell replication, receptor characteristics and influence of secreted proteins on ligand binding. *J Cell Physiol* 139: 469–476
- Orlowski CC, Ooi GT, Rechler MM (1990) Dexamethasone stimulates transcription of the insulin like growth factor binding protein 1 gene in H4 II E rat hepatoma cells. *Mol Endocrinol* 4: 1592–1599
- Price WA, Stiles AD, Moats-Staats BM, d'Ercole AJ (1992) Gene expression of insulin like growth factors (IGFs), the type 1 IGF receptor and IGF binding proteins in dexamethasone induced fetal growth retardation. *Endocrinol* 130: 1424–1432
- Reiss K, Cheng W, Ferber A, Kajstura J, Li P, Li B, Olivetti G, Homey CJ, Baserga R, Anversa P (1996) Overexpression of insulin like growth factor I in the heart is coupled with myocyte proliferation in transgenic mice. *Proc Natl Acad Sci USA* 93: 8630–8635
- Rinderknecht E, Humbel RE (1978a) Primary structure of human insulin like growth factor II. *FEBS letters* 89: 283–286
- Rinderknecht E, Humbel RE (1978b) The amino acid sequence of human insulin like growth factor I and its structural homology with proinsulin. *J Biol Chem* 253: 2769–2776
- Robertson DG, Marino EM, Thule PM, Seneviratne CK, Murphy LJ (1994) Insulin and glucocorticoids regulate IGF BP-1 expression via a common promoter region. *Biochem Biophys Res Comm* 200: 226–232
- Rogler CE, Yang D, Rosetti L, Donohoe J, Alt E, Chang CJ, Rosnfeld RG, Nelly K, Hintz R (1994). Altered body composition and increased frequency of diverse malignancies in insulin like growth factor II transgenic mice. *J Biol Chem* 269: 13779–13784
- Rossetti L, Barzilai N, Chen W, Harris T, Yang D, Rogler CE (1996) Hepatic overexpression of insulin like factor II in adulthood increases basal and insulin stimulated glucose disposal in conscious mice. *J Biol Chem* 271: 203–208
- Salmon WD, Daughaday WH (1957) A hormonally controlled serum factor which stimulates sulphate incorporation by cartilage *in vitro*. *J Clin Lab Med* 49: 825–836
- Schofield PN, Lee , Hill DJ, Cheetham JE, James D, Stewart C (1991) Tumour suppression associated with expression of human insulin like growth factor II. *Br J Cancer* 63: 687–692
- Schofield PN, Tally M, Engström W (1990) Growth Factor synthesis by a human teratocarcinoma cell line. Implications for autocrine growth in the human embryo. In "Activation of hormone and growth factor receptors" Ed by MN Alexis and CE Sekeris, Kluwer Academic Publishers, pp 49–59
- Schofield PN (1992) The Insulin like Growth Factors. Oxford University Press
- Schofield PN, Granerus M, Tally M, Engstrom W (1994) The biological effects of a high molecular weight form of IGF II in a pluripotential human teratocarcinoma cell line. *Anticancer Research* 14: 533–538
- Schule R, Rangarajan P, Kliever S, Ransone LJ, Bolado J, Yang N, Verma IM, Evans RM (1990) Functional antagonism between oncoprotein c-jun and the glucocorticoid receptor. *Cell* 62: 1217–1226
- Scott J, Cowell J, Robertson ME, Priestley J, Wadey R, Hopkins B, Graham CF (1985) Insulin like growth factor II gene expression in Wilms tumour and embryonic tissues. *Nature* 317: 260–262
- Shimasaki S, Ling N (1991). Identification and molecular characterisation of insulin like growth factor binding proteins (IGFBP 1, 2, 3, 4, 5, 6). *Prog Growth Factor Res* 3: 243–266
- Steele-Perkins G, Turner J, Edman JC, Hari J, Pierce SB, Stover C, Rutter WJ, Roth RA (1988) Expression and characterisation of a functional human insulin like growth factor I receptor. *J Biol Chem* 263: 11486–11492
- Suh DS, Zhou YH, Ooi GT, Rechler MM (1996) Dexamethasone stimulation of rat insulin like growth factor binding protein I promoter activity involves multiple cis-elements. *Mol Endocrinol* 10: 1227–1237
- Suh DS, Rechler MM (1997) Hepatocyte nuclear factor 1 and the glucocorticoid receptor synergistically activate transcription of the rat insulin like growth factor binding protein 1 gene. *Mol Endocrinol* 11: 1822–1831
- Suwanichkul A, Allander SV, Morris SL, Powell DR (1994) Glucocorticoids and insulin regulate expression of the human gene for insulin like growth factor binding protein 1 through proximal promoter elements. *J Biol Chem* 269: 30835–30841
- Swolin D, Branting C, Matejka G, Ohlsson C (1996) Cortisol decreases IGF I mRNA levels in human osteoblast like cells. *J Endocrinol* 149: 397–403
- Urban RJ, Bodenbun YH, Nagamani M, Pierce J (1994) Dexamethasone potentiates IGF I actions in porcine granulosa cells. *Am J Physiol* 267 (1): E115–E123
- van Buul-Offers SC, de Haan K, Reijnen-Gresnigt MG, Meinsma D, Jansen M, Oei SL, Bonte EJ, Sussenbach JS, van der Brande JL (1995) Overexpression of human insulin like growth factor II in transgenic mice causes increased growth of the thymus. *J Endocrinol* 144: 491–502
- Wang ZQ, Fung MR, Barlow DP, Wanger EF (1994) Regulation of

- embryonic growth and lysosomal targeting by imprinted IGF2-MPR genes. *Nature* 372: 464–467
- Ward A, Bierke P, Pettersson E, Engström W (1994) Insulin like growth factors-Growth, transgenes and imprinting. *Zool Sci* 11: 167–174
- Werner H, Stannard B, Bach MA, Roberts CT, LeRoith D (1991) Regulation of the insulin like growth factor I receptor gene in normal and pathological states. *Adv Exp Med Biol* 293: 263–272
- Villafuerte BC, Koop BL, Pao CI, Phillips LS (1995) Glucocorticoid regulation of insulin like growth factor binding protein 3. *Endocrinol* 136: 1928–1933
- Wright AP, Zilliacus J, McEwan IJ, Dahlman Wright K, Almlöf T, Carlstedt-Duke J, Gustafsson JÅ (1993) Structure and function of the glucocorticoid receptor. *J Ster Biochem* 47: 11–19
- Yang Yen HF, Chambard JC, Sun YL, Smeal T, Schmidt TJ, Drouin J, Karin M (1990) Transcriptional interference between c-jun and the glucocorticoid receptor. Mutual inhibition of DNA binding due to direct protein-protein interaction. *Cell* 62: 1205–1215
- Yoshiko Y, Hirao K, Maeda N (1998). Dexamethasone regulates the actions of endogenous insulin like growth factor II during myogenic differentiation. *Life Sci* 63: 77–85
- Yuan W, Lucy MC, Smith MF (1996) Messenger ribonucleic acid for insulin like growth factors I and II, insulin like growth factor binding protein 2, gonadotropin receptors and steroidogenic enzymes in porcine follicles. *Biol Reprod* 55: 1045–1054
- Zetterberg A, Engstrom W, Dafgård E (1984): The relative effects of different types of growth factors on DNA-replication, mitosis and cellular enlargement. *Cytometry* 5: 368–375
- Zetterberg A, Engström W, Larsson O (1982) Growth activation of resting cells. *Ann N Y Acad Sci* 397: 130–147
- Zhang Q, Tally M, Larsson O, Kennedy RT, Huang L, Hall K, Berggren PO (1997) Insulin like growth factor II signalling through the insulin like growth factor II-mannose 6 phosphate receptor promotes exocytosis in insulin secreting cells. *Proc Natl Acad Sci USA* 94: 6232–6237
- Zilliacus J, Carlstedt-Duke J, Gustafsson JÅ, Wright AP (1994) Evolution of distinct DNA binding specificities within the nuclear receptor family of transcription factors. *Proc Natl Acad Sci USA* 91: 4175–4179

(Received October 29, 1998 / Accepted March 8, 1999)