



## **Cloning of the Genes for the Pituitary Glycoprotein Hormone $\alpha$ and Follicle-Stimulating Hormone $\beta$ Subunits in the Japanese Crested Ibis, *Nipponia nippon***

Authors: Kawasaki, Daisuke, Aotsuka, Tadashi, Higashinakagawa, Toru, and Ishii, Susumu

Source: Zoological Science, 20(4) : 449-459

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.20.449>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Cloning of the Genes for the Pituitary Glycoprotein Hormone $\alpha$ and Follicle-Stimulating Hormone $\beta$ Subunits in the Japanese Crested Ibis, *Nipponia nippon*

Daisuke Kawasaki<sup>1\*</sup>, Tadashi Aotsuka<sup>2</sup>, Toru Higashinakagawa<sup>1</sup>,  
and Susumu Ishii<sup>2</sup>

<sup>1</sup>Department of Biology, School of Education, Waseda University, Tokyo 169-8050, Japan

<sup>2</sup>Department of Biological Science, Graduate School of Science, Tokyo Metropolitan University, Tokyo 192-0397, Japan

**ABSTRACT**—We have isolated a part of the gene for the pituitary glycoprotein hormone common  $\alpha$  subunit (PGH $\alpha$ ) and the whole gene for the follicle-stimulating hormone  $\beta$  subunit (FSH $\beta$ ) in the Japanese crested ibis (*Nipponia nippon*), a critically endangered bird species in East Asia. The nucleotide sequence of a part of the PGH $\alpha$  gene (5026 bp) contained three exons holding the whole coding and 3' untranslated regions, but lacked a 5' untranslated region. Its exon-intron structure was similar to that in mammals, but different from that in teleosts in the location of the second intron. For the FSH $\beta$  gene, the nucleotide sequence of 7633 bp was assembled from two phage clones. The exon-intron structure of three exons and two introns was similar to that observed in mammals and teleosts. In the putative promoter region of the ibis FSH $\beta$  gene, a progesterone responsive element (PRE)-like sequence and two AP-1 responsive element-like sequences reported in the ovine FSH $\beta$  gene were not conserved in complete form. The increased number of ATTTA motifs in the putative 3' untranslated region in comparison with those in Japanese quail and chicken FSH $\beta$  cDNA suggested that more rapid degradation of FSH $\beta$  mRNA occurs in this species. Deduced amino acid sequences of the ibis PGH $\alpha$  and FSH $\beta$  showed high similarities with those of the corresponding subunits of other avian species. This is the first report on the genomic sequences of the PGH $\alpha$  and FSH $\beta$  in an avian species.

**Key words:** gonadotropin, PGH $\alpha$ , FSH $\beta$ , gene, the Japanese crested ibis

## INTRODUCTION

Follicle-stimulating hormone (FSH) and luteinizing hormone (LH) are the main members of gonadotropins and they also belong to the pituitary glycoprotein hormone family which also includes thyroid-stimulating hormone (TSH). All these hormones are heterodimeric molecules composed of a common  $\alpha$  subunit and a hormone-specific  $\beta$  subunit. The  $\alpha$  and  $\beta$  subunits are encoded by different genes and synthesized as separate peptides. So far, the genes encoding the gonadotropin subunits have been characterized in mammals and teleosts. As for gonadotropin subunits in avian species, cDNA sequences have been reported in Japanese quail (Ando and Ishii, 1994; Kikuchi *et al.*, 1998), chicken (Noce *et al.*, 1989; Foster *et al.*, 1992; Shen and Yu, 2002),

turkey (Foster and Foster, 1991; You *et al.*, 1995) and two species of duck (Hsieh *et al.*, 2001), and amino acid sequences have been chemically determined in ostrich (Koide *et al.*, 1996). However, genomic information such as exon-intron structure and regulatory sequences in the 5' flanking region has not been available in avian species. Accordingly, cloning of avian gonadotropin subunit genes will provide us important information on the evolution of gonadotropin genes and their regulation.

The Japanese crested ibis *Nipponia nippon* (order Ciconiiformes, family Threskiornithidae) is a critically endangered species in East Asia. This species was formerly widespread throughout Japan, Korea, China and an adjacent part in southeastern Russia. At present, the only wild population of approximately 170 individuals survives in Yang Xian, Shaanxi Province, China. In addition, captive populations are kept in China and Japan. Since 1980s, Ishii and his collaborators (Ishii, 1999; Wingfield *et al.*, 2000) have developed endocrinological methods for the artificial breeding of

\* Corresponding author: Tel. +81-426-77-1111;  
FAX. +81-426-77-2559.  
E-mail: kawasakid@ruri.waseda.jp

endangered birds. They have shown that hormone therapy using gonadotropins from closely related species in reproductively quiescent female Japanese quails resulted in gonadal maturation, egg-laying and production of reproductively active offspring (Wakabayashi *et al.*, 1992; Ishii, 1999). Thus, homologous gonadotropin administration is considered to be effective in stimulating reproductive activity in the ibis. However, chemical isolation of gonadotropins from the ibis or related species for the hormone therapy program is almost impossible. Rather, it seems practical to clone the genes for the gonadotropin subunits of the ibis and to generate large quantities of gonadotropins using the recombinant technique.

Thus, cloning of the genes for the gonadotropin subunits in the Japanese crested ibis is invaluable because it provides not only the first information on the genomic structure of these subunits in avian species, but also the molecular information which is necessary for recombinant gonadotropins. Here, we report the isolation and characterization of a nucleotide sequence of a part of the gene for the pituitary glycoprotein hormone common  $\alpha$  subunit (PGH $\alpha$ ) and the sequence of the whole gene for the follicle-stimulating hormone  $\beta$  subunit (FSH $\beta$ ) in the Japanese crested ibis.

## MATERIALS AND METHODS

### Isolation of the ibis PGH $\alpha$ gene

When the last male Japanese crested ibis of the Japanese origin (named Midori) died in 1995, most of his organs were preserved in liquid nitrogen (Ishii, 1999). Genomic DNA was extracted from approximately 15 mg of the preserved kidney using a GenomicPrep Cells and Tissue DNA Isolation Kit (Amersham Biosciences, NJ). The extracted DNA was digested with *EcoRI* and separated in agarose gel electrophoresis. DNA fragments of 4–10 kb long were recovered from the gel and ligated with the Lambda ZAPII vector (Stratagene, CA). An original library of  $2.5 \times 10^5$  clones was generated and amplified once to  $1.3 \times 10^{11}$  clones. Approximately  $2.5 \times 10^5$  clones in the amplified library were screened by plaque hybridization. To prepare a hybridization probe, the PGH $\alpha$  cDNA fragment of Japanese quail (Ando and Ishii, 1994) was randomly labeled with [ $\alpha$ - $^{32}$ P]dCTP (Amersham Biosciences, UK) and used as a probe. Prehybridization was performed at 42°C for more than 2 hr with denatured salmon sperm DNA (0.2 mg/ml) in a hybridization buffer containing 6 $\times$ SSC (1 $\times$ SSC: 150 mM NaCl, 15 mM sodium citrate, pH7.0), Denhardt's solution (0.02% Ficoll, 0.02% bovine serum albumin, and 0.02% polyvinylpyrrolidone), and 0.1% SDS. Hybridization was carried out at 55°C overnight in the hybridization buffer containing the labeled probe. Membranes were washed once with 1 $\times$ SSC containing 0.1% SDS at 60°C for 20 min and hybridization signals on the membranes were analyzed with a BAS-2000II Bio-Imaging Analyzer (Fuji Photo Film, Japan). After third round of screening, ten positive clones were isolated and the insert fragments were subcloned into the pBluescript phagemid vector. After identification of these clones by restriction enzyme digestion, a clone (pIA1) was selected for sequence analysis. Nucleotide sequence was determined with a Thermo Sequenase Cycle Sequencing Kit (USB Corporation, OH) with a DNA sequencer Model 4000L (LI-COR, NE).

### Isolation of the ibis FSH $\beta$ gene

In order to obtain a nucleotide sequence of a part of the FSH $\beta$  gene, PCR was carried out with genomic DNA as a template. Prim-

ers used were as follows: FSH-F, 5'-TG(T/C)TCIGGITA(C/T)TG(C/T)T(A/T)(A/C)ACIA(A/G)(A/G)G-3' and FSH-R, 5'-CA(A/G)TCIGT(A/G)CT(A/G)TCI(C/G)T(A/G)TC(A/G)CAI(G/T)TI(C/T)C(A/G)CA(A/G)TG(A/G)C-3', for the sense and antisense, respectively. They were originally used for amplifying a part of the FSH $\beta$  cDNA in Japanese quail (Kikuchi *et al.*, 1998). The PCR was performed with 30 cycles of 1 min at 95°C, 1 min at 50°C, 3 min at 72°C using a Premix Ex Taq (Takara Shuzo, Japan). The PCR products were subjected to agarose gel electrophoresis and then cloned into pCR2.1 plasmid vector (Invitrogen, CA). After sequencing, a clone (pIF-PCR) was selected for further study.

Another genomic library was constructed in the Lambda EMBL3 phage vector (Stratagene, CA) with genomic DNA partially digested with *BamHI*. An original library of  $5.0 \times 10^4$  clones was amplified once to  $8.0 \times 10^9$  clones, and  $1.0 \times 10^5$  clones were screened by plaque hybridization. The pIFPCR insert was randomly labeled with [ $\alpha$ - $^{32}$ P]dCTP and used as a probe. Prehybridization was performed as described above, and hybridization was carried out at 60°C overnight. Membranes were washed once with 1 $\times$ SSC containing 0.1% SDS at 60°C for 20 min and once with 0.1 $\times$ SSC containing 0.1% SDS at 60°C for 20 min. After third round of screening, one positive clone (IF1) was isolated. The insert fragment was digested with *EcoRI* and subcloned into pBluescriptII phagemid vector (Stratagene, CA) for sequencing.

Because IF1 was found to cover only a 5' part of the FSH $\beta$  gene, screening was repeated to obtain the remaining 3' part. The pIFPCR insert was digested with *BamHI* and then a fragment of approximately 180 bp was recovered to exclude a part overlapping with IF1. This fragment was randomly labeled with [ $\alpha$ - $^{32}$ P]dCTP and used as a probe. Since the genomic library did not give any positive signal, we constructed other genomic library with the Lambda EMBL3 vector and *BamHI*-digested genomic DNA. The resultant library was  $1.6 \times 10^5$  clones and was screened without amplification. Hybridization was carried out at 50°C overnight, and membranes were washed once with 3 $\times$ SSC containing 0.1% SDS at 50°C for 20 min. Consequently, three positive clones were isolated. Based on identification by restriction enzyme digestion, a clone (IF4) was selected and a region of this insert was subcloned into the pBluescriptII phagemid vector. Nucleotide sequence was determined with a BigDye Terminator v3.0 Ready Reaction Cycle Sequencing Kit and an ABI PRISM 377 DNA Sequencing System (Applied Biosystems, CA).


### Analysis of the second intron in the ibis FSH $\beta$ gene in two individuals

Nucleotide sequences in the second intron of the FSH $\beta$  gene were determined in two individuals. One was Midori of Japanese origin, and the other was Long-Long which had been lent from China to Japan and died in Japan in 1994. The genomic DNA of Long-Long was extracted from the cryo-preserved kidney (Ishii, 1999). Genomic DNA extraction, PCR using the primers FSH-F and FSH-R, cloning and sequencing were performed as described above. After sequencing, the PCR products were digested with *EcoRI* and subjected to electrophoresis.

## RESULTS

### Isolation and nucleotide sequence of the ibis PGH $\alpha$ gene

Ten positive clones were isolated from the genomic library and nucleotide sequence of a clone (pIA1) was determined (Fig. 1). The determined sequence of 5026 bp was compared with PGH $\alpha$  cDNA sequences of turkey (Foster and Foster, 1991), chicken (Foster *et al.*, 1992), Japanese quail (Ando and Ishii, 1994) and two species of duck (Hsieh *et al.*, 2001). Comparison results showed remarkable



**Fig. 1.** Structure and sequencing strategy of a part of the gene for the PGH $\alpha$  in the Japanese crested ibis. Solid and open boxes indicate the coding and the untranslated region of exons, respectively. Lines represent introns and 3' flanking region. Arrows indicate the direction and extent of sequencing. Restriction sites for *EcoRI* are indicated with *E*.

sequence similarities which enabled us to infer the exon-intron structure (Fig. 2). We could recognize three putative exons containing the whole coding region and 3' untranslated region within this sequence, but we were unable to find a sequence corresponding to most of 5' untranslated region. Exon-intron junctions were determined according to the GT-AG rule (Breathnach and Chambon, 1981). Then, amino acid sequence was deduced and aligned with those of PGH $\alpha$  of turkey (Foster and Foster, 1991), chicken (Foster *et al.*, 1992), Japanese quail (Ando and Ishii, 1994), ostrich (Koide *et al.*, 1996), and two species of duck (Hsieh *et al.*, 2001) (Fig. 3). Three amino acid residues in the putative ibis PGH $\alpha$  precursor were different from those in each of the other birds over a signal peptide of 24 residues, while 96 residues in a mature protein were completely identical to those of the other birds except for ostrich (ostrich mature protein differed in three amino acid residues from those of the other birds). Ten cysteine residues forming disulfide bonds and two putative N-linked glycosylation sites were conserved among all these birds. These findings assured that the acquired nucleotide sequence encoded the ibis PGH $\alpha$  precursor molecule.

All the PGH $\alpha$  genes previously reported in mammals and teleosts consist of four exons and three introns (Fiddes and Goodman, 1981; Goodwin *et al.*, 1983; Burnside *et al.*, 1988; Gordon *et al.*, 1988; Kato *et al.*, 1990; Golos *et al.*, 1991; Huang *et al.*, 1992; Suzuki *et al.*, 1995). The exon-intron structure in the ibis PGH $\alpha$  gene was compared with that of carp (Huang *et al.*, 1992), mouse (Gordon *et al.*, 1988), bovine (Goodwin *et al.*, 1983) and human (Fiddes and Goodman, 1981) (Fig. 4). The result indicated that the clone we obtained lacked a 5' part corresponding to the first exon of other vertebrates. By aligning amino acid sequences, locations of the introns in the coding region were compared. The location of the first intron in the part of the ibis PGH $\alpha$  gene was identical with that of the corresponding intron (the second intron) in mammals, but different from that in teleosts with an amino acid residue (Fig. 5), while the location of the second intron (the third in mammals and teleosts) was conserved through the ibis, mammals and teleosts.

### Isolation and nucleotide sequence of the ibis FSH $\beta$ gene

Strategy to obtain the ibis FSH $\beta$  gene is illustrated in Fig. 6. The PCR gave a single band of approximately 800 bp and the determined 780 bp sequence was compared with FSH $\beta$  cDNA sequences of Japanese quail (Kikuchi *et al.*, 1998) and chicken (Shen and Yu, 2002). This sequence was found to contain an intron of 582 bp flanked by partial exons of 23 and 175 bp of the putative FSH $\beta$  gene. A clone (IF1), which was isolated from the genomic library, contained an insert of approximately 10 kb, of which the nucleotide sequence of 4059 bp around the 3' part was determined. Comparison of the sequence with the FSH $\beta$  cDNA sequences of the two bird species revealed that IF1 contained a 5' part of the FSH $\beta$  gene, but lacked its 3' part. Then, three clones were isolated by repeated screening of the other genomic library. A partial nucleotide sequence of a clone (IF4), 3580 bp in size, was compared with the FSH $\beta$  cDNA sequences of the two bird species. IF4 was shown to contain a 3' part of the FSH $\beta$  gene which was missing in IF1.

By collecting the results from the two phage clones (IF1 and IF4) and the PCR fragment (pIFPCR), a sequence of 7633 bp was acquired (Fig. 7). Exon-intron structure was inferred by sequence similarities with the FSH $\beta$  cDNA in Japanese quail and chicken and exon-intron junctions were determined based on the GT-AG rule (Breathnach and Chambon, 1981). Particularly, we inferred the putative transcriptional start site and the polyadenylation site referring to the 5' and 3' end of the chicken FSH $\beta$  cDNA determined by rapid amplification of cDNA end (RACE) method. Finally, the assembled nucleotide sequence was found to contain 2820 bp of the 5' flanking region followed by three exons of 34, 160 and 2549 bp, separated by two introns of 458 and 582 bp. An amino acid sequence was deduced and aligned with those of Japanese quail (Kikuchi *et al.*, 1998), chicken (Shen and Yu, 2002) and ostrich (Koide *et al.*, 1996) (Fig. 8). The ibis amino acid sequence showed remarkable similarities to those of FSH $\beta$  of Japanese quail (94.7%) and chicken (95.4%). High similarity (93.4%) was also observed with the ostrich FSH $\beta$  except for a deletion of five residues in the C-terminus. Twelve cysteine residues and two putative N-linked glycosylation sites were conserved through these four bird species. These results enabled us to conclude that the nucleotide sequence we obtained encoded the FSH $\beta$  gene of the Japanese crested ibis.

The ibis FSH $\beta$  gene showed a structure with three exons and two introns. The similar exon-intron structure has been previously reported in mammals and teleosts (Jameson *et al.*, 1988; Kim *et al.*, 1988; Gharib *et al.*, 1989; Hirai *et al.*, 1990; Guzman *et al.*, 1991; Kumar *et al.*, 1995; Sohn *et al.*, 1998; Rosenfeld *et al.*, 2001). Locations of the first intron in the 5' untranslated region and the second intron between amino acid residue 33 and 34 of mature protein in the ibis FSH $\beta$  gene were also similar to locations reported for mammalian and teleostean counterparts. Comparison of the nucleotide sequence in the 5' flanking region between the ibis and non-avian species (teleosts and mammals)

gaattcattttttaaaaaatcaggcaaaagaatttgggacaatattcgtacagttgcaatttacatatttgcacatctttaagtgtgttc  
 cccagttattcatatcagagatcacgcttaccctttgatctggatcaaatatagccacacaactgcatttctgttgaattaccagaga  
 acatgcaatatgctatttacggccacttcatcttgtgagctagcagtagaaggtgatgtgcagtgctgcatgtagcctctttcaciaa  
 aattaatctattgaaaaaatttgtaaaagtttgagctgactggggtgtgttaccagcaccacaaaagtcaattatttgcgtgatcag  
 gaaatgaagcctgaaagaaatcttggtccagaaggctgcagagatgctgaactcccaggaccacctgacctgagctgcctgacga  
 ggctccatgcatgaggtggcccccagggtgtttcaagccctctgagctctctcagcaccgtgtccctcgggtgctctggagca  
 cagcatatagtcctgtctaatatacagggtcagaggagcttttggcactgaggacaggactgggaaaatctgtttatggggactttg  
 atgctaaagcagtcagtggtgaacaaaataatctgactcttggttcaggaaattatgtctgatgtgaaattatttcccttaaatag  
 ctttcatggctcagattctgcagattctctatctcctccataagcatttctggtcttatccagctaccgtaaatattaggcagttat  
 gacgtttgaagttatgtgctgtgtttatgttaaatattcagaaaaaactgttaaggccagaagaaggcaaatattgttatagtggt  
 gaagtcaattttctgagcctataaaatgcacatcagtagcactgcaatttctaatctgttctaaattgtctctgttacttgcg  
 ctgaatgctctcagtgctgtagctagatgctgttactgtacccaactgcaaacagctoccttaaaaaatgtatttattgtattccagA  
 AAGATC ATG GGT TGC TAC GGG AAG TAT GCA GCT GTC ACT TTT G ACC ATT TTG TCT GTA TTT CTG CAT CTT  
 Met Gly Cys Tyr Gly Lys Tyr Ala Ala Val Thr Leu Thr Ile Leu Ser Val Phe Leu His Leu  
 -20 -10

CTT CAT GCT TTC CCA GAT GGA GAG TTT CTC ATG CAG Ggtaagctgctttcagcattcagaaatagggcaattgttcgt  
 Leu His Ala Phe Pro Asp Gly Glu Phe Leu Met Gln  
 +1 +9

aaagagtattaataaaatttcacatgtttataaatgcagtcataaatcttatcattcagttgcatttattgacactgagatatgaagg  
 tataataaatcaaggtaatttgcataaatatacttccatttcttctctaaacttgatttctccagggtcaagagcctcacagatgact  
 gtgtcctcatgttgacatgggtaaccttaactctgaacattatcaaatagaacattttggtttcttctgatatcctgagatcctaattg  
 caacgtaacatcaggaccagcagaacctatgctctctcataaatatgcagaaaatttgcctctgtagatccagaatattgggactttg  
 gtgtgtagcattctagaaagagcatatttaggaagatttaggggtgttttggaggttgctgttttttttttaattcagcctcg  
 aactgaatgtagaaaaatgtgagctacaaaaatctcgggatcactggcatttttaaaacttgataacagagctcttgcctgaaaacagac  
 aatgtttaagaacactacaagctcctgtgtgttatattaaccatctgtagctagtaggaagtttctagggacagctagtgtagtagac  
 atcaaaaaaagcctccacaactgtcaaaaaataggttagaaaaccagaaaagatacaatgatgctcgtgccttaccctagaaagcc  
 accctctagttttacagacgggtgcatgttgcaggggtgaaggaccccaagaggagagctttgctgtctgtctttctgttctgcaggg  
 atcaagcagcaaaaatgtggccagtgcttctcttactgcccgcactttccctttctttatgaaaatgaaaatacagctggagaccaagt  
 aaaaggttctcttatatttttatggcag GT TGT CCA GAG TGC AAG CTA GGG GAG AAC AGA TTC TTT TCC AAG  
 Gly Cys Pro Glu Cys Lys Leu Gly Glu Asn Arg Phe Phe Ser Lys  
 +10 +20

CCA GGA GCC CCC ATT TAC CAG TGC ACT GGG TGC TGT TTC TCC CGG GCC TAT CCC ACT CCG ATG AGG  
 Pro Gly Ala Pro Ile Tyr Gln Cys Thr Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Met Arg  
 +30 +40

TCC AAG AAG ACC ATG CTC GTT CCA AAG AAC ATT ACA TCA GAA GCA ACG TGC TGC GTA GCA AAG GCT  
 Ser Lys Lys Thr Met Leu Val Pro Lys Asn Ile Thr Ser Glu Ala Thr Cys Cys Val Ala Lys Ala  
 +50 +60

TTT ACC AAG gtgagctgtgaatgagaccgtttaagccttgtttcaggatgcaatgtttagctgagccagtgaacagaaaaaata  
 Phe Thr Lys  
 +70


gtttttgtcggaggacattggagaactatttgggaaacatccttggcatttatatcctttgctcttgggttctttcttttctaaaga  
 ggaattaaaaatgacagcagcagctgactgtatattcagttgggagtggaaattgatgactgtcactgactcagtgaaactacttt  
 gactcatgacacctgaagttctggtgagcttgggttcccttgttttagcctctcctggatagcaccctccttaagtactatca  
 agtgtgtatttttaattagtcattttccttgtgtcttctcctgcataagcagaagttctctgggaaggtggctgtcaatactatca  
 attacatgtatgggaatgaggaccacaaacctctgtgctcattagatttaattgaggttgtgtttacttttaatagtataaccaagagg  
 aactgtaaatgtgtgcaactatttgcatttaattgctctcctggaaggctctcttaaagagagatctgcctaaagctgactgtttg  
 ggagtgattcagctccagggtgaaacaaacttaatttctgaaataaattggtgagctagttgaaacaagctaaaggtgtctatcctgtggt  
 gatcgaatcagactcattaaatgtgttgcactagaggtcaggtcagttaccactcacagatttgggtgcatatgaaataccctggggt  
 actcaagacatctgtgttcggctggcagacacaggaccataaaaagaccttctggaatggtagctggaggcttgggtgggacctctga  
 tatctcaaatggcagttgacagctctgtgtgagcagctgaacaaagtgtgggtctcactggctgtgatgagagcccagggtacctgact  
 tactatagacacctacagctgaacctggctgaacctggctctcctctgagtagggcatagattcttttctagaccactgacattat  
 acttgcctgaggcagcaactagaatttgggtgagattaaaaataaaaatgactcatcgtgtttgtgtctctgtttttataatcgccgt  
 gtgaaaaactctaagaacacttctgattttcatagattggttacttgcacacttttgaatatcagacacttaagaagtgcacaaatta  
 aggacccaactatgtgcctttgaaacttctcttattgctctgaaagccactgatgtgcagcctgcaggttggccttcagagg  
 ctgttctagatatacgtacccacagagcctgtgtaccttattggttaagaaaaaggcttcagacatctttactcgagaatgtgtttta  
 ctgtctagctaactaattatttcttagtggtgtgtgttttctttttaaag ATT ACC CTT AAG GAC AAT GTG AAG ATA  
 Ile Thr Leu Lys Asp Asn Val Lys Ile  
 +72 +80

GAG AAC CAT ACA GAT TGT CAC TGC AGT ACC TGC TAC TAT CAT AAA TCC TAA AGCCTGTCCTTTTGTAAATGAT  
 Glu Asn His Thr Asp Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser stop  
 +90 +96


CAAGGACAACGGTGAATGGAATATTTGTTTTTCAGCTTTTATAGCACTGTGTGTAATAATCTTATGTTTTCTGATCAAGACACTGGGTAG  
 ACCTTTGAATAAGATGGATGGCTGTTTTATTTCCTCTTTGCTTTCATGCGTTTAAAGTAAAGTTAATTAATTTCCATTAGGGATAAAAATA  
 CAGCACTTGCAATGACAAACCGAAGCTTATTTTTAAAAATAACTGTGTCAGTTAAATCATCaatgttcttgaggaaaaatgattatt  
 taatagcttaaaacagaaaaagattctgctgcagttaatcttagatttcttttttggggagaggagggaaggagggaaggaggga  
 tgggaaagtggatagaaaaagaagcaatcttcccttattcatttagtgatgattttttggaagactcaacatttcttctcctg  
 tttctctcggtttttcttaccctttaggcagaaagtttaacagtgatcatctatgcacagtgctctgtcctctgtgcatctttt

at t t t t c t t t g a a c t t a a t c t c t c t c a t a t a g a t a c a t a a a t g g a t g c a g a a t c a g a c c t t a t a a a a a t t g t a c a g t t t g g t g t a c a  
 g t g g t a g g c t a g a g g a g t g c t a c a c a a t a t t a g c t a t t t a a g g t g t g c t t g a g c a g c a t g g t a t a g g g c t t a g t a a t t c c c a a t a a a  
 c a a a t g g g a a g g t a a g g a a t a t g t t c c t g g g g a a c a g g c t g c c c a g g a a g t g g t t g a g t c c c c a t c c c t g g a g g t a t t t a a a g a c t t  
 g t a g a t a t g g t g c t a g g g a c a t g g t t a g t g g t g g a c t t g g c a g t g c t a g g t t a a t g g t t g g g c t g g a t g a t c t t a a a g g t o t t t t c c a  
 a c c t a a a t g a t t c t g t a t t c t g t a t t c t a t g t t a g t g a g a a g t g g a c c a c a g a g t a t g t a t t c a g a a t a c a t g a c c t t a a t a t c a t a  
 a a c t g g g a t t t a g a g a t a g c a c t t g t t c a a t t g t a g t g g g t t g g a t c g g g g a c t c t g g g a a c t g c t t g c a g g t c t g g a c c c g c t c a  
 t g g a a c c g a a g t g a a g g a a t t c


**Fig. 2.** Nucleotide sequence of the part of the PGH $\alpha$  gene in the Japanese crested ibis. Exons are shown in capital letters, and introns and flanking regions are shown in lowercase letters. The polyadenylation signal, AATAAA, is underlined. Deduced amino acid residues are represented with three letter codes below the nucleotide sequence and numbered sequentially from the N-terminus. The nucleotide sequence data in this figure is available in the EMBL/GenBank/DBJ Data Bank with Accession No. AB089503.




**Fig. 3.** Alignment of amino acid sequences of signal peptides (A) and mature proteins (B) of the PGH $\alpha$  precursor molecules of the Japanese crested ibis, duck (Hsieh *et al.*, 2001), chicken (Foster *et al.*, 1992), Japanese quail (Ando and Ishii, 1994), turkey (Foster and Foster, 1991) and ostrich (Koide *et al.*, 1996). Residues identical to those in the Japanese crested ibis are indicated with dots. Ten conserved cysteine residues and two putative N-linked glycosylation sites are denoted by  $\bullet$  and  $\nabla$ , respectively.



**Fig. 4.** Comparison of the exon-intron structure of the part of the PGH $\alpha$  gene in the Japanese crested ibis with those of PGH $\alpha$  gene of carp (Huang *et al.*, 1992), mouse (Gordon *et al.*, 1988), bovine (Goodwin *et al.*, 1983) and human (Fiddes and Goodman, 1981). Solid and open boxes represent the coding and the untranslated regions of exons, respectively. Solid lines indicate introns and flanking regions. Thin broken lines between boxes link corresponding exons. A thick broken line in the Japanese crested ibis represents part whose nucleotide sequence is not available at present.



**Fig. 5.** Alignment of a part of amino acid sequences of PGH $\alpha$  for comparison of location of the second intron. Locations of the intron are shaded. Numbers at the top represent positions of amino acid residues in the PGH $\alpha$  in the Japanese crested ibis. References are as follows; carp (Huang *et al.*, 1992), salmon (Suzuki *et al.*, 1995), rat (Burnside *et al.*, 1988), mouse (Gordon *et al.*, 1988), bovine (Goodwin *et al.*, 1983), porcine (Kato *et al.*, 1990), rhesus (Golos *et al.*, 1991), and human (Fiddes and Goodman, 1981).



**Fig. 6.** Structure and sequencing strategy of the FSH $\beta$  gene in the Japanese crested ibis. Insert fragments of pIFPCR, IF1 and IF4 are shown with arrows which indicate the direction and extent of sequencing. At the bottom, the structure of ibis FSH $\beta$  gene is shown with exons indicated as boxes. Solid and open boxes represent coding and untranslated regions, respectively. Restriction sites for *Bam*HI are indicated with *B*.

gaattccaggagatggcttcaataattccactgtgacattctgggcaacacctagcattcagcctggaagcataatgacaactgctaaa  
tctaggatgggcaactactgcttaagaactagatagataaaaccactacttaacgcttgaaagtgtgctgtagcatgtgaagaggagcgtt  
cttccctctgacattattccctaattgaaatataaacctttaacatggaaaacatttctatttctcaaatagatacaaaaatcagtaa  
cgcagaacctatgtataatgttcacatatatatgaaaattatagctttctggataaataacatttgctattcccttcacaattaccataa  
ccttttttaaatctaataaggaaataaaacttgaaggagtatgatatagcttagcatacaacatttcccagaggtatgaaactaatctt  
agaagacataacttacacaacatcttgcccatcatgagaagtaattccttagcgtatctttctacaaaagggtgtgggaacaaaatca  
aagtcaaccaagtaacaactcaccaaaattaccactcctcagaggatagcgtggtggttagttgattccagtttgaatttctattacc  
taaaaatgcattatcatatattcacacaatgaatttagcagagctgtagctgggagaaccaactagaactaatgattccagaacaaaat  
gcccttgaagtcagtacaggacgaaatcctcctgttaaaaagtttgcatgtgtttacaatgtggcttcatgagctctccagtgagaaca  
ctctgtgaaagcgttctctgtcaacctccccctcattcccactccccaatcaatcccacatctacgctgctacagctgttcatttcatt  
gtggaacaaaatgatatggattaacaacaatgctgtaaagacagctgggtgggtttttgcttttaatacagaatgatttccctaat  
gtgtgccaagtgtgagggtgagtggtggccagacaaggaactagggagggaagaacagcagtgtaggggctaaacttgactctc  
tgcctgtgaaattaggttttagtgaatgtatgacttctcaaacatgaaacagatcattgcaactgagagattagatatttctgt  
attctcctgcaaatgaagtttattgggttttatttaatttgacttagaaaaggcaagatcctcagtagggatataactaatgtaatcag  
ctttgtctggcatctctggcattaatgatttacagcagctgagaatcttctctgacaagacagatcatgttgacagtaaaagacacaaa  
agacaattatcaaaaatcacagagtttatattagtaagaggaagatgtctgctttcttgactctttatacatttaattgagctgtat  
tctcattcaacaaaattattattacaggatattgcccaggcaagttgtagatcacttcttaatttgggtgataggaaattcattaac  
aaataaaggtattgcagactggatggaaactggaacagatggccagctctgcaagaccaggaaggtctgagacacttaagagctcatt  
tgtaaaagccattttggaagaatcctgtctgcaggacagagagaatttctttcatgataaaaatgctgtatgcttctgctagttttct  
tttgggtgatttctgtaaaagtcagactgaccataatagaacagctatacaaaagcaagactggctgttctgggcagatgatagtaa  
atcagctgctactgtgagtcattttcccagaagtgatgtgaagctcaacaagttatttctctgctcattatctggtgagagggcac  
accgtaactgtgaaatgcagaaaaggcagtgagcaggggaacactattcatagaatcttttattatcatgatgaactagggaaatgca  
ttcaagctgagtttgcaggtttatctgttaaaaatattagcattttccctgtaacatgacaactactgtatcagatcactaacagaaga  
cacagatctggaccctcactgtgacatctttatagtcctttccacaatacagacatcccttacataagtttaaaaaaatgtgacctgt  
gacagacaggaaccaggatgacacatactgggatagaaatctgaattttgaaaagccacagtggtattttaaattggaagaaagtcga  
gttacggttttaacaaaactctcaatactagctttctggaccacagatcaataaacaagtatatatgcagctcactgggtaagacctaat  
tcagagaaaagcaattaagattgataccctgaaaaactcaggaacatgttcttagccaagaagcgtctagcaatatctctgctatgaa  
ttaccaaaggacaacttgcaccccaagcctatcagtttgagggtgacataaaacacagttaaaagacttgaactttctttccaaaa  
ataggcttctcagccttattcaaatgtggcaggttccagctatttctaaatgccactgaatcaaatccagggaacttctaaaaagac  
attgtgacagtaacctcaccatcacttctcagtggtgttttgcctttcttttttgccttaacaagcgttgatctcactgtctggctga  
acattaatccactttccaagtaaccaacgaatgcttctgctgcattaacgctgcaaatcctcaaggtatataattacaccaatttagta

taaagggtgagcaggggctttcttttctacTACAGCTACTACCAGGAAGGCAAGTAGAATACAGGtaagtttctcagatcctgatcta  
gaaatctttttcaaatgcaaacactgtatgtaattaattgcttctgagaatacattatcctttgctcttctctcccttgcctctctt  
tctttctccttaagacaggacacattctctgtatggaaacttagataaaaaatattcattgaaagtggtatggtactctgctattatt  
attctcagagattgctgacattgcctcactcagctgcttggaaaatcaaacctcggtaacaataaacacagatagatttttctcattt  
ctttatctatgttctcactgatgtttctgaaagcttgcattatagaaccacttgaggataatgtgaaggctcaacttctctctatttc  
tataaatatttctgttgaaaataactatacatgtttctctggtatagatacatttttgccttatacacagG ATG AAG ACA GTT

Met Lys Thr Val

-20

AAT TGT TAT GTG CTG TTA TTT TGC TGG AAA GCA ATT TGT TGC AAT AGC TGT CAG CTT ACC AAC ATT  
Asn Cys Tyr Val Leu Leu Phe Cys Trp Lys Ala Ile Cys Cys Asn Ser Cys Gln Leu Thr Asn Ile

-10

+1

ACC ATA GCA GTG GAA AGA GAA GAA TGT GAA TTC TGT ATT ACA GTG AAT GCC ACG TGG TGC TCA GGA  
Thr Ile Ala Val Glu Arg Glu Glu Cys Glu Phe Cys Ile Thr Val Asn Ala Thr Trp Cys Ser Gly

+10

+20

TAC TGC TTC ACG AGG gtgagaatcttaagtttaatttctaaataaactcatttctccagtccttaataaacacatagggcaggatga  
Tyr Cys Phe Thr Arg

+30

gccttgtgaaacaacaggttgagtaacttctgtgctatttgacaggaaagtagatacacagccagctgaaactacaaagcgaagggtgaa  
aagatgccagtaaaattcaggtcttcttaactggctgcacacatggttatattgtataactagagttgaaagtaatacaatctggtatct  
ctgcttctgttcaatttgggttgcaaatggatacaataaagcagcctttagactcctctacttttccaaaatattgaaagaaatcaa  
gtaattgacctgaaagattgtctggtgatggtatgacacaaaatttgccatcagctcctcctgaaaggagaaatctaaggatctaggtta  
aattataagagctctgatgcaaaccttctgatctcagtggttctcccatgggttaattagtttccacatcaggtcctttattttgaa  
aaatttgaattctgctcatacaaaaatttcaaaaacagatatttcttctgttaattttcag GAT CCA GTA TAT AAA TAT CCA

Asp Pro Val Tyr Lys Tyr Pro +34 +40

CCA GTA TCA TCT GTT CAG CAA ACA TGT ACT TTC AAA GAG GTT GTG TAT GAA ACA GTG AAG ATC CCA  
Pro Val Ser Ser Val Gln Gln Thr Cys Thr Phe Lys Glu Val Val Tyr Glu Thr Val Lys Ile Pro

+50

+60

GGC TGT GGT GAC CAT CCC GAA TCT TTT TAT TCG TAC CCA GTA GCT ACT GAG TGC CAT TGT GAG ACC  
Gly Cys Gly Asp His Pro Glu Ser Phe Tyr Ser Tyr Pro Val Ala Thr Glu Cys His Cys Glu Thr

+70

+80

TGC GAC ACT GAC AGC ACT GAC TGC ACC GTG CGA GGG CTG GGG CCA TCC TAC TGT TCT TTC AGT CAG  
Cys Asp Thr Asp Ser Thr Asp Cys Thr Val Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Ser Gln

+90

+100

AAT GGA AGT AAC CAA TGA AGGGTACTTGAGATGGCAGCTTGGCTTTACATGTTCACTTCTAAATAAAAGGTACTGATCGGGCTTA  
 Asn Gly Ser Asn Gln Stop  
 +110  
 AGTGAAGATAATAGGCAAGGCTATTTAGAAACTGCCAAGATTGAAACAAAGATTTTTAAAGGCCAAAATGGAGAGCTACTGACTAAACTT  
 CTCTTCAGGCCTTCCCTACTTATCCCACTCAGTTTCCCTTAAATCATTTTCATATGTCTATAGAACACTGCTTCCGATTCCTTCTGCCCTT  
 ACCTCCTCTTCTCTATTACACCTTCATTCCTTATAGTCTGTATTTCCACTGCCTAGTTCACCTTAGATTATTTCTATGTCTATCCTATG  
 CTTTCAAACAGCTTCCCAATTTCCAAGTCTTTATATATCACCTGCTTCCATTTCTCCTCAGAACGCACCTATTTAAAAAGCCTTGCAGCA  
 CTGGTTTTCTTACAACCAACAGCTATGCCCTCACTTTCTATTAGCTGGACTGTAAACCTCTCTGAGCCGAGATCCTGACTTTTTTTTTTT  
 TTGGCAAATATTTTCATATAGTCAATTATGCTCTGTAATAATAACAGCATGTTATTTACAAGCCAAAATCTATGTTCAAATGTTGAATG  
 TCACCTGAACTCTTGTGCTTTCTCAGTAAGGCTTGTAGTCTAGGCTCAGTGACACAATAACGCCAAATAGCAATGAACTCTAGTTTATGCC  
 TGGCTAECTTGA AAAAAGAATCATCAGCCATCTACTTTCACTGAGTAAACATATCCTAAGTTAGTTAAACAATCTGAAAATTTTGTCTG  
 CATCTCAGAAACACTGGATTTCCCTCTCATTTCTGCAACAAAATGTAAGAATGAGAATTTTTTAAACACTATTTAACATAGGCCGCTCACAA  
 ATTATTTAGGGATGAATCTAGAGGCACCTGATCCAGTAGTGCAAACATGGTGTGATAACATGCCTTGAAAGTAGCTCACAGTAGTCCCT  
 CTGAGAATGGAAGTTTATATAGAGTTCTGACGCAACCTAATTTGCAATGGCCAATATGAAGTTGGGATGGTTGGGAAGCAAGCTGGAGG  
 ACCGGCTGAATCAGAATGAAAATTTATCGCACGTTATTTGGGAAAATTTATCTGAAACCACAGAAATGAAACCTTATCAGGAAAATGAAAAG  
 ATACCGCTCTTTCATCAGAAAACATCACCTACACAAACACAGAAATGGAGAGCAACTGGTTTACGACGAGTTCGGCACAAAGACTTGGCGG  
 TTATATGAGCTAAGTATGAGTCAACAGTATGACACCCTGCGA AAAAGTGGGAATGTAAGGAGTAGAGACTACAAGACACAACCTACTTC  
 CACCATTCTGCTCAACATGACAAGAGTCTCAGCGGAGTTGTTATTTCCAGCTTTGGACACAGAACTTCAAGGCATCGAAAAACGGAGAA  
 AATCCACAGAAGAGTATAAAAATGGAGAGAGGCCATAAAAATATCTTAGGAGAAAAGATTTAATAAAAATGTTGTTGTTTTGCTTACGG  
 GAGGTAACACTTAGAGGGGGGAAAATAGTGGAACTTTTACATGCTTGAAGACTGTTACAAGAGAAAAGGAACAGTCAGTTCTCCATG  
 TTCATAGTGAACAGAACAATAATGAAAGATTGAGAGTAGCCACTAGCAAATCTTTGGAAAGTTTACAGACGCCAGACGCTGAATAAAA  
 TGTTAGGGGGTTTAGGAAGGAGGAGAGAGAGTGGGGTCTGTGGAGTAGTGACCCTACAGGCTCTTCAAGAACACATTTGAACCTGTTGG  
 GAATGACAGAGATAAAGTTAATTTGCCATGAGGAAGGGAATGAAGTAGACGCTCTTGGGGTGCCTGACATGTCTGCTTTTGGACCC  
 TTACAGACATGTTTATTTAAAGGAATTTGTCGGTTACTCTGGATGTCTCTTACAATAGATATTTAACTTATTTCAATTTTACTGCTTTT  
 TTACTGAAAGATATAGATCTCTTAACTTTATTTATAAGATTTATCTGCATGGACATTTTCTTTGGTCTACTGAAGTTTAAAGTGTA  
 AAGTAATAGGTAATCAAATGTCCTCTCACTCATTTCAGTTTTAAACATGAGTTGACGTTGGTTAAACTAAGCATAATTATTAATTCAG  
 GAATTTCCATTTCTGGATGAGGGAATTTGAATTAATAATGCAAAACAGAGTAGCTTTGTAGTAAAAAAGATGGA AAAAGATACGCTCT  
 CAAAATAGATCTGTAACACCTGTATTGCTGCTGTGCATTTTGAACAATAAAAATAATTTTGGTGCCTTCTGTTATGTGCTTaaata  
 agccaattaaaaaccttataagtccacctgcctacttcaatgctttaaatcacacttccccccacacacataatttttttcatataa  
 ctaatacactcatctgagccatatttgcaattatagacctgcagtactgtgtccagctctggggtcctcagcactataaagacatggacct  
 gttggagcgggtccagagggggccacaaaatgatcagagggctggaacacctctgctatgaggaaaggctgaaagagctggggctctt  
 tagcctggagacaagaaggctgccccgagaccttattgcccctttcaatacttaagggggcttataagaaagatggggagcagactttt  
 tagcagggcccaatagcgaataggacgaggggtaatgggttttaactaaaagagggtagattcagactagatataaggaagaaacttttac  
 aatgagggtagcgaacactggaacgggtgtccagagaggtggttagatgccccacttggaaacattcaaggacagggctggatggggc  
 tctgagcaacctgatctagttgtgatgtccctgccaatgacagggggatggacttagatgaccttaagagctcccttccatcccaaacg  
 attctatgattttatattgcccacatcaataaaagcctggatatttttagcctgaccagtaataagatggtgaaatattatgta  
 gttgtaacacaacagaactgtggatgagaataaagattatcagtaaatattgtatttaactatagggatagaacctctccagagctgtg  
 tccaatctctgtcacaacttagctgtgataaaacaaaaattatagctctcaatttccatccatcagacaaatgacacacactcattt  
 tcattttgcaaatgaatctattcagaaccttggaaataaaactgtaactaatgcagctcaaaagagatttgaggtaatctcagaatgctg  
 gcaaaaaaatcctaatcagcagcactgcaagctt

**Fig. 7.** Nucleotide sequence of the FSH $\beta$  gene and its flanking region in the Japanese crested ibis. Exons are shown in capital letters, and introns and flanking regions are shown in lowercase letters. The two potential TATAAA sequences and the five potential polyadenylation signals, AATAAA, are underlined. The bent arrow indicates the putative transcriptional start site deduced from 5' end of chicken FSH $\beta$  cDNA (Shen and Yu, 2002). Eight ATTTA motifs in the putative 3' flanking region are boxed. Deduced amino acid residues are represented with three letter codes below the nucleotide sequence and numbered sequentially from the N-terminus. The nucleotide sequence data in this figure is available in the EMBL/GenBank/DDBJ Data Bank with Accession No. AB089502.

revealed that the ibis sequence could be well aligned with the sequences in mammals within the promoter region (Fig. 9).

**Analysis of the second intron of the ibis FSH $\beta$  gene in two individuals**


Results of sequencing the PCR fragments of Midori and Long-Long showed that two types of sequences were present in the second intron of the FSH $\beta$  gene (Fig. 10). The difference between the two types was the 32 bp sequence in the 5' portion. Ten and five clones were sequenced for Midori and Long-Long, respectively. Only the shorter type was observed in Midori, whereas both the shorter (in two clones) and longer (in three clones) types were found in Long-Long. In addition to the length difference, three nucleotide substitutions were observed. A substitution, which was located in the most 3' position, changed the sequence

“GAATTC” (the recognition site of *EcoRI*) in the shorter type into “CAATTC” in the longer type. Utilizing this sequence difference, we could differentiate the two types by *EcoRI* digestion of the PCR products. Namely, the PCR product containing only the shorter type (approximately 780 bp) was divided into two fragments of 550 and 230 bp, while the PCR product of the longer type (approximately 810 bp), was not cleaved. The digestion results showed that only two fragments of 550 and 230 bp were observed for Midori and three fragment of 810, 550 and 230 bp for Long-Long, indicating that Midori had only the shorter type and Long-Long had both types.

**DISCUSSION**

In the present study, we have isolated and characterized a nucleotide sequence of a part of the PGH $\alpha$  gene and





**Fig. 8.** Alignment of the amino acid sequences of signal peptides (A) and mature proteins (B) of FSH $\beta$  precursor molecules of the Japanese crested ibis, chicken (Shen and Yu, 2002), Japanese quail (Kikuchi *et al.*, 1998), and ostrich (Koide *et al.*, 1996). Residues identical to those in the Japanese crested ibis are indicated with dots. Twelve conserved cysteines and two putative N-linked glycosylation sites are denoted by  $\blacktriangledown$  and  $\bullet$ , respectively.

the sequence of the entire FSH $\beta$  gene in the Japanese crested ibis. This is the first isolation and characterization of gonadotropin subunit genes in avian species. The exon-intron structure of the ibis PGH $\alpha$  and FSH $\beta$  genes was basically similar to that of the corresponding gene in mammals and teleosts. This result suggests that the same exon-intron structure exists generally through all avian species. However, there was a minor difference among animal groups. It was the location in the second intron (corresponding to the first intron in the partial ibis gene) in the PGH $\alpha$  gene. This intron was shifted with an amino acid residue in teleosts compared to the ibis and mammals (Fig. 5). It is difficult to conclude from this fact that the teleostean type is more primitive than avian and mammalian type, since Arai *et al.* (1998) reported that most teleosts are considered to be genetically highly specialized group. Arai *et al.* (1998) aligned amino acid sequences of PGH $\alpha$  precursors in various vertebrates and revealed that only Australian lungfish has an insertion of an amino acid residue (alanine) around a position corresponding to the second exon-intron boundary. At present, the genomic structure of lungfish PGH $\alpha$  gene has not been clarified, it may provide some information to explain the location difference of this intron.

In the ibis FSH $\beta$  gene, we deduced the location of the putative transcriptional start site and the polyadenylation site referring to the 5' and 3' end of the chicken FSH $\beta$  cDNA (Shen and Yu, 2002). However, two TATAAA sequences in

the putative promoter region and five potential polyadenylation signals in the putative 3' untranslated region were observed (Fig. 7). The proximal TATAAA sequence, which was 27 bp apart from the putative transcriptional start site, was conserved through the ibis and mammals, whereas the distal sequence was unique to the ibis and did not exist in mammals (Fig. 9). Then, the proximal sequence is considered to be more possible than the distal one, but the other possibility that the distal sequence is utilized cannot be excluded. Further studies are required to obtain a definite conclusion of the transcriptional start site and the polyadenylation site of the ibis FSH $\beta$  gene.

Kikuchi *et al.* (1988) and Shen and Yu (2002) have reported that five ATTTA motifs were found in the 3' untranslated region of the FSH $\beta$  cDNA in Japanese quail and chicken, respectively. This motif has been characterised as a signal for rapid degradation of mRNA coding for cytokine or proto-oncogenes (Shaw and Kamen, 1986; Chen *et al.*, 1995). We found eight ATTTA motifs in the putative 3' untranslated region of the ibis FSH $\beta$  gene. Akashi *et al.* (1994) have demonstrated that the number of this motif in the 3' untranslated region correlates with instability of mRNA using the chimeric rabbit  $\beta$ -globin gene. It is possible that the increased number of this motif in the ibis FSH $\beta$  gene induces more rapid degradation of the mRNA than the other birds.

The nucleotide sequence in the 5' flanking region of the ibis FSH $\beta$  gene showed no appreciable similarity to those in teleosts and mammals apart from those in the promoter region in mammals (Fig. 9). In the promoter region of the ovine FSH $\beta$  gene, a progesterone responsive element (PRE)-like sequence and two AP-1 responsive element-like sequences have been demonstrated to be responsible for progesterone and gonadotropin-releasing hormone (GnRH) regulation of the gene expression, respectively (Webster *et al.*, 1995; Huang *et al.*, 2001). In the bovine and porcine FSH $\beta$  genes, these sequences were completely identical, but several substitutions occurred in each of the sequences in the rat and human FSH $\beta$  genes. Function of these sequences in rat and human has not been clear. The corresponding sequences in the ibis had several substitutions from the mammalian sequences and the consensus sequence. These results suggest a possibility that the ibis and other vertebrates (teleosts and mammals) apply different regulation mechanisms to the FSH $\beta$  gene expression. Further analysis on the 5' flanking region will reveal the mechanism of ibis FSH $\beta$  gene expression.

Among avian species we compared, the amino acid sequences of the PGH $\alpha$  and FSH $\beta$  showed remarkable similarities. According to the taxonomy, extant birds are classified into two lineages, the Palaeognathae and the Neognathae. The Palaeognathae includes ostrich, and the Neognathae consists of two clades; the first clade contains two orders of Galliformes (chicken, Japanese quail and turkey) and Anseriformes (duck), and the second clade contains the rest orders (van Tuinen *et al.*, 2000). Our study has



**Fig. 9.** Comparison of nucleotide sequences of the promoter region of FSH $\beta$  genes of the Japanese crested ibis, ovine (Guzman *et al.*, 1991), bovine (Kim *et al.*, 1988), porcine (Hirai *et al.*, 1990), rat (Gharib *et al.*, 1989) and human (Jameson *et al.*, 1988). The nucleotide sequence is numbered from the putative transcriptional start site in the ibis FSH $\beta$  gene. A progesterone responsive element (PRE)-like sequence (Webster *et al.*, 1995) and two AP-1 responsive element-like sequences (Huang *et al.*, 2001) reported in ovine FSH $\beta$  gene are boxed with solid line with consensus sequence attached in parenthesis. Parts in which TATAAA sequences are observed in the ibis are shaded.



**Fig. 10.** Nucleotide sequences of the second intron of the FSH $\beta$  gene in the Japanese crested ibis, (L) the longer type and (S) the shorter type. Dashes indicate a gap inserted to maximize identity. Three nucleotide mismatches are marked with a reverse triangle. Part recognized by *EcoRI* in the short type is boxed.

provided the first information on the amino acid sequences of the PGH $\alpha$  and FSH $\beta$  in a bird from the second clade of the Neognathae. High similarities of the PGH $\alpha$  and FSH $\beta$  among ibis, chicken, Japanese quail and ostrich indicate that FSH is generally highly conserved over wide range of avian species.

Lastly, we found two (the shorter and longer) types of the sequence in the second intron of the FSH $\beta$  gene in the Japanese crested ibis. The shorter type was observed in both of two individuals we used, Midori of the Japanese origin and Long-Long of the Chinese origin, whereas the longer type was found only in Long-Long. These results suggested that geographic divergence occur in this species. At present, we are extending the sequence analysis of this intron to other individuals of both the Japanese and Chinese origins. Sequence analysis of the mitochondrial DNA of this species is also proceeding.

### ACKNOWLEDGMENTS

This work is partly supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

### REFERENCES

- Akashi M, Shaw G, Hachiya M, Elstner E, Suzuki G, Koeffler P (1994) Number and location of AUUUA motifs: role in regulating transiently expressed RNAs. *Blood* 83: 3182–3187
- Ando H, Ishii S (1994) Molecular cloning of complementary deoxyribonucleic acids for the pituitary glycoprotein hormone  $\alpha$ -subunit and luteinizing hormone  $\beta$ -subunit precursor molecules of Japanese quail (*Coturnix coturnix japonica*). *Gen Comp Endocrinol* 93: 357–368
- Arai Y, Kubokawa K, Ishii S, Joss JMP (1998) Cloning of cDNA encoding the common alpha subunit precursor molecule of pituitary glycoprotein hormones in the Australian lungfish, *Neoceratodus forsteri*. *Gen Comp Endocrinol* 110: 109–117
- Breathnach R, Chambon P (1981) Organization and expression of eucaryotic split genes coding for proteins. *Ann Rev Biochem* 50: 349–383
- Burnside J, Buckland PR, Chin WW (1988) Isolation and characterization of the gene encoding the  $\alpha$ -subunit of the rat pituitary glycoprotein hormones. *Gene* 70: 67–74
- Chen CY, Shyu AB (1995) AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem Sci* 20: 465–470
- Fiddes JC, Goodman HM (1981) The gene encoding the common alpha subunit of the four human glycoprotein hormones. *J Mol Appl Genet* 1: 3–18
- Foster DN, Foster LK (1991) Cloning and sequence analysis of the common  $\alpha$ -subunit complementary deoxyribonucleic acid of turkey pituitary glycoprotein hormones. *Poult Sci* 70: 2516–2523
- Foster DN, Galehouse D, Giordano T, Min B, Lamb IC, Porter DA, Intehar KJ, Bacon WL (1992) Nucleotide sequence of the cDNA encoding the common  $\alpha$  subunit of the chicken pituitary glycoprotein hormones. *J Mol Endocrinol* 8: 21–27
- Gharib SD, Roy A, Wierman ME, Chin WW (1989) Isolation and characterization of the gene encoding the  $\beta$ -subunit of rat follicle-stimulating hormone. *DNA* 8: 339–349
- Golos TG, Durning M, Fisher JM (1991) Molecular cloning of the rhesus glycoprotein hormone  $\alpha$ -subunit gene. *DNA Cell Biol* 10: 367–380
- Goodwin RG, Moncman CL, Rottman FM, Nilson JH (1983) Characterization and nucleotide sequence of the gene for the common  $\alpha$  subunit of the bovine pituitary glycoprotein hormones. *Nucleic Acids Res* 11: 6873–6882
- Gordon DF, Wood WM, Ridgeway EC (1988) Organization and nucleotide sequence of the mouse  $\alpha$ -subunit gene of the pituitary glycoprotein hormones. *DNA* 7: 679–690
- Guzman K, Miller CD, Phillips CL, Miller WL (1991) The gene encoding ovine follicle-stimulating hormone  $\beta$ : Isolation, characterization, and comparison to a related ovine genomic sequence. *DNA Cell Biol* 10: 593–601
- Hirai T, Takigawa H, Kato Y (1990) The gene for the  $\beta$  subunit of porcine FSH: absence of consensus oestrogen-responsive element and presence of retroposons. *J Mol Endocrinol* 5: 147–158
- Hsieh YL, Chatterjee A, Chien JT, Yu JYL (2001) Molecular cloning of the cDNAs for pituitary glycoprotein hormone  $\alpha$  subunits of two species of duck and their gene regulation. *J Mol Endocrinol* 27: 339–347
- Huang CJ, Huang FL, Wang YC, Chang YS, Lo TB (1992) Organization and nucleotide sequence of carp gonadotropin  $\alpha$  subunit genes. *Biochim Biophys Acta* 1129: 239–242
- Huang HJ, Sebastian J, Strahl BD, Wu JC, Miller WL (2001) Transcriptional regulation of the ovine follicle-stimulating hormone- $\beta$  gene by activin and gonadotropin-releasing hormone (GnRH): Involvement of two proximal activator protein-1 sites for GnRH stimulation. *Endocrinology* 142: 2267–2274
- Ishii S (1999) Application of modern endocrine methods to conservation biology. *Ostrich* 70: 33–38
- Jameson JL, Becker CB, Lindell CM, Habener JF (1988) Human follicle-stimulating hormone  $\beta$ -subunit gene encodes multiple messenger ribonucleic acids. *Mol Endocrinol* 2: 806–815
- Kato Y, Ezashi T, Hirai T, Kato T (1990) The gene for the common  $\alpha$  subunit of porcine pituitary glycoprotein hormone. *J Mol Endocrinol* 7: 27–34
- Kikuchi M, Kobayashi M, Ito T, Kato Y, Ishii S (1998) Cloning of complementary deoxyribonucleic acid for the follicle-stimulating hormone- $\beta$  subunit in the Japanese quail. *Gen Comp Endocrinol* 111: 376–385
- Kim KE, Gordon DF, Maurer RA (1988) Nucleotide sequence of the bovine gene for follicle-stimulating hormone  $\beta$ -subunit. *DNA* 7: 227–233
- Koide Y, Papkoff H, Kawauchi H (1996) Complete amino acid sequences of follitropin and lutropin in the ostrich, *Struthio camelus*. *Eur J Biochem* 240: 262–267
- Kumar TR, Kelly M, Mortrud M, Low MJ, Matzuk MM (1995) Cloning of the mouse gonadotropin  $\beta$ -subunit-encoding genes. I. Structure of the follicle-stimulating hormone  $\beta$ -subunit-encoding gene. *Gene* 166: 333–334
- Noce T, Ando H, Ueda T, Kubokawa K, Higashinakagawa T, Ishii S (1989) Molecular cloning and nucleotide sequence analysis of the putative cDNA for the precursor molecule of the chicken LH-beta subunit. *J Mol Endocrinol* 3: 129–137
- Rosenfeld H, Levavi-Sivan B, Gur G, Melamed P, Meiri I, Yaron Z, Elizur A (2001) Characterization of tilapia FSH $\beta$  gene and analysis of its 5' flanking region. *Comp Biochem Physiol B* 129: 389–398
- Shaw G, Kamen R (1986) A conserved AU sequence from the 3' untranslated region of GM-CSF mRNA mediates selective mRNA degradation. *Cell* 46: 659–667
- Shen ST, Yu JYL (2002) Cloning and gene expression of a cDNA for the chicken follicle-stimulating hormone (FSH)- $\beta$ -subunit. *Gen Comp Endocrinol* 125: 375–386
- Sohn YC, Suetake H, Yoshiura Y, Kobayashi M, Aida K (1998) Structural and expression analysis of gonadotropin I $\beta$  subunit genes in goldfish (*Carassius auratus*). *Gene* 222: 257–267

- Suzuki K, Liu D, Hew CL (1995) A gene encoding chinook salmon (*Oncorhynchus tshawytscha*) gonadotropin  $\alpha$  subunit gene structure and promoter analysis in primary pituitary cells. *Mol Mar Biol Biotechnol* 4: 10–19
- You S, Foster LK, Silsby JL, El Halawani ME, Foster DN (1995) Sequence analysis of the turkey LH $\beta$  subunit and its regulation by gonadotrophin-releasing hormone and prolactin in cultured pituitary cells. *J Mol Endocrinol* 14: 117–129
- van Tuinen M, Sibley CG, Hedges SB (2000) The early history of modern birds inferred from DNA sequences of nuclear and mitochondrial ribosomal genes. *Mol Biol Evol* 17: 451–457
- Wakabayashi S, Kikuchi M, Wada M, Sakai H, Ishii S (1992) Induction of ovarian growth and ovulation by administration of a chicken gonadotrophin preparation to Japanese quail kept under a short-day regimen. *Brit Poultry Sci* 33: 847–858
- Webster JC, Pedersen NR, Edwards DP, Beck CA, Miller WL (1995) The 5'-flanking region of the ovine follicle-stimulating hormone- $\beta$  gene contains six progesterone response elements: Three proximal elements are sufficient to increase transcription in the presence of progesterone. *Endocrinology* 136: 1049–1058
- Wingfield JC, Ishii S, Kikuchi M, Wakabayashi S, Sakai H, Yamaguchi N, Wada M, Chikatsuji K (2000) Biology of a critically endangered species, the Toki (Japanese crested ibis) *Nipponia nippon*. *Ibis* 142: 1–11

(Received December 17, 2002 / Accepted January 14, 2003)