

Role of complement regulatory proteins CD46, CD55 and CD59 in reproduction

Authors: Frolíková, Michaela, Stopková, Romana, Antalíková, Jana, Johnson, Peter M., Stopka, Pavel, et al.

Source: Folia Zoologica, 61(1) : 84-94

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v61.i1.a12.2012>

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Role of complement regulatory proteins CD46, CD55 and CD59 in reproduction

Michaela FROLÍKOVÁ¹, Romana STOPKOVÁ¹, Jana ANTALÍKOVÁ², Peter M. JOHNSON³, Pavel STOPKA¹ and Kateřina DVOŘÁKOVÁ-HORTOVÁ^{1*}

¹ Department of Zoology, Faculty of Science, Charles University in Prague, Viničná 7, 128 44 Prague, Czech Republic; e-mail: hortova@natur.cuni.cz

² Department of Immunogenetics, Institute of Animal Biochemistry and Genetics, Slovak Academy of Sciences, 900 28 Ivanka pri Dunaji, Slovak Republic

³ Division of Immunology, School of Infection and Host Defence, Duncan Building, University of Liverpool, L69 3BX, Liverpool, U.K.

This paper was presented at the 3rd Conference on Genes, Gene Expression and Behaviour held in Hrubá Skála, The Czech Republic, 4.–7.11. 2010

Received 1 July 2011; Accepted 4 October 2011

Abstract. In humans, CD46 has been detected on the acrosomal membrane in sperm, in contrast to widespread surface expression on somatic cells where it plays a key role in the protection from complement attack. In rodents, CD46 is expressed solely on the acrosomal membrane of mature sperm and their immediate precursors, spermatids. A monoclonal antibody against the short consensus repeat (SCR1) ectodomain of CD46 blocks binding of human sperm to zona-free oocytes *in vitro*. However, CD46-knockout mice are fertile and have an accelerated spontaneous acrosome reaction. Wild-caught field mice (*Apodemus*) also exhibit a rapid acrosome reaction and CD46 is not expressed in *Apodemus* sperm. CD46 may, therefore, play a role in stabilization of the acrosomal membrane. Two other complement regulatory proteins, CD55 and CD59, are localized on the plasma membrane of mammalian sperm. In human sperm, CD55 and CD59 are expressed also on the inner acrosomal membrane. It remains to be clarified what is the role of CD46, CD55 and CD59 during fertilization and what are the advantages of not expressing CD46 in field mice sperm.

Key words: CD46, CD55, CD59, sperm, rodent, human

Introduction

CD46 (membrane cofactor protein) is a membrane-associated glycoprotein that is present in human cells, including sperm, and in cells of other mammals. In somatic cells, CD46 is a cofactor facilitating the degradation of C3b convertase and, therefore, plays key role in protecting host cells from complement attack (Seya & Atkinson 1989, Liszewski et al. 1991, Liszewski et al. 1996). Another role of the CD46 protein is activation of several intracellular pathways in different types of cells, including T-lymphocytes (Wong et al. 1997, Wang et al. 2000, Liszewski et al. 2005). On the other hand, expression of CD46

protein was demonstrated solely on sperm in rat and mouse (Mizuno et al. 2004, Johnson et al. 2007), with the exception of field mice (*Apodemus*) where spermatozoal CD46 protein is not expressed (Johnson et al. 2007, Clift et al. 2009a). Several pathogens, such as measles virus (Dorig et al. 1993, Naniche et al. 1993), human herpes virus-6 (Santoro et al. 1999), group B and D adenoviruses (Gaggar et al. 2003, Wu et al. 2004), group A *streptococcus* (Okada et al. 1995) and *Neisseria* (Kallstrom et al. 1997), bind to CD46 and use it as receptor enabling them to enter the cell. The loss of sperm CD46 may be an advantage in the case of promiscuous field mice that are exposed to

a large amount of genital pathogens. The aim of this mini-review is to summarize present knowledge about CD46, as well as two other complement regulatory (CReg) proteins CD55 and CD59, and discuss their role in fertilization.

CD46 and its isoforms

In humans, CD46 protein is expressed by all cells except erythrocytes and exists predominantly in four different isoforms – these isoforms are the products of alternative splicing of a single gene (Liszewski et al. 1994). They consist of four extracellular domains on the amino-terminus called short consensus repeats (SCRs), a serine-threonine-proline (STP)-rich area, transmembrane domain (TM), intracytoplasmic anchor and one of two forms of cytoplasmic tail (CYT-1 of 16 amino acids and CYT-2 of 23 amino acids) that arise by alternative splicing at the carboxyl-terminus (Liszewski & Atkinson 1992). The SCR1, two and four domains are sites of N-linked glycosylation. The STP region is a site of extensive O-linked glycosylation and is composed of 14 or 29 amino acids. The size of the STP region depends on the presence of STP exon B that is often spliced out. The absence of STP exon

B leads to lower molecular weight CD46 isoforms of CD46, while its presence leads to higher molecular weight CD46 isoforms (Liszewski et al. 1991, Post et al. 1991). However, an even lower molecular weight hypoglycosylated isoform of CD46 is expressed in mammalian sperm. In humans, polymorphisms of the *Cd46* gene were detected in normal individuals using *HindIII* and *EcoRI* restriction enzymes (Risk et al. 1991).

Mizuno et al. (2004) demonstrated that CD46 protein is expressed exclusively on the acrosomal membrane of mature sperm and on their immediate precursors, spermatids, in rodents. There are only 11 functional exons in the murine *Cd46* gene compared to the human which contains 14. The murine *Cd46* gene contains exons encoding four SCRs domains, an STP-rich area, intracytoplasmic anchor, transmembrane domain and cytoplasmic tail (Miwa et al. 1998, Tsujimura et al. 1998). Differences in tissue-specific mRNA expression of rat *Cd46* are rather quantitative than qualitative, reaching the highest level in testicular tissue (Mead et al. 1999). Our results in mice confirm the presence of *Cd46* in various tissues (Fig. 1) but the significantly highest expression was




Fig. 1. The level of mRNA abundance of *Cd46* in three male and three female *Mus musculus domesticus* was studied in the following tissues: vomeronasal organ (VNO), nasal-associated lymphoid tissue (NLT), main olfactory epithelium (OE), Harderian gland (HG), extraorbital lacrimal gland (LG), submandibular salivary gland (SG), preputial gland (PG), liver (L), ovary (O), uterus (U), vagina including cervix (V), testes (T), cauda epididymis (CE) and prostate (P). An efficiency-sensitive model with *Gapdh* as a reference gene was used for the calculation of mRNA abundance (Pfaffl 2001). Low expression was detected in all tested tissues; only in testes did the amount of *Cd46* mRNA reach the level of housekeeping gene.

detected in testes. The testicular mRNA abundance of *Cd46* is elevated such that the expression level is almost 4-fold higher than the level typical for other tissues, where expression is far below the level of the common house-keeping gene *Gapdh*. Johnson et al. (2007) demonstrated that field mice of three species, *Apodemus sylvaticus*, *A. flavicollis* and *A. microps*, produce alternatively spliced transcripts of testicular *Cd46* mRNA lacking exons 5-7 or 6-7, together with a 3'-untranslated region of 54nt sequence extension and an often truncated a 5'-terminal sequence deletion (Fig. 2), resulting in failure to express spermatozoal CD46 protein. Similar results were described in the case of *A. agrarius* that produces two alternatively spliced transcripts of testicular *Cd46* mRNA, both lacking exon 7. However, these transcripts differ from transcripts found in other *Apodemus* species. The larger *A. agrarius* *Cd46* transcript has an insert between exons 10 and 11 which, if translated, would result in a novel cytoplasmic tail (Clift et al. 2009b).




Fig. 2. Comparison of CD46 protein structure in mouse and rat, with possible CD46 protein structure in wood mouse in case the protein was translated.

CD46 in sperm

CD46 is exposed to the extracellular environment only after the acrosomal reaction due to its localization to the acrosomal membrane (Anderson et al. 1989, Riley et al. 2002a) (Fig. 3). The SCR1 ectodomain of CD46 especially plays an important role in fertilization. This is indicated also by the absence of the SCR1 ectodomain in all cell types of New World monkeys except sperm (Riley et al. 2002b). A monoclonal antibody against the SCR1 ectodomain of CD46 blocks binding of human sperm to zona-free oocytes *in vitro* (Okabe et al. 1990, Taylor et al. 1994). This finding suggested the possibility that CD46 could be a candidate fusion protein in sperm binding to the complementary egg surface binding protein. Further, Miyado et al. (2000) and Kaji et al. (2000) revealed in CD9 knock-out mice




Fig. 3. Localization of CD46 protein in sperm.

that it is the egg CD9 protein that plays a key role in sperm-egg fusion. In rodents, CD46 was discovered exclusively on the acrosomal membrane of mature sperm and their immediate precursors, spermatids (Mizuno et al. 2004). Surprisingly, antibodies against CD46 did not block fertilization in rat (Mizuno et al. 2007). Inoue et al. (2003) produced CD46-knockout mice and showed that the males had normal testes and, moreover, fertile sperm. However, in contrast with the wild-type sperm, sperm of CD46-deficient males showed an accelerated spontaneous acrosome reaction. The protein IZUMO was later discovered in mouse and human sperm by Inoue et al. (2005) and described as the first sperm membrane protein shown to be essential for sperm-egg fusion.

CD46 in field mice

As well as CD46 knockout mice, wild-caught field mice (*Apodemus*) also exhibit a more rapid acrosome reaction and, surprisingly, it was discovered that field mice do not translate CD46 protein during spermiogenesis; therefore, CD46 protein is not expressed in *Apodemus* sperm (Johnson et al. 2007). Minor nucleotide sequence polymorphism was detected between *A. sylvaticus*, *A. flavicollis* and *A. microps*, and polymorphisms both in noncoding and coding regions were identified between individuals of *A. sylvaticus* mice. These changes in the coding region lead to non-synonymous changes and may be the basis for the unique CD46 splicing (Johnson et al. 2007).

Although CD46 plays a key role in protecting the host cells from complement attack in somatic cells, data published by Inoue et al. (2003) and Johnson et al. (2007) suggested a role for CD46 in sperm relating to integrity of the acrosome vesicle or regulation of the acrosome reaction rather than protection of sperm from complement attack. This led to questions as to what is the role of CD46 during fertilization and what

are the advantages/disadvantages of not expressing CD46 in field mice sperm?

Sperm trains in field mice

Field mice are highly territorial and there is pronounced sperm competition, which is reflected by their disproportionately large testis (Breed & Taylor 2000, Bryja & Stopka 2005). Relative testis mass is associated with the shape of the apical hook in the falciform head of murine rodent sperm (Moore et al. 2002). Sperm of *A. sylvaticus* have a very long apical hook and Moore et al. (2002) demonstrated that these sperm form sperm “trains” with significantly increased motility in comparison with an individual spermatozoon. Dissociation of sperm trains is a prerequisite for fertilization and it is believed that the spontaneous acrosomal reaction of part of the involved sperm plays a role in this process (Moore et al. 2002). A significantly faster spontaneous acrosome reaction rate in field mice compared to house mice (Johnson et al. 2007) may be important for the dissociation of sperm trains and, moreover, may help bypass the primary binding of acrosome-unreacted sperm to zona pellucida (Myles et al. 1987, Yamagata et al. 2002). It is possible that a release of the acrosomal content of many sperm at the same time may affect the zona pellucida (ZP) structure and help sperm to pass faster through the ZP, especially since the acrosome reaction is a time-consuming process lasting for up to an hour (Shur et al. 2006). It has been shown that acrosomal exocytosis specifically induced by ZP3-dependent GalT aggregation is not critical for fertilization (Talbot et al. 2003). Huang et al. (1981) showed that only acrosome-reacted sperm can bind to and penetrate the ZP in the guinea pig. These data suggest that acrosomal instability may provide a novel genus-specific strategy to favour rapid fertilization and competitive advantage in the promiscuous reproductive behaviour of wild field mice (Johnson et al. 2007).

CD46 and cytoskeleton

The way in which CD46 may stabilize the acrosomal membrane is still unclear. One of the possibilities is an interaction of CD46 with β 1-integrins. Integrins are transmembrane receptors involved in cell-cell adhesion and cell migration, proliferation and differentiation as well as cellular apoptosis. These receptors may mediate a signal both into and out of the cell (Liu et al. 2000). CD46 binding to β 1-integrins (Lozahic et al. 2000, Kurita-Taniguchi et al. 2002, Rezcallah et al. 2005), as well as β 1-integrin interaction with the actin cytoskeleton (Calderwood

et al. 2000), has been demonstrated. β 1-integrin interaction with the actin cytoskeleton is realized by direct and indirect binding of integrin β -tails and actin-binding proteins (Critchley 2000, Liu et al. 2000). Actin and actin polymerization play a key role during sperm capacitation and acrosome reaction (Brenner et al. 2003, Dvořáková et al. 2005). Filamentous F-actin, part of cortical cytoskeleton, is localized between the plasma and acrosomal membrane where it protects these two membranes from connection. Moreover, this cytoskeleton plays a pivotal role in relocalization of proteins during the acrosome reaction, e.g. the primary fusogenic protein IZUMO (Sosnik et al. 2009). It appears that CD46 stimulation may activate protein kinase signalling leading to actin reorganization (Wong et al. 1997, Zaffran et al. 2001). These facts may lead to the concept that the absence of CD46 enables faster F-actin breakdown and thus rapid sperm acrosome reaction.

Role of CD46 in sperm protection

Complement components are extensively present in the female reproductive tract, where they protect host cells against pathogens. Hence, it is expected that sperm should have some kind of protection against complement attack in female reproductive tract. This protection is provided by complement regulatory proteins, of which CD46 is one of them.

CD46 is localized on the acrosomal membrane (Anderson et al. 1989, Riley et al. 2002a) in mammalian sperm. For this reason, it may only protect sperm against complement attack after the acrosome reaction when the sperm plasma membrane and other plasma membrane-associated CReg proteins are lost and the inner acrosome membrane is exposed to the surrounding of the female reproductive tract. However, the inner acrosome membrane and CD46 are exposed for a very short time before the actual sperm-egg fusion. On the other hand due to the fact that the huge majority of sperm never get to the egg, and CD46 continues to be expressed, CD46 could have a role in the phagocytic clearance of redundant time-expired sperm. The primary role of sperm CD46 may be expected to be different from protection of sperm against complement attack. Therefore, other plasma membrane-associated Creg proteins are likely to play the main role in protecting sperm against complement-mediated damage prior to fertilization (Clift et al. 2009b).

CD55 and CD59

Besides the presence of CD46 in the acrosome membrane of mammalian sperm, there are two

other CReg proteins (CD55 and CD59) localized in the plasma membrane of mammalian sperm. CD55, termed decay accelerating factor (DAF), inhibits the assembly of C3 convertases (Fujita et al. 1987) similarly to CD46. However CD59 inhibits the assembly of the membrane attack complex on the cellular plasma membrane (Davies et al. 1989).

In humans, in contrast to CD46, CD55 is present only as glycosylphosphatidylinositol (GPI)-anchored molecule. However, the expression of both glycosylphosphatidylinositol (GPI)-anchored and transmembrane (TM)-anchored isoforms of CD55 has been shown in mice (Spicer et al. 1995). CD55 isoforms are the products of two different but highly homologous genes (Spicer et al. 1995), compared to CD46 isoforms which are the products of alternative splicing of a single gene (Liszewski et al. 1994). Lin et al. (2001) showed in mice that GPI-CD55 protein is expressed in most of tissues, while TM-CD55 protein is mainly present in the testis and dendritic cells in spleen. The same distribution of GPI- and TM-CD55 protein was demonstrated in rats, with the difference that GPI- and TM-CD55 are products of alternative splicing of a single copy gene (Hinchliffe et al. 1998, Miwa et al. 2000), such as in case of CD46. CD55 isoforms are also generated by alternative splicing of a single copy gene in *Apodemus sylvaticus* (Clift et al. 2009b).

CD59 is another GPI-anchored protein protecting host cells from complement attack and may also be involved in fertilization. The gene encoding CD59 is duplicated in the mouse and two CD59 isoforms, named CD59a and CD59b, are expressed (Qian et al. 2000). It was shown that CD59a is distributed in most tissues, in contrast to CD59b which is exclusively expressed on mature sperm and their immediate precursors, spermatids (Baalasubramanian et al. 2004, Donev et al. 2008). Only one CD59 isoform was found in the rat (Hughes et al. 1992, Rushmere et al. 1994).

Crry, a rodent-specific membrane CReg protein, takes much of the role of protecting host cells from complement attack because, in contrast to humans, no CD46 isoform is expressed in rodent somatic cells (Hosokawa et al. 1996, Molina 2002).

Overall, the exact role of CD55 and CD59 in sperm is still unclear. In mice and field mice sperm, the presence of these proteins has been shown on the plasma membrane in the acrosome, neck and tail regions (Clift et al. 2009b). The presence of these two proteins in the main domains of rodent sperm plasma membrane indicates that CD55 and CD59 could play the key role in the protection of sperm from

complement attack in female reproductive tract. On the other hand, Donev et al. (2008) published that a monoclonal antibody against CD59b inhibits mouse sperm motility, and it has also been demonstrated that CD55 and CD59 are expressed on the inner acrosomal membrane in human sperm (Cummerson et al. 2006). These facts suggest that CD55 and CD59, as well as CD46, may play a more complex role in process of reproduction.

Complement regulatory proteins and the tetraspanin web

As indicated, the primary function of sperm CD46 is probably different from protection of sperm against complement attack. One of the possibilities is that CD46 mediates specific binding between the spermatozoa and egg. However, CD46 may not be involved directly in initial fusion events, as fusion begins in the equatorial or post-acrosomal regions where the membrane remains intact after the acrosome reaction, rather than the inner acrosomal membrane region where CD46 is expressed (Taylor et al. 1994). In relation to sperm-egg fusion studies, it is relevant that there may be the ability of CD46 to bind to CD9 antigen, which was originally shown in HeLa cell lines (Lozahic et al. 2000) as well as in macrophages (Kurita-Taniguchi et al. 2002). The crucial role of CD9 in sperm-egg fusion was demonstrated by the inability of CD9 knock-out mouse oocytes to fuse with sperm (Le Naour et al. 2000, Miyado et al. 2000, Kaji et al. 2000) and by the ability of anti-CD9 antibodies to inhibit this fusion (Chen et al. 1999, Le Naour et al. 2000, Miller et al. 2000). While the CD9 molecule belongs to the tetraspanin family of proteins forming complexes with each other and with various membrane proteins within a network of molecular interactions called the "tetraspanin web" (Boucheix & Rubinstein 2001, Berditchevski 2001, Hemler 2003), it could be proposed that the association with CD46 mediates binding between spermatozoa and the egg. However, the fact that recombinant proteins including the large extracellular loop of CD9 partially inhibit fusion when preincubated with eggs, but not with sperm, indicates that CD9 is probably not a receptor for a sperm molecule and has drawn attention to molecules that associate with CD9 at the oocyte surface and thus to the tetraspanin web (Zhu et al. 2002). The preferred binding partner of CD9 within the tetraspanin web is CD81. Human oocytes express CD81 and CD151, as well as alpha6 and beta1 integrin subunits (Ziyyat et al. 2006). The expression of CD81 has also been demonstrated on mouse eggs (Takahashi et al. 2001),

but these three tetraspanins behave differently in the mouse with respect to sperm-egg fusion than in human. In any event, CD9 controls the formation cluster that contains the tetraspanin CD151 and the $\alpha 6 \beta 1$ integrin (Ziyyat et al. 2006).

Treatment of mouse oocytes with phosphatidylinositol-specific phospholipase C (an enzyme cleaving GPI-anchored structures) during *in vitro* fertilization studies has shown a dramatically reduced ability of the oocyte to bind and fuse with a sperm (Coonrod et al. 1999). CD55 and CD59 have been identified as egg GPI-anchored proteins (Taylor & Johnson 1996), but mice that have targeted deletions in these genes are fertile (Sun et al. 1999, Holt et al. 2001). These data suggest that sperm-egg interactions depend on the presence of GPI-linked egg surface proteins, and possibly on the presence of other proteins on the plasma membrane. Lefevre et al. (2010) assume the lipid-modulated participation of GPI-anchored proteins and/or the CD9 tetraspanin with other necessary proteins in the formation of a particular constitution making the membrane adequate for fusion; the absence of one of them disrupting the membrane organization.

The majority of tetraspanin CD9 protein interactions in the oolema occur in the *cis* configuration: however, it has been suggested that CD9 may additionally bind in *trans* configuration to a PSG17-related ligand (Ellerman et al. 2003) present on the sperm surface which is an immunoglobulin superfamily (IgSF) member, such as IZUMO. Sutovsky (2009) assume, therefore, that sperm IZUMO may bind to oocyte tetraspanins CD9 and/or CD81 to mediate adhesion and fusion. Within the tetraspanin web on the oolema, CD9 and CD81 may interact with each other and also with integrins that are able to bind to CD46 and interconnect the actin cytoskeleton. Ellerman et al. (2009), in co-immunoprecipitation, studies showed the presence of other sperm proteins associated with IZUMO. They favour the model that, rather than directly mediating interaction between sperm and the egg, IZUMO is essential for formation of complexes with other sperm proteins required for gamete fusion. Miyado et al. (2008) reported that sperm-egg fusion is mediated by vesicles containing CD9 proteins that are released from the egg and interact directly with sperm. CD9 is transferred from the oocyte membrane to the spermatozoon present in the perivitelline space, and it may induce a sperm membrane molecular reorganization (Lefevre et al. 2010). Furthermore, CD9 molecules in the inner acrosomal membrane and also on the plasma membrane covering the equatorial segment of mouse sperm have been detected by Ito et

al. (2010). Therefore, the existence of a tetraspanin web on sperm may be envisaged. It could be that the protein IZUMO, as well as CD46, are part of it. Moreover, CD46 stimulation may activate protein kinase signalling which leads to actin reorganization. CD9 is an organizer of membrane functionality through association with EWI-2, a member of IgSF8, as recently reported (Glazar & Evans 2009). EWI-2 and EWI-F are major partners of CD9 and CD81 and, through their direct interaction with ERM (ezrin/radixin/moesin) proteins, act as linkers to connect tetraspanin-associated microdomains to the actin cytoskeleton (Sala-Valdes et al. 2006). Thus, according to the current state of knowledge, the tetraspanin web may play a role in acrosome stabilization.

Conclusion

The present knowledge about the complement regulatory proteins CD46, CD55 and CD59 suggests that role of these proteins in process of reproduction is not only in protection of sperm against complement attack, but that it is more complex. Moreover, reported data suggest that the role of CD46 protein in rodents may be different compared to humans and other mammals. The absence of CD46 protein in field mice is surprising and leads to the question as to what are the advantages of not expressing CD46 in sperm? There is experimental data suggesting that CD46 may play a role in stabilization of the acrosome vesicle due to its interaction with the cytoskeleton and the fact that CD46 may be a part of the predicted tetraspanin web in sperm. The stability of the acrosome vesicle is believed to be an important feature of sperm quality since only an intact sperm is generally able to bind to the zona pellucida of the egg and consequently fertilize it. On the other hand, in highly promiscuous species such as field mice, the simultaneous release of the acrosomal content of a majority of the sperm population, occurring independently of the zona pellucida but in its immediate surrounding, could weaken the zona pellucida structure. In this case, not having the acrosome would represent for sperm an advantage bypassing the primary zona pellucida binding, therefore accelerating the actual process of fertilization. Another advantage of not expressing sperm CD46 protein in this highly promiscuous environment could be result of selective pressure towards protection of sperm against a certain pathogens that use CD46 as a cell surface receptor. However, field mice sperm lacking CD46 are still protected against complement-mediated attack by expressing

on their surface the CD55 and CD59 complement regulatory proteins (Clift et al. 2009b). It has been also shown that CD59 may not only be involved in sperm protection but may also play a role in sperm motility (Donev et al. 2008). It appears that all three complement regulatory proteins, CD46, CD55 and CD59, are not involved in only one particular process but their contribution to sperm fertilization ability is more general and needs yet to be further clarified. Detailed investigation of this broad field is needed to

understand fully and comprehend the complexity of the subject.

Acknowledgements

This work was supported by grants from the Ministry of Education of the Czech Republic nos. VC IM06011, VZ 0021620828, the Institutional Research Support no. SVV263207, and by Slovak Research and Development Agency of the Slovak Republic no. APVV-0137-10.

Literature

- Anderson D.J., Michaelson J.S. & Johnson P.M. 1989: Trophoblast/leukocyte-common antigen is expressed by human testicular germ cells and appears on the surface of acrosome-reacted sperm. *Biol. Reprod.* 41: 285–293.
- Baalasubramanian S., Harris C.L., Donev R.M., Mizuno M., Omidvar N., Song W.C. & Morgan B.P. 2004: CD59a is the primary regulator of membrane attack complex assembly in the mouse. *J. Immunol.* 173: 3684–3692.
- Berditchevski F. 2001: Complexes of tetraspanins with integrins: more than meets the eye. *J. Cell. Sci.* 114: 4143–4151.
- Boucheix C. & Rubinstein E. 2001: Tetraspanins. *Cell. Mol. Life Sci.* 58: 1189–1205.
- Breed W.G. & Taylor J. 2000: Body mass, testes, and sperm size in murine rodents. *J. Mammal.* 81: 758–768.
- Brener E., Rubinstein S., Cohen G., Shternall K., Rivlin J. & Breitbart H. 2003: Remodeling of the actin cytoskeleton during mammalian sperm capacitation and acrosome reaction. *Biol. Reprod.* 68: 837–845.
- Bryja J. & Stopka P. 2005: Facultative promiscuity in a presumably monogamous mouse *Apodemus microps*. *Acta Theriol.* 50: 189–196.
- Calderwood D.A., Shattil S.J. & Ginsberg M.H. 2000: Integrins and actin filaments: reciprocal regulation of cell adhesion and signaling. *J. Biol. Chem.* 275: 22607–22610.
- Chen M.S., Tung K.S., Coonrod S.A., Takahashi Y., Bigler D., Chang A., Yamashita Y., Kincade P.W., Herr J. & White J.M. 1999: Role of the integrin-associated protein CD9 in binding between sperm ADAM 2 and the egg integrin alpha6beta1: implication for murine fertilization. *Proc. Natl. Acad. Sci. USA* 96: 11830–11835.
- Clift L.E., Andrlíková P., Frolíková M., Stopka P., Bryja J., Flanagan B.F., Johnson P.M. & Dvořáková-Hortová K. 2009a: Absence of spermatozoal CD46 protein expression and associated rapid acrosome reaction rate in striped field mice (*Apodemus agrarius*). *Reprod. Biol. Endocrinol.* 7: 29–38.
- Clift L.E., Dvořáková-Hortová K., Frolíková M., Andrlíková P., Salman S., Stopka P., Flanagan B.F. & Johnson P.M. 2009b: CD55 and CD59 protein expression by *Apodemus* (field mice) sperm in the absence of CD46. *J. Reprod. Immunol.* 81: 62–73.
- Conrod S.A., Naaby-Hansen S., Shetty J., Shibahara H., Chen M., White J.M. & Herr J.C. 1999: Treatment of mouse oocyte with PI-PLC releases 70-kDa (pI5) and 35- to 45-kDa (pI5.5) protein clusters from the egg surface and inhibits sperm-ooolema binding and fusion. *Dev. Biol.* 207: 334–349.
- Critchley D.R. 2000: Focal adhesions – the cytoskeletal connection. *Curr. Opin. Cell Biol.* 12: 133–139.
- Cummerson J.A., Flanagan B.F., Spiller D.G. & Johnson P.M. 2006: The complement regulatory proteins CD55 (decay accelerating factor) and CD59 are expressed on the inner acrosomal membrane of human spermatozoa as well as CD46 (membrane cofactor protein). *Immunol.* 118: 333–342.
- Davies A., Simmons D.L., Hale G., Harrison R.A., Tighe H., Lachmann P.J. & Waldmann H. 1989: CD59, an LY-6-like protein expressed in human lymphoid cells, regulates the action of the complement membrane attack complex on homologous cells. *J. Exp. Med.* 170 (3): 637–654.
- Donev R.M., Sivasankar B., Mizuno M. & Morgan B.P. 2008: The mouse complement regulator CD59b is significantly expressed only in testis and plays roles in sperm acrosome activation and motility. *Mol. Immunol.* 45: 534–542.
- Dorig R.E., Marciel A., Chopra A. & Richardson C.D. 1993: The human CD46 molecule is a receptor for measles virus (Edmonston strain). *Cell* 75: 295–305.
- Dvořáková K., Moore H.D.M., Šebková N. & Paleček J. 2005: Cytoskeleton localization in the sperm head prior to fertilization. *Reproduction* 130: 61–69.

- Ellerman D.A., Ha C., Primakoff P., Myles D.G. & Dveksler G.S. 2003: Direct binding of the ligand PSG17 to CD9 requires a CD9 site essential for sperm-egg fusion. *Mol. Biol. Cell.* 14: 5098–5103.
- Ellerman D.A., Pei J., Gupta S., Snell W.J., Myles D. & Primakoff P. 2009: Izumo is part of a multiprotein family whose members form large complexes on mammalian sperm. *Mol. Reprod. Develop.* 76: 1188–1199.
- Fujita T., Toshima T., Ogawa K., Iida K. & Tamura N. 1987: The mechanism of action of decay-accelerating factor (DAF). DAF inhibits the assembly of C3 convertases by dissociating C2a and Bb. *J. Exp. Med.* 166: 1221–1228.
- Gaggar A., Shayakhmetov D.M. & Lieber A. 2003: CD46 is a cellular receptor for group B adenoviruses. *Nat. Med.* 9: 1408–1412.
- Glazar A.I. & Evans J.P. 2009: IgSF8 (EWI-2) and CD9 in fertilization: evidence of distinct functions for CD9 and a CD9-associated protein in mammalian sperm/egg interaction. *Reprod. Fertil. Dev.* 21: 293–303.
- Hemler M.E. 2003: Tetraspanin proteins mediate cellular penetration, invasion, and fusion events and define a novel type of membrane microdomain. *Annu. Rev. Cell Dev. Biol.* 19: 397–422.
- Hinchliffe S.J., Spiller O.B., Rushmere N.K. & Morgan B.P. 1998: Molecular cloning and functional characterization of the rat analogue of human decay-accelerating factor (CD55). *J. Immunol.* 161: 5695–5703.
- Holt D.S., Botto M., Bygrave A.E., Hanna S.M., Walport M.J. & Morgan B.P. 2001: Targeted deletion of the CD59 gene causes spontaneous hemolysis and hemoglobinuria. *Blood* 98: 442–449.
- Hosokawa M., Nonaka M., Okada N., Nonaka M. & Okada H. 1996: Molecular cloning of guinea pig membrane cofactor protein: preferential expression in testis. *J. Immunol.* 157 (11): 4946–4952.
- Huang T.T., Fleming A.D. & Yanagimachi R. 1981: Only acrosome-reacted spermatozoa can bind to and penetrate zona pellucida: a study using the guinea pig. *J. Expl. Zool.* 217: 287–290.
- Hughes T.R., Piddlesden S.J., Williams J.D., Harrison R.A. & Morgan B.P. 1992: Isolation and characterization of a membrane protein from rat erythrocytes which inhibits lysis by the membrane attack complex of rat complement. *Biochem. J.* 284: 169–176.
- Inoue N., Ikawa M., Nakanishi T., Matsumoto M., Nomura M., Seya T. & Okabe M. 2003: Disruption of mouse CD46 causes an accelerated spontaneous acrosome reaction in sperm. *Mol. Cell. Biol.* 23: 2614–2622.
- Inoue N., Ikawa M., Isotani A. & Okabe M. 2005: The immunoglobulin superfamily protein Izumo is required for sperm to fuse with eggs. *Nature* 434: 234–238.
- Ito C., Yamatoya K., Yoshida K., Maekawa M., Miyado K. & Toshimori K. 2010: Tetraspanin family protein CD9 in the mouse sperm: unique localization, appearance, behavior and fate during fertilization. *Cell. Tissue Res.* 340: 583–594.
- Johnson P.M., Clift L.E., Andrlíková P., Jursová M., Flanagan B.F., Cummerson J.A., Stopka P. & Dvořáková-Hortová K. 2007: Rapid sperm acrosome reaction in the absence of acrosomal CD46 expression in promiscuous field mice (*Apodemus*). *Reproduction* 134: 739–747.
- Kaji K., Oda S., Shikano T., Ohnuki T., Uematsu Y., Sakagami J., Tada N., Miyazaki S. & Kudo A. 2000: The gamete fusion process is defective in eggs of Cd9-deficient mice. *Nat. Genet.* 24: 279–282.
- Kallstrom H., Liszewski M.K., Atkinson J.P. & Jonsson A.B. 1997: Membrane cofactor protein (MCP or CD46) is a cellular pilus receptor for pathogenic *Neisseria*. *Mol. Microbiol.* 25: 639–647.
- Kurita-Taniguchi M.K., Murabayashi N., Fukui A., Tsuji S., Matsumoto M., Toyoshima K. & Seya T. 2002: Molecular assembly of CD46 with CD9, alpha3-beta1 integrin and protein tyrosine phosphatase SHP-1 in human macrophages through differentiation by GM-CSF. *Mol. Immunol.* 38: 689–700.
- Lefevre B., Wolf J.P. & Ziyat A. 2010: Sperm egg interaction: is there a link between tetraspanin(s) and GPI-anchored protein(s)? *BioEssays* 32: 143–152.
- Le Naour F., Rubinstein E., Jasmin C., Prenant M. & Boucheix C. 2000: Severely reduced female fertility in CD9-deficient mice. *Science* 287: 319–321.
- Lin F., Fukuoka Y., Spicer A., Ohta R., Okada N., Harris C.L., Emancipator S.N. & Medof M.E. 2001: Tissue distribution of products of the mouse decay-accelerating factor (DAF) genes. Exploitation of a Dafi knock-out mouse and site-specific monoclonal antibodies. *Immunol.* 104: 215–225.
- Liszewski M.K., Post T.W. & Atkinson J.P. 1991: Membrane cofactor protein (MCP or CD46): newest member of the regulators of complement activation gene cluster. *Annu. Rev. Immunol.* 9: 431–455.
- Liszewski M.K. & Atkinson J.P. 1992: Membrane cofactor protein. *Curr. Top. Microbiol. Immunol.* 178: 45–60.

- Liszewski M.K., Tejda I. & Atkinson J.P. 1994: Membrane cofactor protein (CD46) of complement. Processing differences related to alternatively spliced cytoplasmic domains. *J. Biol. Chem.* 269: 10776–10779.
- Liszewski M.K., Farries T.C., Lublin D.M., Rooney I.A. & Atkinson J.P. 1996: Control of the complement system. *Adv. Immunol.* 61: 201–283.
- Liszewski M.K., Kemper C., Price J.D. & Atkinson J.P. 2005: Emerging roles and new functions of CD46. *Semin. Immunopathol.* 27: 345–358.
- Liu S., Calderwood D.A. & Ginsberg M.H. 2000: Integrin cytoplasmic domain-binding proteins. *J. Cell Sci.* 113: 3563–3571.
- Lozahic S., Christiansen D., Manie S., Gerlier D., Billard M., Boucheix C. & Rubinstein E. 2000: CD46 (membrane cofactor protein) associates with multiple $\beta 1$ integrins and tetraspans. *Eur. J. Immunol.* 30: 900–907.
- Mead R., Hinchliffe S.J. & Morgan B.P. 1999: Molecular cloning, expression and characterization of the rat analogue of human membrane cofactor protein (MCP/CD46). *Immunol.* 98: 137–143.
- Miller B.J., Georges-Labouesse E., Primakoff P. & Myles D.G. 2000: Normal fertilization occurs with eggs lacking the integrin $\alpha 6\beta 1$ and is CD9-dependent. *J. Cell. Biol.* 149: 1289–1296.
- Miwa T., Nonaka M., Okada N., Wakana S., Shiroishi T. & Okada H. 1998: Molecular cloning of rat and mouse membrane cofactor protein (MCP, CD46): preferential expression in testis and close linkage between the mouse *Mcp* and *Cr2* genes on distal chromosome 1. *Immunogenet.* 48: 363–371.
- Miwa T., Okada N. & Okada H. 2000: Alternative exon usage in the 3 FORMTEXT region of a single gene generates glycosylphosphatidylinositolanchored and transmembrane forms of rat decay-accelerating factor. *Immunogenet.* 51: 129–137.
- Miyado K., Yamada G., Hasuwa H., Nakamura Y., Ryu F., Suzuki K., Kosai K., Inoue K., Onura A., Okabe M. & Mekada E. 2000: Requirement of CD9 on the egg plasma membrane for fertilization. *Science* 287: 321–324.
- Miyado K., Yoshida K., Yamagata K., Sakakibara K., Okabe M., Wang X., Miyamoto K., Akutsu H., Kondo T., Takahshi Y., Ban T., Ito C., Toshimori K., Nakamura A., Ito M., Miyado M., Mekada E. & Umezawa A. 2008: The fusing ability of sperm is bestowed by CD9-containing vesicles released from eggs in mice. *Proc. Natl. Acad. Sci. USA* 105: 12921–12926.
- Mizuno M., Harris C.L., Johnson P.M. & Morgan B.P. 2004: Rat membrane cofactor protein (MCP; CD46) is expressed only in the acrosome of developing and mature spermatozoa and mediates binding to immobilized activated C3. *Biol. Reprod.* 71: 1374–1383.
- Mizuno M., Harris C.L. & Morgan B.P. 2007: Immunization with autologous CD46 generates a strong autoantibody response in rats that targets spermatozoa. *J. Reprod. Immunol.* 73: 135–147.
- Molina H. 2002: The murine complement regulator Crry: new insights into the immunobiology of complement regulation. *Cell. Mol. Life Sci.* 59: 220–229.
- Moore H., Dvořáková K., Jenkins N. & Breed W. 2002: Exceptional sperm cooperation in the wood mouse. *Nature* 418: 174–177.
- Myles D.G., Hyatt H. & Primakoff P. 1987: Binding of both acrosome-intact and acrosome-reacted guinea pig sperm to the zona pellucida during in vitro fertilization. *Dev. Biol.* 121: 559–567.
- Naniche D., Varior-Krishnan G., Cervoni F., Wild T.F., Rossi B., Rabourdin-Combe C. & Gerlier D. 1993: Human membrane cofactor protein (CD46) acts as a cellular receptor for measles virus. *J. Virol.* 67: 6025–6032.
- Okabe M., Nagira M., Kawai Y., Matzno S., Mimura T. & Mayumi T. 1990: A human sperm antigen possibly involved in binding and/or fusion with zona-free hamster eggs. *Fertil. Steril.* 54: 1121–1126.
- Okada N., Liszewski M.K., Atkinson J.P. & Caparon M. 1995: Membrane cofactor protein (CD46) is a keratinocyte receptor for the M protein of the group A streptococcus. *Proc. Natl. Acad. Sci. USA* 92: 2489–2493.
- Pfaffl M.W. 2001: A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acid Res.* 29: 2002–2007.
- Post T.W., Liszewski M.K., Adams E.M., Tedja I., Miller E.A. & Atkinson J.P. 1991: Membrane cofactor protein of the complement system: alternative splicing of serine/threonine/proline-rich exons and cytoplasmic tails produces multiple isoforms that correlate with protein phenotype. *J. Exp. Med.* 174: 93–102.

- Qian Y.M., Qin X., Miwa T., Sun X., Halperin J.A. & Song W.C. 2000: Identification and functional characterization of a new gene encoding the mouse terminal complement inhibitor CD59. *J. Immunol.* 165 (5): 2528–2534.
- Rezcallah M.S., Hodges K., Gill D.B., Atkinson J.P., Wang B. & Cleary P.P. 2005: Engagement of CD46 and $\alpha 5\beta 1$ integrin by group A streptococci is required for efficient invasion of epithelial cells. *Cell. Microbiol.* 7: 645–653.
- Riley R.C., Kemper C., Leung M. & Atkinson J.P. 2002a: Characterization of human membrane cofactor protein (MCP; CD46) on spermatozoa. *Mol. Reprod. Dev.* 62: 534–546.
- Riley R.C., Tannenbaum P.L., Abbott D.H. & Atkinson J.P. 2002b: Inhibiting measles virus infection but promoting reproduction: an explanation for splicing and tissue-specific expression of CD46. *J. Immunol.* 169: 5405–5409.
- Risk J.M., Flanagan B.F. & Johnson P.M. 1991: Polymorphism of the human CD46 gene in normal individuals and in recurrent spontaneous abortion. *Hum. Immunol.* 30: 162–167.
- Rushmere N.K., Harrison R.A., Van Den Berg C.W. & Morgan B.P. 1994: Molecular cloning of the rat analogue of human CD59: structural comparison with human CD59 and identification of a putative active site. *Biochem. J.* 304: 595–601.
- Sala-Valdes M., Ursa A., Charrin S., Rubinstein E., Hemler M.E., Sanchez-Madrid F. & Yanez-Mo M. 2006: EWI-2 and EWI-F link the tetraspanin web to the actin cytoskeleton through their direct association with ezrin-radixin-moesin proteins. *J. Biol. Chem.* 281: 19665–19657.
- Santoro F., Kennedy P.E., Locatelli G., Malnati M.S., Berger E.A. & Lusso P. 1999: CD46 is a cellular receptor for human herpesvirus 6. *Cell* 99: 817–827.
- Seya T. & Atkinson J.P. 1989: Functional properties of membrane cofactor protein of complement. *Biochem. J.* 264: 581–588.
- Shur B.D., Rodeheffer C., Ensslin M.A., Lyng R. & Raymond A. 2006: Identification of novel gamete receptors that mediate sperm adhesion to the egg coat. *Mol. Cell. Endocrinol.* 250: 137–148.
- Sosnik J., Miranda P.V., Spiridonov N.A., Yoon S.Y., Fissore R.A., Johnson G.R. & Visconti P.E. 2009: Tssk6 is required for Izumo relocalization and gamete fusion in the mouse. *J. Cell. Sci.* 122: 2741–2749.
- Spicer A.P., Seldin M.F. & Gendler S.J. 1995: Molecular cloning and chromosomal localization of the mouse decay-accelerating factor genes. Duplicated genes encode glycosylphosphatidylinositol anchored and transmembrane forms. *J. Immunol.* 155: 3079–3091.
- Sun X., Funk C.D., Deng C., Sahu A., Lambris D. & Song W.C. 1999: Role of decay accelerating factor in regulating complement activation on the erythrocyte surface as revealed by gene targeting. *Proc. Natl. Acad. Sci. USA* 96: 628–633.
- Sutovsky P. 2009: Sperm-egg adhesion and fusion in mammals. *Expert Rev. Mol. Med.* 11: 1–12.
- Takahashi Y., Bigler D., Ito Y. & White J.M. 2001: Sequence specific interaction between the disintegrin domain of mouse ADAM 3 and murine eggs: role of beta1 integrin-associated proteins CD9, CD81, and CD98. *Mol. Biol. Cell* 12: 809–820.
- Talbot P., Shur B.D. & Myles D.G. 2003: Cell adhesion and fertilization: steps in oocyte transport, sperm-zona pellucida interaction, and sperm-egg fusion. *Biol. Reprod.* 68: 1–9.
- Taylor C.T., Biljan M.M., Kingsland C.R. & Johnson P.M. 1994: Inhibition of human spermatozoon-oocyte interaction *in vitro* by monoclonal antibodies to CD46 (membrane cofactor protein). *Hum. Reprod.* 9: 907–911.
- Taylor C.T. & Johnson P.M. 1996: Complement binding proteins are strongly expressed by human preimplantation blastocysts and cumulus cells as well as gametes. *Mol. Hum. Reprod.* 2: 52–59.
- Tsujimura A., Shida K., Kitamura M., Nomura M., Takeda J., Tanaka H., Matsumoto M., Matsumiya K., Okuyama A., Nishimune Y., Okabe M. & Seya T. 1998: Molecular cloning of a murine homologue of membrane cofactor protein (CD46): preferential expression in testicular germ cells. *Biochem. J.* 330: 163–168.
- Wang G., Liszewski M.K., Chan A.C. & Atkinson J.P. 2000: Membrane cofactor protein (MCP; CD46): isoform-specific tyrosine phosphorylation. *J. Immunol.* 164: 1839–1846.
- Wong T.C., Yant S., Harder B.J., Korte-Sarfaty J. & Hirano A. 1997: The cytoplasmic domains of complement regulatory protein CD46 interact with multiple kinases in macrophages. *J. Leuk. Biol.* 62: 892–900.

- Wu E., Trauger S.A., Pache L., Mullen T.M., von Seggern D.J., Siuzdak G. & Nemerow G.R. 2004: Membrane cofactor protein is a receptor for adenoviruses associated with epidemic keratoconjunctivitis. *J. Virol.* 78: 3897–3905.
- Yamagata K., Nakanishi T., Ikawa M., Yamaguchi R., Moss S.B. & Okabe M. 2002: Sperm from the calmegin-deficient mouse have normal abilities for binding and fusion to the egg plasmamembrane. *Dev. Biol.* 250: 348–357.
- Zaffran Y., Destaing O., Roux A., Ory S., Nheu T., Jurdic P., Roubourdin-Combe C. & Astier A.L. 2001: CD46/CD3 costimulation induces morphological changes of human T cells and activation of Vav, Rac, and extracellular signal-regulated kinase mitogen-activated protein kinase. *J. Immunol.* 167: 6780–6785.
- Zhu G.Z., Miller B.J., Boucheix C., Rubinstein E., Liu C.C., Hynes R.O., Myles D.G. & Primakoff P. 2002: Residues of SFQ (173-175) in the large extracellular loop of CD9 are required for gamete fusion. *Development* 129: 1995–2002.
- Ziyyat A., Rubinstein E., Monier-Gavelle F., Barraud V., Kulski O., Prenent M., Boucheix C., Bomsel M. & Wolf J.P. 2006: CD9 controls the formation of clusters that contain tetraspanins and the integrin $\alpha 6 \beta 1$, which are involved in human and mouse gamete fusion. *J. Cell. Sci.* 119: 416–424.

Appendix. Methods: real-time PCR analysis of *Cd46* expression.

Although this is a review-type of article, we present here original data on *Cd46* expression to demonstrate the prevalence of *Cd46* mRNA in testicular tissue. Total RNA was extracted from 11 different tissues of three male and three female *M. m. domesticus*. Tissues included in the expression analysis were as follows: vomeronasal organ (VNO), nasal-associated lymphoid tissue (NLT), main olfactory epithelium (OE), Harderian gland (HG), extraorbital lacrimal gland (LG), submandibular salivary gland (SG), preputial gland (PG), liver (L), ovary (O), uterus (U), vagina including cervix (V), testes (T), cauda epididymis (CE) and prostate (P). RNA isolation was performed using TriReagent from Sigma Life Science according to the manufacturer's protocol. The purity of RNA was assessed from the ratio of the optical densities at 260 and 280 nm, and the integrity was controlled by electrophoresis on a 1 % agarose gel containing ethidium bromide. Prior to cDNA synthesis, the RNA was cleaned with a DNase protocol (Fermentas) to avoid potential DNA contamination. One microgram of total RNA was used for the synthesis of single-stranded cDNA according to a first-strand cDNA synthesis protocol (Fermentas UAB, Vilnius, Lithuania) with RevertAid™M-MuLV Reverse Transcriptase and oligo(dT)₁₈ primer. Real-time PCR was performed on a Light Cycler 480 (Roche Applied Sciences) using the specific dual hydrolyzation probe method (Universal Probe – Roche Applied Sciences) with the appropriate Probe Master kit (Roche) and protocol according to the manufacturer's protocol. Primers and specific probe number 17 for the *Cd46* sequence were designed by Universal probe library software (<https://www.roche-applied-science.com/sis/rtPCR/upl/index.jsp>) using the NCBI reference sequence NM_010778.3, intron-spanning assay condition and multiplex PCR with reference gene (*Gapdh*) requirement. The resulting primers were: *Cd46* F: 5'-AGCCCTCCGGAGTGTAAGT-3' on the 4th exon and *Cd46* R: 5'-ACATCACTGTTGATTGATAGGAAAAT-3' on the 5th exon. Correct amplification of the primer product was validated by sequencing. PCR amplification was performed with the following conditions: initial denaturation at 95 °C for 10 min, followed by 40 cycles consisting of denaturation at 95 °C for 10 s, annealing at 60 °C for 10 s where fluorescence was acquired, and elongation at 72 °C for 5 s. The data used for calculation are the means of Cp values of triplicate samples. Variation of triplicates did not exceed 0.5 CP. The level of *Cd46* mRNA in each sample was calculated relative to the mRNA level of the reference gene (*Gapdh*) amplified in the same well. A calibration curve was generated for both the target and reference gene, and values for PCR efficiency (E; in both cases, E was equal to 0.99, i.e. 99 % efficiency of PCR reaction) were used in a formula calculating normalized mRNA abundance (Efficiency sensitive model): Normalized amount of mRNA = $(1 + E_{\text{reference}})^{CP_{\text{reference gene}}} / (1 + E_{\text{target}})^{CP_{\text{target gene}}}$

Non-template and non-RT reactions were performed as a control samples.