



Age Estimation of Live Arctic Foxes *Vulpes lagopus* Based on Teeth Condition

Authors: Chevallier, Clément, Gauthier, Gilles, and Berteaux, Dominique

Source: Wildlife Biology, 2017(1)

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/wlb.00304>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Age estimation of live arctic foxes *Vulpes lagopus* based on teeth condition

Clément Chevallier, Gilles Gauthier and Dominique Berteaux

C. Chevallier (chevallier.clement@uqar.ca) and D. Berteaux, Canada Research Chair on Northern Biodiversity and Centre d'Études Nordiques, Univ. du Québec à Rimouski, 300 Allée des Ursulines, Rimouski, QC, G5L3A1, Canada. – G. Gauthier, Dépt de biologie and Centre d'Études Nordiques, Univ. Laval, Quebec city, QC, G1V 0A6, Canada

Aging individuals is a prerequisite to many studies in wildlife biology. In mammals, the most accurate methods are based on cement lines analysis and require tooth extraction. Here, we adapted a method based on tooth wear assessment in live individuals and calibrated against cement lines counts on dead individuals. We developed and tested this less invasive method in arctic foxes *Vulpes lagopus*. We photographed dentition of live individuals captured in 2006–2013 and of skulls collected opportunistically in 2003–2013. Five observers assessed twice the tooth wear of 245 sets of pictures (left, frontal, and right views of the dentition for 179 captures and 66 skulls, with a mean of 7.2 pictures per set) to obtain a teeth condition index (TCI) for each set. TCI was highly repeatable, with an intraclass correlation coefficient averaging 0.89 and 0.83 within and between observers, respectively. We then used 44 known-aged individuals to predict fox age from TCI through linear regression ($\text{age} = 0.62 \times \text{TCI} + 1.04$, $r^2 = 0.64$). We tested through jackknife validation the ability of this model to accurately age foxes. The rate of correct classification of exact age was 39%, while 80% and 100% of predicted ages were within one and two years of the true age, respectively. The model predicted correctly the age class (young, prime-age adult, old) of individuals 75% of the time. We conclude that our proposed method, which is much less invasive than those based on tooth extraction, can allow progress in the study of demography of arctic foxes, and potentially other mammal species.

Age is a key parameter in wildlife biology as it has a strong impact on the behaviour and ecology of individuals, and thus on the demography and dynamics of populations. For example, reproductive and survival rates are strongly age-dependent in mammals. The growth rate of a population is thus intimately linked to the proportion of individuals in particular age classes, called the population age structure. Yet, researchers often lack such information, even when individuals can be captured (Delahay et al. 2011).

More than 10 methods have been reported for aging wild mammals (Morris 1972, Spinage 1973, Harris 1978). Among these, the most effective approach consists in counting incremental lines in tooth cement (Harris 1978). The cement, a calcified tissue surrounding the dentine and deposited seasonally (Santymire et al. 2012), is made of light and dark layers formed respectively during periods of rapid and restricted growth. Each layer pair represents one

year in most cases (Pekelharing 1970, Coy and Garshelis 1992, Lohr et al. 2011). Cement lines count is a common technique for aging mammals in general (Grue and Jensen 1979, Gipson et al. 2000) and carnivores in particular polar bears *Ursus maritimus* (Christensen-Dalsgaard et al. 2009), coyotes *Canis latrans* (Linhart and Knowlton 1967), red fox *Vulpes vulpes* (Harris 1978, Cavallini and Santini 1995), arctic fox *Vulpes lagopus* (Grue and Jensen 1976), gray wolf *Canis lupus* (Landon et al. 1998). Age estimates obtained by counting the number of growth layer groups deposited in teeth cement proved to be accurate for many species (King 1991, Hamlin et al. 2000). However, this method requires a tooth extraction and subsequent laboratory analyses, which generate potential ethical problems as well as delays in age determination of individuals. The need for a faster and less invasive method for age estimation of wild animals is clear.

Accordingly, many mammal studies have used tooth wear for estimating age of individuals (red fox: Harris 1978, badger *Meles meles*: Harris et al. 1992, roe deer *Capreolus capreolus*: Hewison 1999, gray wolf: Gipson et al. 2000, racoon *Procyon lotor*: Grau et al. 1970, black bear *Ursus americanus*: Shimoinaba and Oi 2015). Most of the times, tooth wear was used in dead (Harris 1978, Smuts et al. 1978,

This work is licensed under the terms of a Creative Commons Attribution 4.0 International License (CC-BY) <<http://creativecommons.org/licenses/by/4.0/>>. The license permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Harris et al. 1992, Gay and Best 1996, Hewison 1999, Gipson et al. 2000, Rosatte and Silver 2007), anesthetized (Stander 1997, Van Horn et al. 2003, Olifiers et al. 2010, Delahay et al. 2011) or captive (Grau et al. 1970) animals. To our knowledge, only Lohr et al. (2011) used tooth wear to age non-anesthetized wild mammals, in their case, the brushtail possum *Trichosurus vulpecula*. Moreover, many studies cited above required a detailed expertise in dental morphology, which might necessitate some specific training at costs and constraints not affordable to many projects. An aging method available to non-experts and applicable to live and non-anesthetized animals captured in the wild would represent substantial progress in wildlife biology.

We used the arctic fox as study model to validate a minimally invasive method to age wild carnivores from their teeth condition. This work was conducted in the context of a detailed study on the behaviour and ecology of the species in the Canadian High Arctic. Our studied population is monitored since 1993 through den surveys and since 2003 through capture–mark–recapture, but age of individuals captured as adults is unknown. Given that cement lines count can accurately determine age of arctic foxes due to large differences between winter and summer diet (Grue and Jensen 1976), we combined cement lines count on skulls of dead foxes found opportunistically in the field and analysis of tooth wear on live foxes to test a teeth condition index for arctic foxes. This method was modified from Olifiers et al. (2010) and adapted for our model species. We had three specific objectives: 1) to propose a visual teeth condition index (TCI) for an arctic predator, 2) to determine its repeatability within and across observers and 3) to build and test a predictive model with data from known-age individuals.

Material and methods

Study area

We worked in the south plain of Bylot Island (73°N, 80°W) which is part of Sirmilik National Park, Nunavut, Canada. Our study area increased from 425 km² in 2003 to 520 km² since 2008. The arctic fox is one of the main terrestrial predators of the local food web (Gauthier et al. 2004, Legagneux et al. 2012). Its diet is composed of lemmings, *Lemmus trimucronatus* and *Dicrostonyx groenlandicus*, and greater snow geese *Chen caerulescens atlanticus*, mostly eggs and goslings (B  ty and Gauthier 2002, Careau et al. 2008, Tarroux et al. 2012, Giroux et al. 2012), as well as seal pups and seal carcasses found on the sea ice during winter and spring (Gagnon and Berteaux 2009).

Field protocol

We captured arctic foxes between mid-May and mid-August, from 2006 to 2013. We trapped cubs and adults using collapsible live traps (Tomahawk cage trap no. 205) or padded leghold traps (Softcatch no. 1) visited every six hours (Proulx et al. 1994). We sexed and ear-tagged each captured fox and photographed the dentition of ≥ 1 year-old individuals. One observer maintained the fox mouth closed with the lips pulled apart to expose dentition, while the other took

the pictures. A minimum of three pictures (right lateral, left lateral and frontal views) were taken for each capture event. All pictures taken during a capture event constitute hereafter a ‘set of pictures’, with a mean of 5.5 pictures per set since some pictures were repeated when quality was uncertain.

We anesthetized 60 of 179 captured adults judged too agitated for safe manipulation, using an intramuscular injection of medetomidine (0.05 ml kg⁻¹) and ketamine (0.03 ml kg⁻¹). We used atipamezol (0.05 ml kg⁻¹) as antidote before releasing foxes. The remaining 119 adults were manipulated and photographed carefully with kevlar gloves only.

We collected adult-size skulls in the study area between 2003 and 2013. Some were found on the ground, often with little skin remains, after they had been exposed to scavengers and weathering, whereas others were collected from fresh carcasses of dead foxes. After cleaning skulls, we photographed their dentition as was done for live individuals, with a mean of 11.9 pictures per set. We then extracted a tooth, usually a canine or molar. All extracted teeth were sectioned and aged by the Matson’s Laboratory LLC (Milltown, MT, USA) following the methods described in Grue and Jensen (1976) and Matson and Matson (1981).

Teeth condition index (TCI)

Based on the work of Olifiers et al. (2010) on brown-nosed coati *Nasua nasua* and crab-eating fox *Cerdocyon thous*, we classified teeth condition by tooth type (incisor, canine or premolar/molar) with molar and premolar lumped together because wear appeared similar. Condition was based on tooth state (0 = intact, 1 = if one tooth or more is broken or absent) and wear (0 = sharp, 1 = worn but not flat, 2 = flat; Table 1, Fig. 1). Tooth wear is considered more indicative of age than any other criterion (Wood 1958) but tooth loss and tooth breakage is also age-dependent (Valkenburgh 1988). Tooth condition is not evaluated for each tooth, but rather evaluated across teeth within a given tooth category (e.g. two sharp and two flat canines resulted in a wear index for the tooth category canine of 1; Fig. 1). As shown in Table 1, the TCI is the sum of the various scores described above, and can vary from 0 (excellent teeth condition) to 9 (very bad teeth condition).

In order to evaluate the reliability of the visual assessment of teeth wear and state, we calculated the repeatability of these assessments within and between observers. Five

Table 1. Variables used to build a teeth condition index for arctic foxes *Vulpes lagopus* studied on Bylot Island (Nunavut, Canada) between 2006 and 2013.

Variable		Description
Incisor	state (IS)	complete and intact (0), incomplete or broken (1)
	wear (IW)	sharp (0), worn (1), flat (2)
Canine	state (CS)	complete and intact (0), incomplete or broken (1)
	wear (CW)	sharp (0), worn (1), flat (2)
Premolar/molar	state (MS)	complete and intact (0), incomplete or broken (1)
	wear (MW)	sharp (0), worn (1), flat (2)
Teeth condition index (TCI)		IS + IW + CS + CW + MS + MW




Figure 1. Right lateral (A1, A2, A3), frontal (B1, B2, B3) and left lateral (C1, C2, C3) views of the dentition of three live arctic foxes *Vulpes lagopus* captured on Bylot Island, Nunavut, Canada. Pictures on the top row are from a one year-old arctic fox and show a TCI of 0 (all teeth are sharp and complete). Pictures on the middle row are from an arctic fox of unknown age with worn but complete teeth and show a TCI of 4 (incisives: complete (IS = 0) and worn (IW = 1) ; canines: complete (CS = 0) and worn (CW = 1) ; molars and premolars: complete (MS = 0) and worn (MW = 1); sum (TCI) = 3). Pictures on the bottom row are from an arctic fox of unknown age with flat and incomplete teeth and show a TCI of 9 (incisives: two are absent (IS = 1) and flat (IW = 2); canines: one is broken (CS = 1) and flat (CW = 2); molars and premolars: one is absent (MS = 1) and flat (MW = 2); sum (TCI) = 9).

observers performed the visual assessment of teeth twice for each set of pictures. At least two days elapsed between the assessments of a given set of pictures by a given observer. Order of pictures was random and changed between notation sessions to decrease effects of observer fatigue or learning. Observers were professional biologists but not experts at evaluating teeth condition. All received the same protocol containing an annotated figure similar to Fig. 1. An assessment was considered complete when the state and wear of all tooth types was scored (this was sometimes impossible due to poor quality pictures or tooth loss).

Data analyses

Repeatability can be assessed by the intraclass correlation coefficient (ICC) (Wolak et al. 2012), with

$$ICC = \frac{S_A^2}{S_A^2 + S_W^2}$$

where S_A^2 is the variance among groups and S_W^2 is the variance within groups. To estimate within-observer repeatability, each group was a pair of assessment by a given observer for a given set of pictures. To estimate between-observer repeatability, each group was a set of five assessments for a given set of pictures, one for each observer. Between-observer repeatability was calculated twice, once with the first assessment done by each observer on each set of pictures, and once with the second assessment. ICC varies between 0 (measurements not repeatable) and 1 (measurements fully repeatable). We estimated ICC using the R package ICC (Wolak et al. 2012), with an alpha level of 0.05 to assess the 95% confidence interval.

We used a linear regression to relate real fox age determined by cement lines count or first capture as a cub to our tooth condition index (TCI). We then tested the ability of the resulting model to accurately classify foxes within their respective age categories through jackknife validation. To do

this, we built again our regression model while excluding one fox for which age was to be predicted, repeating this procedure for each fox. When several sets of pictures were available for a given individual due to multiple captures or live capture followed by skull recovery, we used only the first set obtained. Because known ages of foxes were integers, we rounded ages estimated from the model before comparing known and estimated ages. In addition, we tested the ability of the model to classify foxes in age categories, using the following age classes: yearling (1 year-old fox), prime-age adults (2, 3 and 4 years old) and old (≥ 5 years old). All analyses were conducted with the statistical software R (<www.r-project.org>).

Results

Real ages of foxes (≥ 1 year-old) obtained from cement line counts on skulls ranged from 1 to 8 years (mean $\pm$ SD = 3.45 ± 2.04 , $n = 38$). In addition, real ages were known for 6 adult-sized foxes that were captured as cubs in the study area (from 1 to 4 year, mean $\pm$ SD = 2.60 ± 1.03).

Repeatability of the teeth condition index

We succeeded to estimate the TCI of 234 sets of pictures out of 245 (57 out of 66 skulls and 177 out of 179 live captures; Table 2). TCI varied along all possible scores between 0 (excellent condition) and 9 (very bad condition) (median: 3). Within-observer repeatability ranged from 0.83 to 0.96 and averaged 0.89, while between-observer repeatability was 0.81 the first time observers assessed teeth condition and 0.84 the second time (average = 0.83). These repeatability values indicate an excellent precision of the TCI, both within and across observers, with an overall estimation error of 17%, of which 6% was due to differences between observers (0.89–0.83) and 11% to differences within observers (1–0.89).

Table 2. Sample sizes used to build a teeth condition index (TCI) for arctic foxes *Vulpes lagopus* studied on Bylot Island (Nunavut, Canada) between 2006 and 2013.

	Skulls	Captures	Total
Sets of pictures	66	179	245
TCI calculated	57	177	234
Individuals	66	149	205
Individuals of known age (sets of pictures)	63 (63)	13 (19)	69 (82)
Individuals aged through cement lines count (sets of pictures)	63 (63)	7 (12)	63 (75)
Individuals aged through marking as cub (sets of pictures)	0 (0)	6 (7)	6 (7)
TCI with age associated (sets of pictures)	55 (55)	13 (18)	61 (73)
≥ 1 year-old individuals with calculated TCI (sets of pictures)	38 (38)	13 (18)	44 (56)

Validation of the teeth condition index

The linear regression between TCI (x) and real fox age (y) yielded the following predictive equation:

$$y = 0.62 (\pm 0.07 \text{ SE}) x + 1.04 (r^2 = 0.64, n = 44)$$

Tooth wear increased linearly with fox age (Fig. 2). The jack-knife validation indicated that this model predicted correctly the age class (young, prime-age adult, old) of individuals 75% of the time (the age class of 33 out of 44 individuals was correctly assessed; Table 3).

The rate of correct classification decreased to 38.6% (17 out of 44 individuals) when exact age (rather than age class) was predicted. However, in 79.5% (35 of 44) of the cases, the model estimate was within one year of the true age, and in 100% of the cases it was within two years of true age.




Figure 2. Linear regression (with standard error, dotted lines) between the known age in years (y) and the mean of the teeth condition index (x) for 44 individuals in an arctic fox *Vulpes lagopus* population studied on Bylot Island, Nunavut, Canada. Regression equation: $y = 0.62x + 1.04$ ($r^2 = 0.64$, $p < 0.001$). Size of data points in the plot reflects the number of observations ($n = 1$ or $n = 2$).

Table 3. Comparison of known age with age estimated from the teeth condition index for 44 live arctic foxes *Vulpes lagopus* studied on Bylot Island (Nunavut, Canada). Perfect match between estimates and real ages appears in bold.

		Age predicted by the model					
		Yearling	Prime-age adults				Old
		1	2	3	4	5	≥ 6
Known age	yearling	1	4	3	1	0	0
	prime-age adults	2	2	5	3	1	0
		3	1	2	4	3	1
		4	0	1	0	2	1
old		5	0	0	0	1	2
		6	0	0	0	1	3
		7	0	0	0	0	2

Discussion

Our proposed teeth condition index allowed an estimation of the age of individuals captured from a wild population of carnivores without relying on tooth extraction. Use of distinguishable wear categories following Olifiers et al. (2010) allowed us to limit subjectivity when evaluating tooth attrition, which likely improved the repeatability of our assessments. Moreover, the constant increase of tooth attrition with fox age indicated the relevance of a linear model to predict age of individuals, whereas the strong relation ($r^2 = 0.64$) between tooth wear notation and known age indicates that individuals could be reliably aged from tooth wear in our studied population. For comparison, the r^2 of the rank regression between tooth wear and cement lines count was only 0.10 for males and 0.35 for females in brushtail possums (Lohr et al. 2011).

The model predicted the exact age of 39% of foxes, an estimated age that was within one year of true age for 77.3% of foxes and the correct age category (yearling, prime-age adult, old adult) for 75% of individuals. Such a level of accuracy is higher than that achieved in other studies where individuals were aged from tooth wear by multiple observers. In roe deer, the age estimated by tooth wear of 50% of individuals was within one year of true age (Hewison 1999), and in gray wolf, the estimated age of 75% of individuals was within two years of true age (Gipson et al. 2000). The accuracy for age categories matches the results obtained for crab-eating foxes by Olifiers et al. (2010), where 67 to 100% of individuals were correctly classified according to definite age categories.

We observed a maximum age of 8 years in our population, while the oldest individuals were 7 or 8 years-old in other Canadian populations (Bradley et al. 1981, Hiruki and Stirling 1989), 8 years-old on Saint Lawrence Island in Alaska (Fay and Rausch 1992), and 10 years-old in Iceland (Hersteinsson 1992) and Svalbard (Eide et al. 2012). This indicates that maximum longevity seems to be a rather constant parameter across arctic fox populations. Published population age structures of arctic foxes were inferred from multiple aging techniques and presented using age classes defined in multiple ways (Macpherson 1969, Bradley et al. 1981, Hiruki and Stirling 1989, Fay and Rausch 1992, Hersteinsson 1992, Pagh and Hersteinsson 2008, Eide et al.

2012). However, these population age structures were all based on analyses of carcasses from fur trapping stations (Macpherson 1969, Bradley et al. 1981, Hiruki and Stirling 1989, Fay and Rausch 1992, Hersteinsson 1992, Pagh and Hersteinsson 2008, Eide et al. 2012), and were therefore all likely biased toward young individuals given that this age class is easier to trap (Smirnov 1968 in Bradley et al. 1981, Hiruki and Stirling 1989). In addition, age structures built from dead individuals originating from multiple cohorts do not necessarily represent the age structure of live individuals in the population. Our method, which estimates the age of live animals, may therefore help establish more precise age structures for arctic fox populations.

Age evaluation based on external teeth condition offers promising avenues for wildlife biology as it is relatively cheap, can be done without specialized equipment or detailed expertise, and is less invasive than most other aging methods. A better knowledge of the tooth wear patterns of individuals is needed, however, to improve predictive models. This can be obtained through a combination of long-term monitoring of known-aged individuals, systematic photography of dentition of captured animals, and development of teeth condition indexes adapted to each species. One difficulty, however, is that calibration of tooth wear patterns must be done at the population level, because various diets might generate various tooth wear patterns (Morris 1972, Hewison 1999, Ozaki et al. 2010). For example, Daitch and Guralnick (2007) found a difference of tooth shape among circumpolar populations of arctic foxes, which could be explained by dietary differences (see also Szuma 2008). Another difficulty is that tooth wear might vary through time in some populations. For example, it is possible that the between-individual variation in tooth wear observed in our study was partly due to the phases of the lemming cycle that individuals were exposed to during their lifetime, given that lemming abundance strongly influences the diet of many tundra predators (Legagneux et al. 2012). Sex could also impact tooth wear in some species (Coy and Garshelis 1992, van Deelen et al. 2000) but we could not test this due to sample size limitation. However, the diet of male and female arctic foxes is generally similar (Pagh and Hersteinsson 2008, Tarroux et al. 2012), so we do not expect any sex-related difference in tooth wear for this species. Clearly, both fundamental research on the behaviour and population dynamics of several wildlife species and applied research on their management and conservation would benefit from advancing aging methods.

Acknowledgements – We thank the many field workers who collected data on Bylot Island, and Sandra Lai, Nicolas Casajus, Marie-Jeanne Rioux, Camille Morin and Florence Lapierre-Poulin who assessed tooth state and wear. Sandra Lai and Nicolas Casajus provided statistical advice.

Funding – This study was supported by (alphabetical order): Canada Foundation for Innovation, Canada Research Chairs Program, Fonds de recherche du Québec - Nature et technologies (FRQNT), International Polar Year program of Indian and Northern Affairs Canada, Kenneth M Molson Foundation, Mittimatalik Hunters and Trappers Organization, Natural Sciences and Engineering Research Council of Canada (NSERC), Network of Centers of Excellence of Canada ArcticNet, Northern Ecosystem Initiative (Environment Canada), Northern Scientific Training Program

(Indian and Northern Affairs Canada), Nunavut Wildlife Management Board, Parks Canada Agency, Polar Continental Shelf Program (Natural Resources Canada), and Université du Québec à Rimouski (UQAR).

Permits – All animals were treated and handled according to permits: CPA15-02-01 : 2003-2007, CPA32-08-62 : 2008-2011, CPA48-12-100 : 2012-2013.

References

- Bêty, J. and Gauthier, G. 2002. Shared predators and indirect trophic interactions: lemming cycles and arctic-nesting geese. – *J. Anim. Ecol.* 71: 88–98.
- Bradley, J. et al. 1981. Age determination in the arctic fox (*Alopex lagopus*). – *Can. J. Zool.* 59: 1976–1979.
- Careau, V. et al. 2008. Hoarding of pulsed resources: temporal variations in egg-caching by arctic fox. – *Ecoscience* 15: 268–276.
- Cavallini, P. and Santini, S. 1995. Age determination in the red fox in a Mediterranean habitat. – *Int. J. Mamm. Biol.* 60: 136–142.
- Christensen-Dalsgaard, S. N. et al. 2009. Accuracy and precision in estimation of age of Norwegian Arctic polar bears (*Ursus maritimus*) using dental cementum layers from known-age individuals. – *Polar Biol.* 33: 589–597.
- Coy, P. and Garshelis, D. 1992. Reconstructing reproductive histories of black bears from the incremental layering in dental cementum. – *Can. J. Zool.* 70: 2150–2160.
- Daitch, D. and Guralnick, R. 2007. Geographic variation in tooth morphology of the arctic fox, *Vulpes (Alopex) lagopus*. – *J. Mammal.* 88: 384–393.
- Delahay, R. et al. 2011. Using lifetime tooth-wear scores to predict age in wild Eurasian badgers: performance of a predictive model. – *J. Zool.* 284: 183–191.
- Eide, N. E. et al. 2012. Reproductive responses to spatial and temporal prey availability in a coastal Arctic fox population. – *J. Anim. Ecol.* 81: 640–648.
- Fay, F. and Rausch, R. 1992. Dynamics of the arctic fox population on St. Lawrence Island, Bering Sea. – *Arctic* 45: 393–397.
- Gagnon, C. and Berteaux, D. 2009. Integrating traditional ecological knowledge and ecological science: a question of scale. – *Ecol. Soc.* 14: 19.
- Gauthier, G. et al. 2004. Trophic interactions in a high arctic snow goose colony. – *Integr. Comp. Biol.* 44: 119–129.
- Gay, S. and Best, T. 1996. Age-related variation in skulls of the puma (*Puma concolor*). – *J. Mammal.* 77: 191–198.
- Gipson, P. et al. 2000. Accuracy and precision of estimating age of gray wolves by tooth wear. – *J. Wildl. Manage.* 64: 752–758.
- Giroux, M.-A. et al. 2012. Benefiting from a migratory prey: spatio-temporal patterns in allochthonous subsidization of an Arctic predator. – *J. Anim. Ecol.* 81: 533–542.
- Grau, G. et al. 1970. Age determination of raccoons. – *J. Wildl. Manage.* 34: 364–372.
- Grue, H. and Jensen, B. 1976. Annual cementum structures in canine teeth in arctic foxes (*Alopex lagopus* (L.)) from Greenland and Denmark. – *Dan. Rev. Game Biol.* 10: 1–16.
- Grue, H. and Jensen, B. 1979. Review of the formation of incremental lines in tooth cementum of terrestrial mammals. – *Dan. Rev. Game Biol.* 11: 3–44.
- Hamlin, K. L. et al. 2000. Evaluating the accuracy of ages obtained by two methods for Montana ungulates. – *J. Wildl. Manage.* 64: 441–449.
- Harris, S. 1978. Age determination in the red fox (*Vulpes vulpes*) – an evaluation of technique efficiency as applied to a sample of suburban foxes. – *J. Zool.* 184: 91–117.

- Harris, S. et al. 1992. Age determination of badgers (*Meles meles*) from tooth wear: the need for a pragmatic approach. – J. Zool. 228: 679–684.
- Hersteinsson, P. 1992. Demography of the arctic fox (*Alopex lagopus*) population in Iceland. – In: McGullogh, D. R. and Barrett, R. H. (eds), Wildlife 2001: populations. Springer, pp. 954–964.
- Hewison, A. 1999. Tests of estimation of age from tooth wear on roe deer of known age: variation within and among populations. – Can. J. Zool. 77: 58–67.
- Hiruki, L. and Stirling, I. 1989. Population dynamics of the arctic fox, *Alopex lagopus*, on Banks Island, Northwest Territories. – Can. Field-Nat. Ottawa 103: 380–387.
- King, C. 1991. A review of age determination methods for the stoat *Mustela erminea*. – Mamm. Rev. 21: 31–49.
- Landon, D. et al. 1998. Evaluation of age determination techniques for gray wolves. – J. Wildl. Manage. 62: 674–682.
- Legagneux, P. et al. 2012. Disentangling trophic relationships in a High Arctic tundra ecosystem through food web modeling. – Ecology 93: 1707–16.
- Linhart, S. and Knowlton, F. 1967. Determining age of coyotes by tooth cementum layers. – J. Wildl. Manage. 31: 362–365.
- Lohr, C. A. et al. 2011. Estimating the age structure of a population of brushtail possums (*Trichosurus vulpecula*) by comparing tooth wear and cementum lines. – Aust. Mammal. 33: 52–56.
- Macpherson, A. 1969. The dynamics of Canadian arctic fox populations. – Can. Wildl. Serv. Rep. Ser. 8: 1–50.
- Matson, G. and Matson, J. 1981. Workbook for cementum analysis. – Matson's.
- Morris, P. 1972. A review of mammalian age determination methods. – Mamm. Rev. 2: 69–104.
- Olifiers, N. et al. 2010. Estimating age of carnivores from the Pantanal region of Brazil. – Wildl. Biol. 16: 389–399.
- Ozaki, M. et al. 2010. The relationship between food habits, molar wear and life expectancy in wild sika deer populations. – J. Zool. 280: 202–212.
- Pagh, S. and Hersteinsson, P. 2008. Difference in diet and age structure of blue and white Arctic foxes (*Vulpes lagopus*) in the Disko Bay area, West Greenland. – Polar Res. 27: 44–51.
- Pekelharing, C. 1970. Cementum deposition as an age indicator in the brush-tailed possum, *Trichosurus vulpecula* Kerr (Marsupiala). – Aust. J. Zool. 18: 71–76.
- Proulx, G. et al. 1994. Field evaluation of the number 1½ steel-jawed leghold and the Sauvageau 2001-8 traps to humanely capture arctic fox. – Wildl. Soc. Bull. 22: 179–183.
- Rosatte, R. and Silver, A. 2007. Correlation between age estimates for elk, *Cervus elaphus*, using tooth wear/eruption patterns and counts of annuli in tooth cementum. – Can. Field-Nat. 121: 214–215.
- Santymire, R. M. et al. 2012. Using canine width to determine age in the black-footed ferret (*Mustela nigripes*). – Small Carniv. Conserv. 46: 17–21.
- Shimoinaba, S. and Oi, T. 2015. Relationship between tooth wear and age in the Japanese black bear in Hiroshima Prefecture, Japan. – Mammal Study 40: 53–60.
- Smirnov, V. S. 1968. Analysis of arctic fox population dynamics and methods of increasing the arctic fox harvest. – Probl. North 11: 81–101.
- Smuts, G. et al. 1978. Age determination of the African lion (*Panthera leo*). – J. Zool. 185: 115–146.
- Spinage, C. 1973. A review of the age determination of mammals by means of teeth, with especial reference to Africa. – Afr. J. Ecol. 11: 165–187.
- Stander, P. 1997. Field age determination of leopards by tooth wear. – Afr. J. Ecol. 35: 156–161.
- Szuma, E. 2008. Geographic variation of tooth and skull sizes in the arctic fox *Vulpes (Alopex) lagopus*. – Ann. Zool. Fenn. 45: 185–199.
- Tarroux, A. et al. 2012. The marine side of a terrestrial carnivore: intra-population variation in use of allochthonous resources by arctic foxes. – PLoS ONE 7: e42427.
- Valkenburgh, B. Van 1988. Incidence of tooth breakage among large, predatory mammals. – Am. Nat. 131: 291–302.
- van Deelen, T. R. et al. 2000. Sex affects age determination and wear of molariform teeth in white-tailed deer. – J. Wildl. Manage. 64: 1076–1083.
- Van Horn, R. C. et al. 2003. Age estimation and dispersal in the spotted hyena (*Crocuta crocuta*). – J. Mammal. 84: 1019–1030.
- Wolak, M. E. et al. 2012. Guidelines for estimating repeatability. – Meth. Ecol. Evol. 3: 129–137.
- Wood, J. 1958. Age structure and productivity of a gray fox population. – J. Mammal. 39: 74–86.