

First record of the genus *Beierochelifer* (Pseudoscorpiones: Cheliferidae) from north-western Iran

Authors: Nassirkhani, Mahrads, and Doustaresharaf, Mojtaba
Mohammad

Source: Arachnologische Mitteilungen: Arachnology Letters, 59(1) : 30-34

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit5905>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of the genus *Beierochelifer* (Pseudoscorpiones: Cheliferidae) from north-western Iran

Mahrad Nassirkhani & Mojtaba Mohammad Doustaresharaf



doi: 10.30963/aramit5905

Abstract. *Beierochelifer peloponnesiacus* (Beier, 1929) is recorded from Iran for the first time. The species is briefly described and its diagnostic characters are illustrated. Furthermore, a key for recognizing males of the genera of the family Cheliferidae Risso, 1827 reported from Iran is provided.

Keywords: Arachnida, distribution, identification key, Middle East, taxonomy

Zusammenfassung. Erster Nachweis der Gattung *Beierochelifer* (Pseudoscorpiones: Cheliferidae) aus den Nordwesten Irans. *Beierochelifer peloponnesiacus* (Beier, 1929) wird erstmals für den Iran nachgewiesen. Die Art wird kurz beschrieben und ihre diagnostischen Merkmale werden abgebildet. Zusätzlich wird ein Schlüssel für die Gattungen der Familie Cheliferidae Risso, 1827 im Iran erstellt.

The pseudoscorpion genus *Beierochelifer* Mahnert, 1977 is distributed mostly in the Mediterranean region, but – less frequently – was also recorded in the Middle East and its adjacent areas. *Beierochelifer peloponnesiacus* (Beier, 1929) currently contains two subspecies: *B. p. peloponnesiacus* (Beier, 1929), which was originally described from Greece and subsequently reported from Azerbaijan, Bulgaria, Italy and Slovakia, and *B. p. jonius* (Beier, 1932), known from Greece, Spain and Turkey (Harvey 2013, Krajčovičová & Christophoryová 2017, Hernández-Corral et al. 2018).

Brief descriptions and drawings of the pedipalp and tarsus I of this species were presented by Beier (1929, 1932, 1963), Dashdamirov & Schawaller (1992) and Petrov (2004) for the specimens collected from Greece, Azerbaijan and Bulgaria. In addition, the only known figure of the statumen convolutum of *B. p. jonius* was drawn by Mahnert (1977) based on males from Greece. Recently, Krajčovičová & Christophoryová (2017) provided a short description of *B. p. peloponnesiacus* based on two males from Slovakia. The authors figured the pedipalpal segments, details of the pleural membrane, and tarsus I and IV. Moreover, Hernández-Corral et al. (2018) prepared a description of *B. p. jonius* based on the adults from Spain, but did not provide any illustrations.

According to Beier (1963), the subspecies of *B. peloponnesiacus* can only be separated by small morphometric differences between the pedipalpal segments. Therefore the determination of the subspecies is rather difficult and unreliable. In this contribution, the specimens from Iran are considered at the specific level. The illustrations presented here depict males recently found in Iran. The presented locality is thus the easternmost record of the known species range.

A total of seven genera of the family Cheliferidae Risso, 1827 have been reported from Iran including the present study (Beier 1951, 1971, Mahnert 1974, Schawaller 1983, Judson 1990, Nassirkhani & Takaloo zade 2012, 2013a, 2013b,

Nassirkhani & Shoushtari 2014, 2015, Nassirkhani et al. 2016). An identification key to these genera for recognizing the males is provided, together with the provincial distribution of the cheliferid species in Iran (Fig. 1).

Material and methods

The specimens were found in soil and litter at a depth of 6–10 cm. They were collected by sieving samples, and removed by hand with the thin needle of a hypodermic syringe. They were permanently mounted in Swann's fluid, examined with an Olympus CH-2 compound microscope, illustrated with a drawing tube attached to the microscope, and measured by an ocular graticule. Photographs were made using a digital camera (Canon PC1468). All measurements are expressed in millimetres; the given ratios are length/width for individual articles and the length/width for legs; when two articles are compared, the ratio is the length/length index. Morphological terminology and measurements follow Chamberlin (1931), Harvey (1992), Harvey et al. (2012) and Judson (2007). Terminology for chelal lyrifissures follows Zaragoza (2017). Coordinates are given in the Geodetic System WGS 84. The specimens are deposited in Collection of Acarology Laboratory, Islamic Azad University of Arak, Iran (IAUA).

Abbreviations

<i>dat</i> = dorsal accessory tooth	<i>ls</i> = lateral seta
<i>ddp</i> = dorso-distal projection	<i>ma₁</i> = retrolateral lyrifissures
<i>ds</i> = discal seta	<i>ma₂</i> of movable
<i>fa</i> = retrolateral lyrifissure of fixed chelal finger,	<i>ma₃</i> chelal finger
<i>fb</i> = dorso-retrolateral lyrifissure of fixed chelal finger,	<i>ms</i> = median seta
<i>fd</i> = dorso-distal lyrifissure of fixed chelal finger	<i>sdt</i> = serrate dorsal teeth
<i>L</i> = length	<i>sc</i> = statumen convolutum
	<i>se</i> = sensillum
	<i>T</i> = tactile seta
	<i>W</i> = width


Trichobothriotaxy as in Chamberlin (1931):

<i>eb</i> = external basal	<i>ist</i> = internal sub-terminal
<i>esb</i> = external sub-basal	<i>it</i> = internal terminal
<i>est</i> = external sub-terminal	<i>t</i> = terminal
<i>et</i> = external terminal	<i>st</i> = sub-terminal
<i>ib</i> = internal basal	<i>sb</i> = sub-basal
<i>isb</i> = internal sub-basal	<i>b</i> = basal


Mahrad NASSIRKHANI, Entomology Department, Faculty of Agriculture and Natural Resources, Islamic Azad University, Arak branch, Arak, Iran;
E-mail: greenartificialturfgrass@gmail.com
Mojtaba Mohammad DOUSTARESHARAF, Department of Plant Protections, Faculty of Agriculture, University of Maragheh, Maragheh, Iran;
E-mail: mojtaba.doostar@gmail.com

Academic editor: Theo Blick

submitted 25.6.2019, accepted 11.2.2020, online 6.3.2020



Chelicera: hand with 5 setae, *b* and *sb* denticulate, the other setae smooth (Fig. 7); rallum with 3 blades, distalmost blade laterally denticulate, median blade with a few (1–2) lateral denticulations and proximalmost blade smooth (Fig. 8) [in one specimen, proximalmost blade with some lateral denticulations (Fig. 9)].



Figures 2–16: *Beierochelifer peloponnesiacus* (Beier, 1929), ♂. **2.** carapace, dorsal view (some setae magnified, right half showing granulation pattern, left half showing chaetotaxy, position of furrows and lyrifissure); **3.** left hemitergites III–IV, dorsal view; **4.** right coxae and sternites II–III, ventral view; **5.** internal genitalia, in part; **6.** statumen convolutum, magnified (microscopic picture); **7.** left chelicera, dorsal view; **8.** rallum, magnified (showing proximalmost blade smooth); **9.** rallum, magnified (showing proximalmost blade denticulate); **10.** right pedipalp (chela omitted), dorsal view (one seta magnified); **11.** right chela, dorsal view (some setae magnified); **12.** right chela, retrolateral view; **13.** tibia and tarsus I, retrolateral view; **14.** posterior claw of tarsus I, magnified; **15.** distal part of tarsus I, retrolateral view (microscopic picture); **16.** tarsus IV, retrolateral view. See Material and methods for abbreviations

Pedipalps: granulation pattern as in Figs 10–11; trochanter 1.50–1.73×; femur 2.32–2.61×; patella with short and stout pedicel (L = 0.25–0.27 mm), 2.16–2.21×; Chela with pedicel 3.60–3.86×, without pedicel 3.32–3.57×; hand with pedicel 2.22–2.28×, without pedicel 1.95–2.04×; hand (with pedicel) 1.32–1.39× longer than movable finger; hand (without pedicel) 1.14–1.23× longer than movable finger; trichobothriotaxy as in Figs 11–12.

Legs: coxal chaetotaxy: 8:7–8:7–8:18–19; sub-terminal setae simple; aroliae shorter than claws. Leg I: femur 1.61–1.69×; patella 2.26–2.35×; patella 1.35–1.38× longer than femur; tibia 2.44–2.55×; tarsus 2.10–2.27× [in one specimen 2.40×]; claws asymmetric, posterior claw with a dorsal accessory tooth, which bears some serrate dorsal teeth with different sizes (Figs 13–15). Leg IV: femur 1.60–1.75×; patella 2.62–2.84×; femur + patella 3.06–3.39×; tibia 3.70–3.89×; tarsus with a tactile seta situated distinctly distal to middle of the segment (T = 0.64–0.67), 3.69–3.85×; claws symmetric and smooth (Fig. 16).

Measurements (mm): Body length: 3.75–4.07. Carapace: 1.10–1.15/0.97–1.05. Pedipalp: trochanter 0.58–0.60/0.34–0.40; femur 1.00–1.10/0.39–0.44; patella 0.93–0.98/0.43–0.46; chela (with pedicel) 1.62–1.72/0.42–0.46; chela (without pedicel) 1.50–1.57; hand (with pedicel) L. 0.96–1.02; hand (without pedicel) L. 0.86–0.89; movable finger L. 0.71–0.77. Leg I: femur 0.37–0.39/0.23; patella 0.50–0.54/0.22–0.23; tibia 0.44–0.46/0.18; tarsus 0.40–0.41/0.17–0.19. Leg IV: femur 0.32–0.35/0.20–0.21; patella 0.84–0.88/0.31–0.32; femur + patella 0.98–1.05; tibia 0.74–0.75/0.19–0.20; tarsus 0.48–0.50/0.13.

Discussion

Beierochelifera peloponnesiacus (Beier, 1929) can be easily separated from *B. anatolicus* (Beier, 1949) from Turkey and Greece, and from *B. geoffroyi* Heurtault, 1981 from France on the basis of the shape of the male tarsus I and the morphometric characteristics. In *B. anatolicus*, the male tarsus I has no dorso-distal projection (see Beier 1965: fig. 11). In *B. geoffroyi*, the dorso-distal projection of the male tarsus I is not conspicuous and not clearly protruded (see Heurtault 1981: fig. 5). Comparatively, the male tarsus I of *B. peloponnesiacus* bears a very robust and prominent dorso-distal projection (see Beier 1929: fig. 11B, 1932: figs 287, 289, 1963: figs 299–300, Krajčovičová & Christophoryová 2017: fig. 5, present study: Fig. 13).

Moreover, *B. peloponnesiacus* is clearly larger than the other *Beierochelifera* species, e.g. the pedipalpal femur length is 0.77–1.00 mm, the patellar length is 0.71–0.95 mm, and length of the movable chelal finger is 0.54–0.69 mm for *B. peloponnesiacus* (Petrov 2004, Krajčovičová & Christophoryová 2017, Hernández-Corral et al. 2018), while these are 0.59–0.69 mm, 0.54–0.65 mm, and 0.38–0.44 mm for *B. anatolicus* and *B. geoffroyi* respectively (Beier 1949, Heurtault 1981, Petrov 2004). The newly collected specimens from Iran are slightly larger than the specimens of *B. peloponnesiacus* previously recorded from different localities, e.g. the pedipalpal femur length is 1.00–1.10 mm, and the length of the movable chelal finger is 0.71–0.77 mm.

Noticeably, the granulation pattern of the pedipalpal is different between the specimens studied before under *B. peloponnesiacus*. The pedipalpal trochanter and femur are entirely,

and the patella is partly, granulated and the chela is completely smooth in the specimens from Bulgaria (judging from Petrov 2004: fig. 9). The pedipalpal segments, even the chelal fingers, are entirely granulated in *B. aff. peloponnesiacus* from Azerbaijan (judging from Dashdamirov & Schawaller 1992: fig. 13D). Moreover, both surfaces of the pedipalpal femur and the prolateral surface of the patella and the chelal hand are granulated in the specimens from Spain (Petrov 2004, Hernández-Corral et al. 2018). In the newly collected specimens from Iran, only the retrolateral surface of each pedipalpal segment is partially, and the chelal fingers are entirely smooth. The other parts of the pedipalp are distinctly granulated (Figs 11–12).

The other important regional difference between the specimens is the shape of the statumen convolutum of the male internal genitalia, e.g. it has apically a shallow incision in the specimens from Greece (see Mahnert 1977: fig. 12a), whereas it is apically rounded and without any notch in the newly found specimens from Iran (Figs 5–6).

The position of the trichobothrium *isb* varies within the species, e.g. *isb* is located at the same level as *it* in the adults from southern and the male from western Greece (see Beier 1932: figs 286, 288), while it is located proximal to *it* in the female from western Greece (see Beier 1932: fig. 288), the female from Azerbaijan (see Dashdamirov & Schawaller 1992: fig. 13D), and the newly collected males from Iran (Figs 11–12).

Beierochelifera peloponnesiacus can be mostly found under bark of *Quercus* spp. and *Platanus* sp. (Petrov 2004, Krajčovičová & Christophoryová 2017), and it was rarely collected from litter or tree hollows of *Quercus* spp., *Q. cerris* L. and *Q. pyrenaica* Willd (Petrov 2004, Krajčovičová & Christophoryová 2017, Hernández-Corral et al. 2018). Herein, the males were collected from soil and litter consisting of leaf and wood pieces of *Quercus macranthera* Fisch. & Mey. ex Hohen. and *Quercus petraea* (Matt.) Liebl.

Key to the genera of the family Cheliferidae Risso, 1827 recorded from Iran (males)

Many cheliferids can only be identified with certainty at the generic or specific level according to the male characters. Therefore, the present key is provided for recognizing males of the cheliferid genera from Iran.

1. Statumen convolutum with a deep apical recess and a sclerotized rod in the middle of this indentation [tribe Cheliferini Risso, 1827] 2
- Statumen convolutum apically rounded, not deeply indented, and without sclerotized rod [tribe Dactylocheliferini Beier, 1932] 3
2. Pedipalpal femur and patella with large and conspicuous conical keels *Strobilochelifera* Beier, 1932
- Pedipalpal femur and patella without large conical keels *Hysterochelifera* Chamberlin, 1932
3. Movable chelal finger with 3 trichobothria *Ellingsenius* Chamberlin, 1932
- Movable chelal finger with 4 trichobothria 4
4. Sub-terminal setae of pedal tarsus I–IV simple 5
- Sub-terminal setae of pedal tarsus I–IV denticulate *Rhacochelifera* Beier, 1932
5. Coxal sac present in coxae IV; tarsus IV without a tactile seta *Dactylochelifera* Beier, 1932

- Coxal sac absent in coxae IV; tarsus IV with a long/slightly long tactile seta 6
- 6. Claws of tarsus I asymmetric, one claw with a large smooth ventral accessory tooth; all setae on cheliceral hand simple *Gobichelifer* Krumpál, 1979
- Claws of tarsus I asymmetric, one claw with a large dorsal accessory tooth plus some small teeth on its outer edge; *b* and *sb* on cheliceral hand denticulate *Beierochelifer* Mahnert, 1977

Acknowledgements

The authors wish to thank Dr. Reza Vafai Shoushtari (IAUA) for his support and Mr. Mahmoud Nassirkhani for his assistance.

References

- Beier M 1929 Die Pseudoscorpione des Wiener Naturhistorischen Museums. II. Panctenodactyli. – Annalen des Naturhistorischen Museums in Wien 43: 341-367
- Beier M 1932 Pseudoscorpionidea II. Subord. C. Cheliferinea. – Das Tierreich 58: 1-294
- Beier M 1949 Türkiye Pseudoscorpion'lari hakkında. Türkische Pseudoscorpione. – Revue de la Faculté des Sciences de l'Université d'Istanbul (B) 14: 1-20
- Beier M 1951 Ergebnisse der Österreichischen Iran-Expedition 1949/50. Pseudoscorpione und Mantiden. – Annalen des Naturhistorischen Museums in Wien 58: 96-102
- Beier M 1963 Ordnung Pseudoscorpionidea (Afterscorpione). Bestimmungsbücher zur Bodenfauna Europas, Band 1. Akademie-Verlag, Berlin. 313 pp.
- Beier M 1965 Anadolu'nun Pseudoscorpion faunası. Die Pseudoscorpioniden-Fauna Anatoliens. – Istanbul Üniversitesi Fen Fakültesi Mecmuası 29(B): 81-105
- Beier M 1971 Pseudoscorpione aus dem Iran. – Annalen des Naturhistorischen Museums in Wien 76: 357-366
- Chamberlin JC 1931 The arachnid order Chelonethida. – Stanford University Publications, Biological Sciences 7(1): 1-284
- Dashdamirov S & Schawaller W 1992 Pseudoscorpions of the Caucasian fauna (Arachnida: Pseudoscorpionida). – Arthropoda Selecta 1(4): 31-72.
- Harvey MS 1992 The phylogeny and classification of the Pseudoscorpionida (Chelicerata: Arachnida). – Invertebrate Taxonomy 6: 1373-1435 – doi: [10.1071/IT9921373](https://doi.org/10.1071/IT9921373)
- Harvey MS 2013 Pseudoscorpions of the World, version 3.0. Western Australian Museum, Perth. – Internet: <http://www.museum.wa.gov.au/catalogues-beta/pseudoscorpions> (17. Jan. 2019)
- Harvey MS, Ratnaweera PB, Udagama PV & Wijesinghe MR 2012 A new species of the pseudoscorpion genus *Megachernes* (Pseudoscorpiones: Chernetidae) associated with a threatened Sri Lankan rainforest rodent, with a review of host associations of *Megachernes*. – Journal of Natural History 46: 2519-2535 – doi: [10.1080/00222933.2012.707251](https://doi.org/10.1080/00222933.2012.707251)
- Hernández-Corral J, Zaragoza JA & Micó E 2018 New species of Pseudoscorpiones (Arachnida) from tree hollows in a Mediterranean oak forest in Spain. – Zootaxa 4497: 201-225 – doi: [10.11646/zootaxa.4497.2.3](https://doi.org/10.11646/zootaxa.4497.2.3)
- Heurtault J 1981 Présence et signification dans la France méditerranéenne des espèces des genres *Beierochelifer*, *Cheirochelifer* et *Calocheiridius* (Arachnides, Pseudoscorpions). – Atti della Società Toscana di Scienze Naturali, Memorie B 88: 209-222
- Judson MLI 1990 Redescription of the bee-associate *Ellingsenius fulleri* (Hewitt and Godfrey) (Arachnida, Chelonethi, Cheliferidae) with new records from Africa, Europe and the Middle East. – Journal of Natural History 24: 1303-1310 – doi: [10.1080/00222939000770771](https://doi.org/10.1080/00222939000770771)
- Judson MLI 2007 A new and endangered species of the pseudoscorpion genus *Lagynochthonius* from a cave in Vietnam, with notes on chelal morphology and the composition of the Tyrannochthoniini (Arachnida, Chelonethi, Chthoniidae). – Zootaxa 1627: 53-68 – doi: [10.5281/zenodo.179321](https://doi.org/10.5281/zenodo.179321)
- Krajčovičová K & Christophoryová J 2017 First record of *Beierochelifer* Mahnert, 1977 (Pseudoscorpiones: Cheliferidae) from Slovakia. – Check List 13 (2, 2017): 1-4 – doi: [10.15560/13.2.2074](https://doi.org/10.15560/13.2.2074)
- Mahnert V 1974 *Roncus viti* n. sp. (Arachnida, Pseudoscorpiones) aus dem Iran. – Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck 61: 78-91
- Mahnert V 1977 Über einige Atemnidae und Cheliferidae Griechenlands (Pseudoscorpiones). – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 50: 67-74 – doi: [10.5169/seals-401837](https://doi.org/10.5169/seals-401837)
- Nassirkhani M & Takalloo zade HM 2012 First report of two species of false-scorpions (Pseudoscorpions–Arachnida) from Kerman province. In: 17th National & 5th International Biology Conference; 4-6 September 2012, Kerman, Iran. 4 pp. – Internet: <http://www.takalloozadeh.ir/upload/pdf/paper-mahrad.pdf> (31. Jan. 2020)
- Nassirkhani M & Shoushtari RV 2014 Two new records of the genus *Dactylochelifer* (Pseudoscorpiones: Cheliferidae) from Iran. – Acta Societatis Zoologicae Bohemicae 78: 219-227
- Nassirkhani M & Shoushtari RV 2015 A redescription of *Rhacochelifer melanopygus* (Redikorzev 1949) (Pseudoscorpiones: Cheliferidae), a new record of pseudoscorpion fauna of Iran. – Acta Arachnologica 64: 21-25 – doi: [10.2476/asjaa.64.21](https://doi.org/10.2476/asjaa.64.21)
- Nassirkhani M, Shoushtari RV & Rahmat Abadi M 2016 Notes on *Hysterochelifer afghanicus* Beier (Pseudoscorpiones: Cheliferidae), with a key to the species of *Hysterochelifer* Chamberlin. – Zoology in the Middle East 62: 78-84 – doi: [10.1080/09397140.2016.1144281](https://doi.org/10.1080/09397140.2016.1144281)
- Nassirkhani M & Takalloo zade HM 2013a The first report of *Dactylochelifer spasskyi* Redikorzev (Pseudoscorpiones: Cheliferidae) from Iran. – Journal of Entomological Research 4: 335-340
- Nassirkhani M & Takalloo zade HM 2013b The first report of *Gobichelifer chelanops* (Pseudoscorpiones: Cheliferidae) from Iran. – Acta Arachnologica 62: 13-18 – doi: [10.2476/asjaa.62.13](https://doi.org/10.2476/asjaa.62.13)
- Petrov BP 2004 The false scorpions (Arachnida: Pseudoscorpiones) of the Eastern Rhodopes (Bulgaria and Greece). In: Beron P & Popov A (eds) Biodiversity of Bulgaria 2. Biodiversity of Eastern Rhodopes (Bulgaria and Greece). Pensoft and National Museum of Natural History, Sofia. pp. 153-166
- Schawaller W 1983 Pseudoscorpione aus dem Norden des Iran (Arachnida: Pseudoscorpiones). – Senckenbergiana Biologica 63: 367-371
- Zaragoza JA 2017 Revision of the *Ephippiochthonius* complex (Pseudoscorpiones, Chthoniidae) in the Iberian Peninsula, Balearic Islands and Macaronesia, with proposed changes to the status of the *Chthonius* subgenera. – Zootaxa 4246: 1-221 – doi: [10.11646/zootaxa.4246.1.1](https://doi.org/10.11646/zootaxa.4246.1.1)