

Seasonal Diet of the Grey Heron *Ardea cinerea* on an Oceanic Island (Tenerife, Canary Islands): Indirect Interaction with Wild Seed Plants

Authors: Rodriguez, Airam, Rodríguez, Beneharo, Rumeu, Beatriz, and Nogales, Manuel

Source: *Acta Ornithologica*, 42(1) : 77-87

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/068.042.0102>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Seasonal diet of the Grey Heron *Ardea cinerea* on an oceanic island (Tenerife, Canary Islands): indirect interaction with wild seed plants

Airam RODRÍGUEZ^{1,2}, Beneharo RODRÍGUEZ³, Beatriz RUMEU¹ & Manuel NOGALES¹

¹Island Ecology and Evolution Research Group, Instituto de Productos Naturales y Agrobiología (IPNA-CSIC), Astrofísico Francisco Sánchez 3, 38206 La Laguna, Tenerife, Canary Islands, SPAIN, e-mail: airamrguez@ipna.csic.es

²Current address: Department of Evolutionary Ecology, Estación Biológica de Doñana (IPNA-CSIC), Pabellón del Perú, Avda. María Luisa S/N, 41013 Seville, SPAIN

³C/ La Malecita S/N, 38480 Buenavista del Norte, Tenerife, Canary Islands, SPAIN

Rodríguez A., Rodríguez B., Rumeu B., Nogales M. 2007. Seasonal diet of the Grey Heron *Ardea cinerea* on an oceanic island (Tenerife, Canary Islands): indirect interaction with wild seed plants. *Acta Ornithol.* 42: 77–87.

Abstract. In 199 pellets analyzed a total of 7 460 prey items were counted, 96.2% of which were arthropods. Aeshnidae larvae (Odonata) made up 66.1% of the total prey items and were the main invertebrate group. Vertebrates constituted 3.8%, with reptiles and mammals being the main prey of this type (1.8% each). Despite the small size of the invertebrates, this group reached > 60% in terms of biomass. All the main prey items varied significantly among seasons. Odonata was the most important group in all seasons, reaching its maximum value in summer. In the case of vertebrates, reptiles were captured mainly in spring, mammals in winter. With regard to indirect interaction with seeds, a total of 901 seeds associated with lizard remains were found in 77 pellets, indicating that they had previously been consumed by these reptiles. External visual damage of seeds was low and only 1.1% was destroyed. No seeds germinated after the four-month germination experiment and practically all of them were unviable. In conclusion, these results indicate that Grey Heron diet on islands varies in comparison with continental zones, including an important number of invertebrates and reptiles. Furthermore, this bird acts as an opportunistic secondary seed disperser, although its ecological effect does not seem to be very significant for the dynamics of the Canary Island ecosystems.

Key words: Grey Heron, *Ardea cinerea*, feeding ecology, seasonal diet pattern, secondary seed disperser, islands

Received — Nov. 2006, accepted — Febr. 2007

INTRODUCTION

The Grey Heron is mainly a winter visitor in the Macaronesian archipelagos (Snow & Perrins 1998), although recent breeding records have been mentioned in the Cape Verde Islands (Palacios & Barone 2001). In the Canaries, only some unsuccessful nesting attempts have been recorded (Martín & Lorenzo 2001). In these islands (e.g. Tenerife) some individuals are present throughout the year, not only using inland reservoirs as feeding sites, but also fields, water sources in ravines and xeric flat lands. Its winter population ranges between 29–62 individuals (Martín & Lorenzo 2001).

Grey Heron diet, during the breeding season, has been well documented in Europe by using several methods (Owen 1955, Hewson & Hancox

1979, Giles 1981, Moser 1986, Draulans et al. 1987, Marquiss & Leitch 1990, Peris et al. 1995, Lekuona & Campos 1998, Jakubas & Mioduszewska 2005). However, information on its seasonal feeding habits throughout the year remains scarce, probably because in the northern part of its geographical range the species shows a migratory behaviour. Furthermore, very little is known about the feeding ecology of this heron in insular environments, particularly in oceanic islands (Voisin 1991, Sawara et al. 1994, Nanpo & Matsuda 2006).

In general, the Grey Heron shows piscivorous feeding behaviour in continental ecosystems. Nevertheless, preliminary observations carried out in the Canaries have shown that most of the pellets have a high content in Odonata larvae and, less frequently, aquatic beetles (Martín & Lorenzo

2001). Furthermore, this heron preys upon lizards in the Canary archipelago. In this regard, the consumption of fleshy fruits and subsequent seed dispersal process by lizards is a relatively common finding in tropical and subtropical oceanic island ecosystems (see Olesen & Valido 2003 and references therein). This framework suggests that herons could be functioning as secondary seed dispersers of some native plants in the Canary Islands, as occurs with other bird species (Nogales et al. 1998, 2002). However, the ecological significance of this interaction depends greatly on the secondary disperser. Data contained in the two scientific contributions previously cited demonstrate how the Southern Grey Shrike *Lanius meridionalis* is a legitimate disperser for *Lycium intricatum* (Solanaceae) and how the Kestrel *Falco tinnunculus* is an illegitimate disperser of their seeds. One of the main factors that explain these results is the fact that gut enzymatic action is clearly stronger in the diurnal raptor. Furthermore, it is also interesting to note that this ecological interaction depends on the seed coat thickness of the plant species which is participating in the interaction. With regard to the Grey Heron and its digestive seed treatment, it is important to note that its gut enzymatic action is rather strong (Hibbert-Ware 1940, Vinokurov 1960), causing degradation in most of the prey items appearing in the pellets. Thus, it would be very interesting to perform experiments on the viability and ability to germination of the seeds found in heron pellets as a result of its predation on frugivorous lizards.

In the present study, pellets were used to determine the annual Grey Heron diet in one of the most important roosting sites in Tenerife Island (Martín & Lorenzo 2001). This method was selected because it is able to supply a representative sample size (see Methods). Three main aims of this study are: 1) to describe, in detail, the seasonal variation of the Grey Heron diet in an oceanic island environment; 2) to compare these insular trophic patterns with those of different continental populations; and 3) to assess the ecological interaction with wild seed plants from the Canary Islands.

STUDY AREA AND METHODS

The material was collected in the Valle Molina water reservoir (Tegueste), located in the north-east of Tenerife Island (Canarian Archipelago,

27°37'–29°25'N and 13°20'–29°25'W). This locality is situated at 340 m a.s.l. and is surrounded by different types of crops (mainly vines), grassland zones and villages. The climate is subtropical and oceanic, this area receiving around 400–500 mm of mean annual rainfall and having a mean annual temperature ranging between 16–19°C (Marzol 2000).

This reservoir has a maximum area of 7.5 ha and has a perimeter of 1034 m. Its bottom is covered by a plastic layer and its banks have not been colonized by plants. Grey Herons use this place as a roosting site and foraging area. Other heron species such as the Little Egret *Egretta garzetta* and the Cattle Egret *Bubulcus ibis* are regularly present at this place, in addition to many migratory species (ducks, waders, gulls, etc.) (see Martín & Lorenzo 2001). While the pellets were collected (see details in the following section), two-monthly censuses were carried out to assess a minimum size of the study population. A mean of 14.2 ± 7.3 (max and min values = 33 and 1 individuals, respectively) Grey Herons were counted. A mean sighting of 14.2 individuals throughout the year would represent a range of 22.9–48.9% with respect to the total population estimated in Tenerife by Martín & Lorenzo (2001).

Collection, quantification and methodological considerations on pellets

At the beginning of each month throughout 2004, all the margins of the reservoir were cleared, removing or destroying all pellets found, to ensure that only fresh pellets were collected. A minimum of twenty pellets were collected, when possible, at the middle and at the end of each month (two visits per month), but they were seasonally grouped. A total of 60 pellets were gathered in spring and summer, but only 48 and 31 in winter and autumn, respectively. Each pellet was wrapped in aluminium foil, stored in a plastic bag and dried at room temperature prior to analysis. All pellets were broken up by hand under dry conditions, after which different items were separated. Invertebrate remains were identified using a stereo-microscope (8× magnification) (Baéz 1985, Ralph et al. 1985). Some doubtful prey remains were identified by referring to the collection of the Zoology Department (University of La Laguna). The minimum number of individuals present in each pellet was estimated according to the most commonly found body parts, which represented one individual (heads, mandibles, legs, sclerotized parts, caudal appendages, etc.).

Invertebrates less than 1 cm long were not considered in this study because these items occurred in association with reptilian remains ($G_1 = 96.61$, $p < 0.001$). In the case of Oligocheates, due to their soft body, only frequency of occurrence was recorded. In the case of mammal remains, teeth, lower jaws and bones contributed to the identification of some prey items. Since the majority of Grey Heron pellets do not contain these parts (Hibbert-Ware 1940, Draulans et al. 1987, Marquiss & Leitch 1990), mammalian fur samples were identified by comparison with a previously prepared reference collection, and also by using the identification keys developed by Day (1966). From each pellet, ten hairs were sampled for microscopic analysis, following the same protocol used by Jakubas & Mioduszevska (2005). We assumed that a minimum of one mammalian specimen was eaten by a heron when the species occurred in a pellet, following the Amat & Herrera (1978) criteria for other species. In some instances, fish and reptilian remains were mainly identified on the basis of their respective scale traits.

Data on diet are presented in four ways: number of prey items (NP), prey item percentage (%P, the number of items of a particular prey taxon recorded in the diet as a percentage of the total number of prey items identified), percentage frequency of occurrence (%FO, the percentage of pellets containing a particular prey taxon), and percentage contribution to dietary biomass (%B). Wet biomass was calculated using an average weight for reptiles (personal data) and mean weights for mammals and birds were obtained from the literature (Castells & Mayo 1993, Cramp 1998). In the case of the main arthropods and fish, we collected several individuals (five at least) of each identified taxon and took their fresh weight using an electronic balance (precision: 0.01 g).

Several techniques (direct observations, analysis of stomach contents, food regurgitated by nestlings at nest, or undigested remains from regurgitated pellets) have been used to determine Grey Heron diet, but all of them present some degree of bias (see Marquiss & Leitch 1990). The absence of breeding colonies in the Canaries (Martín & Lorenzo 2001) limited us to using prey remnants or chick regurgitation in heronries. Only a small sample of stomach contents from dead birds could be obtained because their corpses are difficult to find. Furthermore, direct observations of herons at their feeding sites was greatly hindered by their erratic feeding habits (van Vessem et al. 1984, pers. obs.), and the

notable diversity of feeding sites identified in a small area in Tenerife. All the above, together with the presence of an accessible communal roosting site, where herons are present throughout the year, forced us to use pellet analysis. However, this method presents a degree of bias based on the differential digestion rate of each prey item (Milstein et al. 1970, Draulans et al. 1987, Marquiss & Leitch 1990). In this sense, fish and amphibians are almost absent in pellets, and mammal quantification is also difficult because of the almost complete lack of key bones (Milstein et al. 1970, Draulans et al. 1987, Marquiss & Leitch 1990).

Ingested seeds, germination and status

All seeds found inside pellets were preserved in a dry, dark place until they were identified using the reference collection of the IPNA-CSIC, at La Laguna. Most of them ($n = 753$) were planted separately in a 4 cm² pot, using a standard substrate (50% culture soil, 25% turf and 25% volcanic sand). To control environmental conditions, a germination experiment was performed in a greenhouse located at Tagoro (Tenerife: 300 m a.s.l.), a locality near (4 km) to the study area where pellets were collected. Therefore, night-day light period conditions were the same as those naturally found at the study site. Each seed was buried about 5 mm deep and watered every two days over four months (December 2005 – March 2006). Data were recorded every five days.

Due to the relatively small number of seeds found in the heron pellets, we decided to plant the majority of the samples. To assess the apparent status of embryos, a sub-sample of seeds ($n = 129$) of the main represented plants ($n = 7$ species) that did not germinate during these four months were extracted from the soil and then observed by using a stereo-microscope (8–50× magnification).

Statistical and index analysis

To study seasonal variation of the main items found in the diet, likelihood ratio tests were applied directly to the number of prey items which appeared in the pellets. These analyses compared a certain prey item with the total number of the remaining prey items identified. In those cases where it was necessary to use the same data set, this statistical effect was reduced by applying a more conservative sequential Bonferroni technique, for assessing statistical significance (Rice 1989). Due to the scarce presence of some prey items (Aranea, Tricoptera, Mantodea, Lepidoptera, Pisces, etc.), it was not feasible to

carry out any statistical inference. Diet similarity or overlap among the different seasons was evaluated using the Morisita index of similarity for percentage of prey, whose values near "0" indicate low similarity and values near "1", a high similarity. Moreover, niche-breadth was assessed using the standardized Levin's niche-breadth index, where values close to "0" indicate dietary specialization and values close to "1", a broad diet (Krebs 1989). To calculate the index values, we used the taxonomical level of order for invertebrates and species for vertebrates. Since it was impossible to assign numbers for Oligochaetes, this prey item was not included in the index analyses.

To test whether or not seeds of fleshy fruits in pellets occurred in association with lizard remains, their presence in each pellet was recorded individually and a likelihood ratio test was used to assess such an association. All analyses were performed by using SPSS (version 12.0).

RESULTS

General diet composition

A total of 7 460 prey items were identified in the 199 pellets analysed, 96.2% corresponding to arthropods and the remainder consisting mainly of vertebrates (Table 1). A high variety of insect families was recorded. Aeshnidae larvae (Odonata), representing 66.1% of the total prey items, was the main invertebrate group. This family was followed in number by Libellulidae larvae (Odonata, 23.8%), Acrididae (Orthoptera, 1.3%) and Carabidae (Coleoptera, 1.1%). With regard to the vertebrates (3.8% of the total prey items), the principal groups corresponded to reptiles and mammals (1.8%, respectively).

In terms of FO, invertebrates clearly predominated in the diet (94.0%); specifically, Aeshnidae was the most important family (59.8%). The next most frequent groups were Acrididae (18.6%), Libellulidae larvae (15.6%), Gryllidae (9.1%), and Carcinophoridae (Dermaptera) and Hymenoptera (7%, respectively). Furthermore, Oligochaetes occurred at 1.1% of pellets and vertebrates at 68.8% of FO. In this case, mammals (53.8%) and reptiles (48.4%) were the most relevant groups (Table 1).

As for biomass, despite the low size of invertebrates in comparison with vertebrates, the most important group was Odonata (61.4%), Aeshnidae (57.3%) being the main family represented (Fig. 1).

Introduced mammals (mice, rats, shrews and rabbits) constituted the second group in importance (27.4%) followed by reptiles (8.9%). Fishes, amphibians and birds were negligible prey items for the total biomass.

Seasonal variation in diet

All the main prey items varied significantly among seasons. Odonata was the most important group in all seasons although its consumption varied significantly, being highest during the summer ($G_3 = 54.19$, $p < 0.001$). The two main groups of this order were Aeshnidae and Libellulidae larvae. The former larvae showed statistical significance among seasons ($G_3 = 10.45$, $p = 0.015$), but their consumption was lower in summer. Seasonal variation was also observed in the case of Libellulidae ($G_3 = 844.24$, $p < 0.001$), but they were more frequently captured during summer. The maximum peak of grasshoppers was observed in autumn ($G_3 = 255.53$, $p < 0.001$). Acrididae were most frequently eaten in winter and spring, while the diet in autumn was characterised by the presence of Tettigonidae and Gryllidae. The consumption of Dermaptera (Carcinophoridae) reached maximum values in spring and winter ($G_3 = 102.20$, $p < 0.001$). With regard to vertebrates, reptiles appeared more often in spring ($G_3 = 156.43$, $p < 0.001$), mammals in winter ($G_3 = 129.54$, $p < 0.001$) and fish in autumn. 70.1% of total mammals corresponded to House Mice *Mus domesticus* which characterised the diet in autumn ($G_3 = 87.51$, $p < 0.001$). Other groups, such as fish, shrews, rabbits, and the




Fig. 1. Percentage of wet biomass for each group of prey items consumed by the Grey Heron on Tenerife (Canary Islands) in different seasons.

Table 1. Seasonal changes in diet composition in the Grey Heron (January 2004–December 2004) on Tenerife (Canary Islands). Numbers in brackets correspond to pellets analysed. NP — number of prey items, %P — percentage of prey, %FO — frequency of occurrence in pellets.

Prey items	Spring (N = 60)			Summer (N = 60)			Autumn (N = 31)			Winter (N = 48)			Total (N = 199)		
	NP	%P	%FO	NP	%P	%FO	NP	%P	%FO	NP	%P	%FO	NP	%P	%FO
Invertebrates	990	89.4	98.3	4654	99.2	100	556	90.4	87.1	975	93.3	85.4	7175	96.2	94.0
Oligochaeta	-	-	31.7	-	-	-	-	-	-	-	-	6.3	-	-	11.1
Araneae	3	0.3	3.3	0	0	0	3	0.5	6.5	4	0.4	6.3	10	0.1	3.5
Diplopoda	7	0.6	1.7	0	0	0	0	0	0	0	0	0	7	0.1	0.5
Odonata	767	69.3	58.3	4583	97.7	93.3	479	77.9	51.6	895	85.6	62.5	6724	90.1	68.8
Orthoptera	63	5.7	28.3	16	0.3	13.3	64	10.4	51.6	39	3.7	33.3	182	2.4	28.6
Dermoptera	24	2.2	21.7	0	0	0	0	0	0	17	1.6	2.1	41	0.5	7.0
Tricoptera	2	0.2	1.7	2	0	3.3	6	1.0	3.2	0	0	0	10	0.1	2.0
Mantodea	0	0	0	0	0	0	1	0.2	3.2	1	0.1	2.1	2	0	1.0
Hemiptera	7	0.6	8.3	32	0.7	3.3	0	0	0	6	0.6	4.2	45	0.6	4.5
Coleoptera	106	9.6	31.7	6	0.1	8.3	2	0.3	6.5	11	1.1	12.5	125	1.7	16.1
Lepidoptera	7	0.6	8.3	1	0	1.7	0	0	0	0	0	0	8	0.1	3.0
Hymenoptera	5	0.5	8.3	10	0.2	11.7	1	0.2	3.2	2	0.2	2.1	18	0.2	7.0
Hexapoda indet.	0	0	0	2	0	3.3	0	0	0	0	0	0	2	0	1.0
Vertebrata	117	10.6	76.7	39	0.8	41.7	59	9.6	80.7	70	6.7	72.9	285	3.8	68.8
Pisces	6	0.6	5.0	1	0	1.7	7	1.1	3.2	0	0	0	14	0.2	2.5
<i>Gambusia affinis</i>	5	0.5	3.3	1	0	1.7	7	1.1	3.2	0	0	0	13	0.2	2.0
Gen. sp. indet.	1	0.1	1.7	0	0	0	0	0	0	0	0	0	1	0	0.5
Amphibia	1	0.1	1.7	0	0	0	0	0	0	0	0	0	1	0	0.5
<i>Rana perezi</i>	1	0.1	1.7	0	0	0	0	0	0	0	0	0	1	0	0.5
Reptilia	66	6.0	70.0	18	0.4	25.0	26	4.2	58.1	25	2.4	43.8	135	1.8	48.4
<i>Gallotia galloti</i>	32	2.9	51.7	12	0.3	20	12	2.0	38.7	11	1.1	22.9	67	0.9	33.2
<i>Chalcides viridanus</i>	34	3.1	56.7	6	0.1	10	14	2.3	45.2	14	1.3	29.2	68	0.9	34.2
Aves	0	0	0	0	0	0	1	0.2	3.2	0	0	0	1	0	0.5
<i>Calidris</i> sp.	0	0	0	0	0	0	1	0.2	3.2	0	0	0	1	0	0.5
Mammalia	44	4.0	56.7	20	0.4	26.7	25	4.1	67.7	45	4.3	75.0	134	1.8	53.8
<i>Mus domesticus</i>	29	2.6	48.3	15	0.3	25.0	20	3.3	64.5	30	2.9	62.5	94	1.3	47.2
<i>Rattus</i> sp.	12	1.1	20.0	3	0.1	5.0	2	0.3	6.5	9	0.9	18.8	26	0.3	13.1
<i>Suncus etruscus</i>	3	0.3	5.0	1	0	1.7	3	0.5	9.7	5	0.5	10.4	12	0.2	6.0
<i>Oryctolagus cuniculus</i>	0	0	0	1	0	1.7	0	0	0	1	0.1	2.1	2	0	1.0
Total of prey	1107			4693			615			1045			7460		

invertebrates (Lepidoptera or Hymenoptera) appeared in small numbers (Table 1). Oligochaetes occurred in two seasons, spring (31.7% of FO) and winter (6.3%).

In terms of biomass, Odonata were the most important dietary component in summer (85.1%), when mammals and reptiles decreased in numbers. In spring and winter, the most relevant group was mammals (38.3% and 47.9%, respectively). Reptiles attained their maximum value in spring (18.7%) (Fig. 1).

Levin's niche breadth (B) was broader in spring (0.063) and narrower in summer (0.003). In the other two seasons this index had medium values (autumn = 0.047 and winter = 0.025). The Morisita index indicates a high trophic overlap throughout practically all seasons, spring and summer showing the clearest differences in values with respect to the other two seasons (Table 2).

Indirect seed interaction

A total of 901 seeds were counted in all analysed pellets. Seeds from at least 12 taxa were identified at species level, representing 90.5%. After extraction from pellets, external visual damage of seeds ingested by the Grey Heron was low (1.1%) (Table 3). No seeds germinated after the four month experiment. Data on embryo viability indicate that only 4.7% seeds could still be alive after the germination experiment, indicating that the Grey Heron digestive gut effect caused significant damage to seed fate (Table 4). Seeds present in pellets were significantly associated with the presence of lizard remains ($G_1 = 75.96$, $p < 0.001$), which indicates that these seeds were previously consumed by these reptiles and indirectly by the herons.

DISCUSSION

Composition diet and seasonal variation

This study provides the first detailed description of the seasonal variation through a year of the

Grey Heron diet on an oceanic island. However, the majority of contributions in continental ecosystems were made during the breeding period (Owen 1955, 1960, Hewson & Hancox 1979, Giles 1981, Moser 1986, Marquiss & Leitch 1990, Peris et al. 1994, 1995, Lekuona & Campos 1998, Lekuona 1999, 2001, Jakubas & Mioduszewska 2005).

Differences in diet composition between continental and insular areas were found (Table 5). It has been suggested that the diet varies considerably according to different habitats and seasons, and the studied species exhibits opportunistic feeding behaviour (Voisin 1991, Fasola et al. 1993, Cramp 1998). In the Canaries, Grey Herons often feed at agricultural ponds, but the physical characteristics of the majority of these reservoirs, which are usually deep with steep high banks, hinder fish capture (Draulans 1987, Voisin 1991). On marine shores, other species of fishes are available, but unlike more northern latitudes (Lekuona 1999, Carss & Elston 2003), the Grey Heron practically does not use this habitat as a foraging site in Tenerife (pers. obs.). Therefore, the low number of fish recorded in the present study may be due to limited access. On the other hand, it was found that the bulk of the diet in Tenerife was composed of Odonata larvae, which are present at both of these sites and in addition to all manner of natural bodies of water (Malmqvist et al. 1995), and may be more accessible than fish to herons because they remain relatively still near the surface on the pond walls until ready to leave the water. This prey item has been poorly represented in several studies carried out in Europe (Hibbert-Ware 1940, Draulans et al. 1987, Campos 1990, Jakubas & Mioduszewska 2005). Reptiles are rarely consumed in continental areas (Hibbert-Ware 1940, Moser 1986, Campos 1990, Peris et al. 1994, 1995, Cramp 1998, Jakubas & Mioduszewska 2005) but in Tenerife they constitute almost 50% of FO. This phenomenon may be related to the high density of lizards in insular environments (Olesen & Valido 2003, and references therein). In other predatory vertebrates studied in the Canaries, the same pattern was recorded (Carrillo et al. 1994, Nogales & Medina 1996, Padilla et al. 2005). Furthermore, contrary to other studies carried out in continental environments (Cuesta et al. 1980, Campos 1990, Peris et al. 1995, Jakubas & Mioduszewska 2005), invertebrates constituted more than 50% of ingested biomass. However, this result is probably a consequence of the

Table 2. Morisita's niche overlap index for Grey Heron diet between seasons in Tenerife (Canary Islands); results are based on percentage of prey for each food item.

Seasons	Summer	Autumn	Winter
Spring	0.934	0.983	0.972
Summer	-	0.967	0.990
Autumn	-	-	0.992
Winter	-	-	-

Table 3. Seeds found in Grey Heron pellets over a year (January–December 2004) on Tenerife (Canary Islands). N — number of pellets, NS — number of seed, %S — percentage of seeds, %FO — frequency of seed occurrence in pellets, %DS — percentage of seeds visually damaged. Nomenclature of plant species follows the check-list of Izquierdo et al. (2004).

Plant species	Family	Spring (N = 60)		Summer (N = 60)		Autumn (N = 31)		Winter (N = 48)		Total (N = 199)	
		NS	%S	NS	%S	NS	%S	NS	%S	NS	%S
<i>Solanum cf. nigrum</i>	Solanaceae	216	40.3	38	53.5	3	1.4	2	2.4	259	28.7
<i>Atriplex semibaccata</i>	Chenopodiaceae	97	18.1	1	1.4	27	12.8	37	44.0	162	18.0
<i>Rubus ulmifolius</i>	Rosaceae	1	0.2	1	1.4	135	64.3	2	2.4	139	15.4
<i>Lycopersicon esculentum</i>	Solanaceae	111	20.7	8	11.3	-	-	-	-	119	13.2
<i>Rubia fruticosa</i>	Rubiaceae	47	8.8	-	-	-	-	1	1.2	48	5.3
<i>Opuntia maxima</i>	Cactaceae	-	-	8	11.3	34	16.2	-	-	42	4.7
<i>Canarina canariensis</i>	Campanulaceae	13	2.4	-	-	-	-	-	-	13	1.5
<i>Lantana camara</i>	Verbenaceae	-	-	-	-	-	-	13	15.5	13	1.5
<i>Jasminum odoratissimum</i>	Oleaceae	9	1.7	-	-	-	-	-	-	9	1.0
<i>Telina sp.</i>	Fabaceae	6	1.1	-	-	-	-	3	3.6	9	1.0
<i>Chenopodium sp.</i>	Chenopodiaceae	3	0.6	1	1.4	5	2.4	-	-	9	1.0
<i>Vitis vinifera</i>	Vitaceae	-	-	7	9.9	1	0.5	-	-	8	0.9
<i>Avena sp.</i>	Poaceae	2	0.4	1	1.4	-	-	-	-	3	0.3
<i>Malva parviflora</i>	Malvaceae	2	0.4	-	-	-	-	-	-	2	0.2
<i>Achyranthes aspera</i>	Amaranthaceae	1	0.2	-	-	-	-	-	-	1	0.1
Unidentified seeds	-	28	5.3	6	8.4	5	2.4	26	30.9	65	7.2
Total		536		71		210		84		901	
											38.7
											1.1

impossibility to quantify mammals (only one individual was assumed to be present in each pellet). All these findings seem to indicate that differences in Grey Heron diet (mainland vs. islands) are based on distinct availability of potential prey items in foraging areas.

In our study area, seasonal variation of diet was high and all main prey items identified varied significantly. The consumption of vertebrates (reptiles and mammals) clearly decreases in summer, when lizards are more abundant and most active. However, this pattern was observed in mammalian prey in other studies performed during the breeding season (Campos 1990). In contrast, peak consumption of Odonata larvae occurs in summer. Dragonfly larvae are present throughout the year in the Canaries, but their demographic explosion takes place in spring and summer (Báez 1985), which would explain vertebrate prey substitution by Odonata larvae. This change in predatory behaviour is probably based on the greater availability of these prey.

The highest value obtained from Levin's niche breadth in spring is a result of an increase of insect abundance after autumn and winter rains. Due to the high abundance of Odonata larvae (Báez 1985) and the fact that herons concentrate their predation on certain prey components (Müller 1984), the niche breadth index decreases to a minimum during the dry season. Considering the mild climate throughout the year in this archipelago (Marzol 2000), and the low diversity of prey on islands, the important trophic niche overlap explains the high diet similarity recorded among seasons.

Seed interaction

Six of 12 plant species identified in seed analyses (*Solanum cf. nigrum*, *Rubus ulmifolius*, *Rubia fruticosa*, *Canarina canariensis*, *Jasminum odoratissimum* and *Malva parviflora*) were native and the remainder corresponded to species introduced by man (see Izquierdo et al. 2004). One species from the former group (*R. fruticosa*) is endemic to the Macaronesian Islands and another (*C. canariensis*) is endemic to the Canarian Archipelago. However, these species together constituted only 6.8% of the total seeds indirectly ingested by the Grey Heron. Furthermore, 57.7% of the total identified seeds corresponded to native species, while a total of 42.3% were from introduced plants.

No germination and visual embryo analysis indicate that the digestive treatments, particularly the strong gut enzymatic effect on seeds in herons

Table 4. Seed status and germination of the different species indirectly consumed by the Grey Heron on Tenerife (Canary Islands).

Plant species	Number of seeds planted	% germinated seeds	Number of seeds extracted after germination	% apparent viable seeds
<i>Solanum cf. nigrum</i>	230	0	-	-
<i>Rubus ulmifolius</i>	135	0	24	12.5
<i>Atriplex semibaccata</i>	130	0	14	0
<i>Lycopersicon esculentum</i>	95	0	19	0
<i>Rubia fruticosa</i>	47	0	21	0
<i>Opuntia maxima</i>	38	0	30	10
Gen. sp. indet.	19	0	-	-
<i>Canarina canariensis</i>	13	0	-	-
<i>Lantana camara</i>	13	0	13	0
<i>Jasminum odoratissimum</i>	9	0	8	0
Unidentified seeds	24	0	-	-
Total	753	0	129	4.7

(Hibbert-Ware 1940, Vinokurov 1960) produce a clear negative effect on all plant species. In contrast seeds from the majority of these species, passed through a lizard gut, presented a considerable germination rate (Nogales et al. 2005, Authors' unpublished data). This negative effect on *L. intricatum* seeds, when the secondary disperser shows a strong enzymatic action, has been observed in other birds, such as some diurnal raptors (Nogales et al. 2002). However, when the second seed interaction is carried out by non-specialised predatory birds (e.g. the Southern Grey Shrike) the fate of the seeds is clearly different in terms of viability and germination (Nogales et al. 1998). Furthermore, another negative ecological effect caused by herons, in the majority of seeds, is that they concentrate great numbers of seeds in their roosting sites. These places, mainly located on the margins of reservoirs and coastal cliffs, are not suitable micro sites for seed germination and seedling establishment.

Comparing seed number and frequency of occurrence per plant species in Grey Heron pellets with those of other birds (e.g. kestrels and shrikes), the incidence was found to be higher in the latter because their predation on lizards is more frequent (see Nogales et al. 1998, 2002) and thus the interaction with wild seeds is more probable. Lastly, with the increasing knowledge of trophic interactions in subtropical islands, such as the Canary Archipelago, it can be seen that the participation of predatory birds (e.g. herons, diurnal and nocturnal raptors, shrikes, etc.), as legitimate/illegitimate secondary

seed dispersers, is a more common phenomenon than it has been thought to be. The intervention of Grey Heron as an illegitimate secondary seed disperser is a rather novel ecological interaction for this species. However, considering the relatively low number of seeds that occurred in pellets in comparison with the millions of seeds annually dispersed by abundant endemic lizards and passerines, the role of herons does not seem to be of great significance in the dynamics of Canarian ecosystems.

ACKNOWLEDGEMENTS

Many friends helped us during the field work, but we would especially like to thank Tinguaro Montelongo, Heriberto López, David P. Padilla, Pedro Oromí and Juan Carlos Rando identified some prey items. Concepción Nieves assisted during the checking of germination experiments and Christian Kowalewski translated some articles from the German. Dariusz Jakubas, Yuji Sawara, Salvador J. Peris and an anonymous reviewer made useful comments on an earlier draft of this paper. Finally, Bernardino Pérez, Juan Pedro Castro and Miguel Fernández del Castillo (Grupo de Ardeidas-SEO/Teydea) permitted us to use inedited information on the status of the Grey Heron in Tenerife. A. Rodríguez and B. Rumeu were partially supported by a postgraduate and a student grant (respectively) from the Spanish National Research Council (CSIC).

Table 5. Differences in Grey Heron diet composition, according to different sites and methods. RF — regurgitated food, GC — gut contents, DO — direct observations, PL — pellets, * — values lesser than 0.1, † — values referred to percentage of prey, § — values referred to frequency of occurrence.

Method	Locality	Main prey groups							Source
		Invertebrates	Fish	Amphibians	Reptiles	Birds	Mammals		
RF	England	26.9	70.9	0.2	0	0	0.1	1.9	Owen 1955 †
	England	26.4	71.4	0.1	0	0	0.1	2	Owen 1960 †
	Southeast France	17.6	81.9	0.3	0.1	0	0	0.1	Moser 1986 †
	Northern Italy	10.7	50.0	4.8	7.1	0	0	1.2	Fasola et al. 1993 †
	Japan	15.9	76.4	7.3	0	0	0	0.4	Sawara et al. 1994 †
	Northern Poland	?	95	1.2	0	0	0.9	Jakubas & Mioduszewska 2005 †	
GC	Northern Italy	68.3	26	21	24	0	0	12.5	Moltoni 1936 in Cramp 1998 §
	Spain	44.4	55.5	0	0	0	0	11.1	Bernis & Valverde 1954 §
	Hungary	25.1	40.1	20.3	0	0	0	14.5	Vasvari 1954 in Cramp 1998 †
	Germany	44	53	1	0	0	0	2	Schlegel 1964 †
	Central and Western Spain	53.7	40.8	1.1	1.1	?	?	2.5	Cuesta et al. 1980 †
	Switzerland	78	73	0	0	0	0	45	Müller 1984 §
	Northeast Spain	44.4	55.6	0	0	0	0	0	González-Martín & González-Solis 1990 †
DO	Southwest France	17.5	82.1	0.4	0	0	0	0	Lekuona 1999 †
	Northern Spain	21.0	73.1	2.8	1.2	0	0	1.8	Lekuona 2001 †
RF & PL	Scotland	12.1	35.4	12.9	0	0	0	39.6	Giles 1981 †
	Scotland	1.6	69.5	0.8	0	0	21.1	7.0	Marquiss & Leitch 1990 †
RF, GC & PL	Central-West Spain	11.9	81.6	0.6	0.8	0.8	0.8	4.2	Campos 1990 †
	Central-West Spain	14.1–20.0	62.9	1.8	0.6	1.5	3.2	3.2	Peris et al. 1995 §
PL	England	?	0	0	0	0	30.0	75.0	Milstein et al. 1970 §
	Northeast Scotland	38.9	0	0	0	0	29.6	98.1	Hewson & Hancox 1979 §
	Northern Poland	26–51	1–5	0.2–0.3	0	0.5–0.9	99.9	99.9	Jakubas & Mioduszewska 2005 §
	Tenerife, Canary Islands	96.2	0.2	*	1.8	*	1.8	1.8	Present study †

REFERENCES

- Amat J. A., Herrera C. M. 1978. [Diet of the Purple Heron (*Ardea purpurea*) in the marismas of the Guadalquivir (Doñana National Park) during the breeding period]. *Ardeola* 24: 95–104.
- Baéz M. 1985. [The Dragonflies of the Canary Islands]. Aula de la Cultura Popular Canaria, Santa Cruz de Tenerife.
- Bernis F., Valverde J. A. 1954. [On the Grey Heron (*Ardea cinerea*) in Spain]. *Bol. R. Sdad. Esp. Hist. Nat.* 50: 201–213.
- Campos F. 1990. [Diet of the Grey Heron (*Ardea cinerea*) in the Duero basin (Spain) during the breeding period]. *Doñana Acta Vert.* 17: 141–151.
- Carrillo J., Hernández E. C., Nogales M., Delgado G., García R., Ramos T. 1994. Geographic variation in the spring diet of *Falco tinnunculus* L. on the islands of Fuerteventura and El Hierro (Canary Islands). *Bonn. zool. Beitr.* 45: 39–48.
- Carss D. N., Elston D. A. 2003. Patterns of association between algae, fishes and grey herons *Ardea cinerea* in the rocky littoral zone of a Scottish sea loch. *Estuar. Coast. Shelf S.* 58: 265–277.
- Castells A., Mayo M. 1993. [Guide of the Mammals at liberty of Spain and Portugal]. Ediciones Pirámide, Madrid.
- Cramp S. 1998. The Complete Birds of the Western Palearctic on CD-ROM. Oxford University Press.
- Cuesta L., Lobón-Cerviá J., González L. M., González J., Llandres C., Garzón J., Palacios F. 1980. [Diet of the Grey Heron (*Ardea cinerea cinerea* L., 1758) in Spain]. II Reunión Iberoamer. Cons. Zool. Vert.: 189–195.
- Draulans D. 1987. The effectiveness of attempts to reduce predation by fish-eating birds: a review. *Biol. Conserv.* 41: 219–232.
- Draulans D., Perremans K., van Vessem J., Pollet M. 1987. Analysis of pellets of the grey heron, *Ardea cinerea*, from colonies in Belgium. *J. Zool.* 211: 695–708.
- Day M. G. 1966. Identification of hair and feather remains in the gut and faeces of stoats and weasels. *J. Zool. Lond.* 148: 201–217.
- Fasola M., Rosa P., Canova L. 1993. The diets of squacco herons, little egrets, night, purple and grey herons in their Italian breeding ranges. *Rev. Ecol. (Terre Vie)* 48: 35–47.
- Giles N. 1981. Summer diet of the grey heron. *Scot. Birds* 11: 153–159.
- González-Martín M., González-Solís J. 1990. [Feeding data of Herons from the Ebro Delta]. *Misc. Zool.* 14: 240–244.
- Hibbert-Ware A. 1940. An investigation of the pellets of the Common Heron (*Ardea cinerea cinerea*). *Ibis* 82: 433–450.
- Hewson R., Hancox M. 1979. Prey remains in grey heron pellets from north-east Scotland. *Bird Study* 26: 29–32.
- Izquierdo I., Martín J. L., Zurita N., Arechavaleta M. (eds). 2004. [Check-list of wild species of the Canary Islands (fungi, plants and terrestrial animals) 2004]. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias, Santa Cruz de Tenerife.
- Jakubas D., Mioduszewska A. 2005. Diet composition and food consumption of the grey heron (*Ardea cinerea*) from breeding colonies in northern Poland. *Eur. J. Wildl. Res.* 51: 191–198.
- Krebs C. J. 1989. *Ecological Methodology*. Harper Collins, New York.
- Lekuona J. M. 1999. Food and foraging activity of grey herons, *Ardea cinerea*, in a coastal area during the breeding season. *Folia Zool.* 48: 123–130.
- Lekuona J. M. 2001. Variations in food composition and foraging behaviour of Grey Heron *Ardea cinerea* in Aragon River (Northern Spain) during the breeding season. *Anu. Ornit. Navarra* 1999, 6: 19–25, Gorosti, Pamplona.
- Lekuona J. M., Campos F. 1998. [Food composition and foraging behaviour of Grey Heron (*Ardea cinerea*) and Purple Heron (*Ardea purpurea*) in the Aragón river during the breeding season]. *Anu. Ornit. Navarra* 1996, 3: 21–28, Gorosti, Pamplona.
- Malmqvist B., Nilsson A.N., Báez M. 1995. Tenerife's freshwater macroinvertebrates: status and threats (Canary Islands, Spain). *Aquat. Conserv.* 5: 1–24.
- Marquiss M., Leitch A. F. 1990. The diet of Grey Herons *Ardea cinerea* breeding at Loch Leven, Scotland, and the importance of their predation on ducklings. *Ibis* 132: 535–549.
- Martín A., Lorenzo J. A. 2001. [Birds of the Canarian Archipelago]. Lemus, La Laguna.
- Marzol V. 2000. [The climate]. In: Morales G., Pérez R. (eds). [Great Thematic Atlas of Canaries]. Editorial Interinsular Canaria, Santa Cruz de Tenerife, pp. 87–106.
- Milstein P. le S., Prestt I., Bell A. A. 1970. The breeding cycle of the Grey Heron. *Ardea* 58: 171–257.
- Moser M. E. 1986. Prey profitability for adult Grey Herons *Ardea cinerea* and the constraints on prey size when feeding young nestlings. *Ibis* 128: 392–405.
- Müller R. 1984. Magenuntersuchungen an Graureihern *Ardea cinerea*. *Orn. Beob.* 81: 159–163.
- Nanpo A., Matsuda K. 2006. [Seasonal changes in foraging sites and diets of the Grey Heron, in riverine environment]. *Strix* 24: 9–22.
- Nogales M., Delgado J. D., Medina F. M. 1998. Shrikes, lizards and *Lycium intricatum* (Solanaceae) fruits: a case of indirect seed dispersal on an oceanic island (Aleganza, Canaries). *J. Ecol.* 86: 866–871.
- Nogales M., Medina F. M. 1996. A review of the diet of feral domestic cats (*Felis silvestris* f. *catus*) on the Canary Islands, with new data from the laurel forest of La Gomera. *Z. Säugetierkd.* 61: 1–6.
- Nogales M., Nieves C., Illera J. C., Padilla D. P., Traveset A. 2005. Effect of native and alien vertebrate frugivores on seed viability and germination patterns of *Rubia fruticosa* (Rubiaceae) in the eastern Canary Islands. *Funct. Ecol.* 19: 429–436.
- Nogales M., Quilis V., Medina F. M., Mora J. L., Trigo L. S. 2002. Are predatory birds effective secondary seed dispersers? *Biol. J. Linn. Soc.* 75: 345–352.
- Olesen J. M., Valido A. 2003. Lizards as pollinators and seeds dispersers: an island phenomenon. *Trends Ecol. Evol.* 18: 177–181.
- Owen D. F. 1955. The food of the Heron *Ardea cinerea* in the breeding season. *Ibis* 97: 276–295.
- Owen D. F. 1960. The nesting success of the Heron *Ardea cinerea* in relation to the availability of food. *Proc. Zool. Soc. Lond.* 133: 597–617.
- Padilla D. P., Nogales M., Pérez A. J. 2005. Seasonal diet of an insular endemic population of Southern Grey Shrike *Lanius meridionalis koenigi* on Tenerife, Canary Islands. *Ornis Fennica* 82: 155–165.
- Palacios C. J., Barone R. 2001. Le Héron cendré *Ardea cinerea*, nouvelle espèce nidificatrice aux îles de Cap Vert. *Alauda* 69: 18.
- Peris S. J., Briz F. J., Campos F. 1994. Recent changes in the food of the Grey Heron *Ardea cinerea* in central-west Spain. *Ibis* 136: 488–489.
- Peris S. J., Briz F. J., Campos F. 1995. Shifts in the diet of the Grey Heron (*Ardea cinerea*) in the Duero Basin, Central-West Spain, following the introduction of exotic fish species. *Folia Zool.* 44: 97–102.
- Ralph C. P., Nagata S. E., Ralph C. J. 1985. Analysis of droppings to describe diets of small birds. *J. Field Ornithol.* 56: 165–174.

- Rice W. R. 1989. Analyzing tables of statistical tests. *Evolution* 43: 223–225.
- Sawara Y., Sakuyama M., Demachi G. 1994. [Diets and foraging site utilization of the Grey Heron, *Ardea cinerea*, in the breeding season]. *Jap. J. Ornithol.* 43: 61–71.
- Schlegel R. 1964. Zur Nahrung des Graureihers (*Ardea cinerea* L.) an Oberlausitzer Karpfenteichen. *Zool. Abh. Mus. Tierk. Dresden* 27: 65–67.
- Snow D. W., Perrins C. M. 1998. *The Birds of the Western Palearctic. Concise Edition. Vol. I.* Oxford Univ. Press.
- van Vessem J., Draulans D., de Bont A. F. 1984. Movements of radio-tagged Grey Herons *Ardea cinerea* during the breeding season in a large pond area. *Ibis* 126: 576–587.
- Vinokurov A. A. 1960. [On the food digestion role in heron *Ardea purpurea*]. *Bull. Otdel. Biol. Moskva* 65: 10.
- Voisin C. 1991. *The Herons of Europe.* T. & A. D. Poyser, London.

STRESZCZENIE

[Pokarm czapli siwej na Teneryfie (Wyspy Kanaryjskie) w cyklu rocznym — pośrednie związki z rozsiewaniem nasion].

Na Teneryfie nie istnieje lęgowa populacja czapli siwej, choć osobniki tego gatunku obserwowane są na tej wyspie przez cały rok. Pokarm ptaków określano analizując wypluwki zbierane na brzegach dużego zbiornika wodnego wykorzystywanego przez ptaki jako miejsce żerowania i noclegu. Łącznie zebrano 199 wypluwek, w których stwierdzono 7 460 okazów zdobyczy, wśród

których 96,2% stanowiły stawonogi. Wśród bezkręgowców dominowały larwy ważek z rodziny żagnicowanych (Aeshnidae), które stanowiły do 66.1% wszystkich zdobyczy. Kręgowce, głównie ssaki i gady stanowiły tylko 3.8% zdobyczy (Tab.1). Zaobserwowano dość znaczne zmiany sezonowe w składzie pokarmu (Fig. 1, Tab. 2). Wazki były najważniejszym pokarmem przez cały rok, a ich największy udział w diecie zanotowano latem. Wśród kręgowców ssaki łapano były najczęściej zimą, zaś gady — wiosną.

Łącznie w wyplawkach znaleziono 901 nasion 12 taksonów (Tab. 3). Obecność nasion związana była z pozostałościami po skonsumowanych jaszczurkach, co wskazywałoby, że zostały one wcześniej zjedzone przez te gady. Zewnętrznie nasiona te nie były prawie uszkodzone. Jednakże żadne z nich nie siewkowały w okresie czterech miesięcy po zasadzeniu, co wskazuje, że straciły żywotność po przejściu przez system trawienny czapli (Tab. 4). Wydaje się więc, że w związku z żerowaniem na jaszczurkach ptaki biorą udział w rozsiewaniu nasion, choć ich rola w tym procesie nie jest istotna w ekosystemie wysp Kanaryjskich.

W pracy zestawiono także dane o pokarmie czapli siwej zebrane w wielu miejscach i różnymi metodami (Tab. 5).