



Encyclia altissima (Orchidaceae) new for Cuba

Authors: Espinosa, María Del Carmen Fagilde, and Quesada, Eddy Martínez

Source: Willdenowia, 32(2) : 319-321

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.32.32213>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MARÍA DEL CARMEN FAGILDE ESPINOSA & EDDY MARTÍNEZ QUESADA

Encyclia altissima* (Orchidaceae) new for Cuba*Abstract**

Fagilde Espinosa, M. del C. & Martínez Quesada, E.: *Encyclia altissima* (Orchidaceae) new for Cuba. – Willdenowia 32: 319-321. 2002. – ISSN 0511-9618.

Encyclia altissima, so far known from the Bahamas, the Turks and Caicos Islands and Haiti, is recorded for Cuba from the Sierra de Mariana, Guantánamo province. For the Cuban population, a description and illustration, distribution data and conservation status are given.

During an expedition along the south coast of the province of Guantánamo a non-flowering orchid species was found by the authors, which could not then be identified. Pseudobulbs were taken into cultivation and brought to flower (Fig. 1). In flower, the plants could be identified with the help of Correll & Correll (1996) and Withner (1966, 1996) as *Encyclia altissima* (Bateman ex Lindl.) Schltr. [Syn.: *E. hodgiana* (Hawkes) Beckner], a species hitherto known only from the Bahamas, the Turks and Caicos Islands and Haiti (Correll & Correll 1996, Withner 1996).

Description of the Cuban plants. – *Perennial*, flowering up to 1.5 m tall, roots white, 3 mm in diameter, pseudobulbs up to 33 cm long and 3 cm thick, with 10-15 cm long imbricating sheaths. *Leaves* very coriaceous, linear to linear-lanceolate, 63-66 × 2.5 cm. *Inflorescence* terminal, 1.5 m long, with up to 30 flowers. *Sepals* 22 × 5 mm, elliptic, acute, greenish tan with reddish brown striping towards apex. *Petals* oblanceolate, somewhat narrower, otherwise similar. *Labellum* deeply 3-lobed, 22 × 20 mm wide; lateral lobes oblong, obtuse, erect, embracing the column, yellow with radiating purple lines, midlobe white with yellow undulating margin and radiating purple lines; two lateral erect callose keels joining at the midlobe, two undulate callose lamellae parallel the keels on the midlobe. *Column* elongate, white, streaked with purple, with membranaceous, incurved, rounded auricles 12 × 8 mm; *anther* 3 × 2 mm, yellow. Flowering: November to March.

Correll & Correll (1996) give the height of the plants with up to 210 cm, and the leaf length (but apparently due to a typographical error) with 6 cm; otherwise the Cuban plants match perfectly the description of plants from the other areas.




Fig. 1. *Encyclia altissima* – A: pseudobulb; B: leaf; C: flower. – Scale bar: A-B = 3 cm, C = 1 cm; drawn by Walfrido Lago based on Martínez & Fagilde 20631 from BSC.

Distribution in Cuba. – *Encyclia altissima* is known so far only from the Guantánamo province, where it is found in the Sierra de Mariana, c. 5.5 km from Baitiquirí, in a microphyllous, semideciduous forest, near a gypsum mine, c. 400 m above sea level. – Specimen: *Martínez & Fagilde 20631* (BSC).

Conservation status. – Because of the localized distribution in an area close to settlements and strongly affected by human activities, we suggest for *Encyclia altissima* in Cuba the status Critically Endangered (CR), according to the criteria A2c and C2b of the Conservation Assessment and Management Plan (CAMP) I for Cuban forest plants (Peña & al. 1998).

References

- Correll, D. S. & Correll, H. B. 1996: Flora of the Bahama archipelago (including The Turks and Caicos Islands). – Vaduz [reprint of orig. ed. Vaduz 1982].
- Peña, E., Lopez, P. I., Lazcano, J., Leiva, A. T. & Seal, U. S. 1998: Memorias del primer taller para la conservacion, analisis y manejo planificado de plantas silvestres cubanas, 13-15 abril. IUCN/SSC Conservation Breeding Specialist Group. – Apple Valley, MN.
- Withner, C. L. 1966: The differences among *Epidendrum altissimum*, *Epidendrum gracile*, and *Epidendrum rufum (bahamense)*. – *Orchidata* **6**: 128-129.
- 1996: The cattleyas and their relatives 4. The Bahamian and Caribbean species. – Portland.

Acknowledgements

We wish to thank Dr Robert Dressler from the Florida Museum of Natural History for helping us to identify the species, and Dr Lulú Rico and John Flanagan from the Royal Botanic Gardens, Kew, for providing us relevant photocopies.

Address of the authors:

María del Carmen Fagilde Espinosa & Eddy Martínez Quesada, Centro Oriental de Ecosistemas y Biodiversidad (BIOECO), Enramadas No. 601 esq. Barnada, Santiago de Cuba 1. CP 90100, Cuba; e-mail: eddy@bioeco.ciges.inf.cu.