

## **Two new combinations in *Morella* (Myricaceae) for species of the Cuban flora**

Authors: Iturralde, Rosalina Berazaín, and Hidalgo, Banessa Falcón

Source: *Willdenowia*, 41(1) : 113-114

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.41.41113>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ROSALINA BERAZAÍN ITURRALDE<sup>1\*</sup> & BANESSA FALCÓN HIDALGO<sup>1</sup>

## Two new combinations in *Morella* (*Myricaceae*) for species of the Cuban flora

### Abstract

Berazaín Iturralde R. & Falcón Hidalgo B.: Two new combinations in *Morella* (*Myricaceae*) for species of the Cuban flora. – Willdenowia 41: 113–114. – Online ISSN 1868-6397; © 2011 BGBM Berlin-Dahlem. doi:10.3372/wi.41.41113 (available via <http://dx.doi.org/>)

All Cuban representatives of *Myricaceae*, formerly treated under *Myrica* sensu lato, pertain to the segregate genus *Morella*. For two of the four Cuban species, *Myrica cacuminis* and *M. shaferi*, the required combinations do not exist as yet and are validly published here; a lectotype is designated for the latter name.

Additional key words: *Myrica*, taxonomy, Greater Antilles

*Myricaceae* is a small family, which has traditionally been considered to be monogeneric. Macdonald (1989) has summarised the reasons for dismembering the Linnean genus *Myrica* and his arguments have gained increased acceptance in the following years. Recent studies (Wilbur 2001; Herbert 2005; Mabberley 2008) treat the family with four genera: *Myrica* L. (temperate North America and Europe), *Morella* Lour. (pantropical), *Comptonia* L'Hér. (northeastern North America) and *Canaomyrica* Guillaumin (New Caledonia).

Several former species of *Myrica* from North America (Wilbur 1994), Africa (Killick & al. 1999), Malesia (Turner 2001), South America (Parra-O. 2002) and some from Central America and the Caribbean (Wilbur 2001; Knapp 2002) have already found their place in *Morella*. As characterised by e.g. Herbert (2005), *Morella* comprises evergreen shrubs or trees with inflorescences borne on the present year's growth and fleshy, papillose fruits; whereas *Myrica* consists of deciduous small shrubs with inflorescences borne on the previous year's growth and dry fruits with adherent spongy bracteoles.

For the Cuban flora, León & Alain (1951) reported four species of *Myricaceae*, which they placed in *Myrica*, but which under the above criteria clearly belong in

*Morella*. Two of them have already been transferred to that genus, as *M. cerifera* (L.) Small and *M. punctata* (Griseb.) J. Herb. For the two others, the required combinations do not as yet exist. They are here proposed, as follows.

*Morella cacuminis* (Britton & P. Wilson) Berazaín & Falcón, **comb. nov.** ≡ *Myrica cacuminis* Britton & P. Wilson in Bull. Torrey Bot. Club 50: 35. 1923. – Holotype: Cuba, Pico Turquino, July 1922, León LS 10973 (NY 73365 [photo!]; isotypes: HAC (3×)!, US 26705 [photo!]).

*Morella cacuminis* can be recognised by its small elliptic, rounded to obtuse, bullate leaves with obtuse base and irregularly toothed margin, the venation being strongly impressed adaxially and prominent abaxially. It is endemic to the high mountain peaks of SE Cuba (Sierra Maestra).

*Morella shaferi* (Urb. & Britton) Berazaín & Falcón, **comb. nov.** ≡ *Myrica shaferi* Urb. & Britton in Urban, Symb. Antill. 7: 190. 1912. – Holotype: Cuba, Baracoa to Florida, Oriente, serpentine hills, 15.3.1910, *Shafer 4331*

<sup>1</sup> Jardín Botánico Nacional, Universidad de La Habana, Carretera del Rocío km 3½, Calabazar, C.P. 19230, La Habana, Cuba;

\*e-mail: malvarosa@fbio.uh.cu (author for correspondence).

(B †). Lectotype (designated here): (NY 73366 [photo!]; isolectotype: HAC!).

*Morella shaferi* can be recognised by its obovate, truncate to apiculate leaves with cuneate to acute base and strongly revolute margin with but few teeth toward the apex. It is endemic to the serpentine mountains of NE Cuba.

Altogether, the Cuban species of *Morella* are four: three endemic to E Cuba (*M. cacuminis*, *M. shaferi* and *M. punctata* (Urb. & Britton) J. Herb.) and one neotropical, widely spread in the island (*M. cerifera* (L.) Small).

### Acknowledgements

The authors gratefully acknowledge the helpful cooperation of Dr W. Greuter.

### References

Herbert J. 2005: New combinations and a new species in *Morella* (*Myricaceae*). – *Novon* **15**: 293–295.

Killick D. J. B., Polhill R. M. & Verdcourt B. 1999: New combinations in African *Myricaceae*. – *Kew Bull.* **53**: 993–995.

Knapp S. 2002: A new combination in *Morella* (*Myricaceae*) in Mesoamerica. – *Novon* **12**: 200.

León [bro.] & Alain [bro.] 1951: Flora de Cuba 2. Dicotiledóneas: Casuarináceas a Meliáceas. – *Contr. Ocas. Mus. Hist. Nat. Colegio “De La Salle”* **10**.

Mabberley D. J. 2008: *Mabberley’s plant-book*, ed. 3. – Cambridge: Cambridge University.

Macdonald A. D. 1989: The morphology and relationships of the *Myricaceae*. – Pp. 147–165 in: Crane P. C. & Blackmore S. (ed.), *Evolution, systematics, and fossil history of the Hamamelidae* **2**. – Oxford: Clarendon.

Parra-O[sorio] C. 2002: New combinations in South American *Myricaceae*. – *Brittonia* **54**: 322–326.

Turner I. M. 2001: New combinations in Malesian *Myricaceae*. – *Gard. Bull. Singapore* **53**: 323–325.

Wilbur R. L. 1994: The *Myricaceae* of the United States and Canada: genera, subgenera, and series. – *Sida* **16**: 93–107.

Wilbur R. L. 2001: Five new combinations in the genus *Morella* (*Myricaceae*) for Neotropical species. – *Rhodora* **103**: 120–122.