

Cryptocoryne ×batangkayanensis (Araceae), a new hybrid from Sarawak

Authors: Ipor, Isa B., Ørgaard, Marian, and Jacobsen, Niels

Source: Willdenowia, 45(2) : 183-187

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.45.45204>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISA B. IPOR¹, MARIAN ØRGAARD² & NIELS JACOBSEN^{2*}

Cryptocoryne ×*batangkayanensis* (Araceae), a new hybrid from Sarawak

Abstract

Ipor I. B., Ørgaard M. & Jacobsen N.: *Cryptocoryne* ×*batangkayanensis* (Araceae), a new hybrid from Sarawak. – Willdenowia 45: 183–187. 2015. – Version of record first published online on 06 July 2015 ahead of inclusion in August 2015 issue; ISSN 1868-6397; © 2015 BGBM Berlin.

DOI: <http://dx.doi.org/10.3372/wi.45.45204>

A new hybrid species, *Cryptocoryne* ×*batangkayanensis* Ipor, Ørgaard & N. Jacobsen (*C. cordata* var. *grabowskii* (Engl.) N. Jacobsen × *C. ferruginea* Engl. var. *ferruginea*), from the Batang Kayan basin, Sarawak, Malaysia, is described and illustrated. It is compared with the parent species and an artificial hybrid of the same parentage.

Additional key words: aroids, taxonomy, relationships

Introduction

Cryptocoryne species have for a long time been relatively well known from Sarawak due to the works of Engler (1879), Schulze (1971), Jacobsen (1985) and Ipor & al. (2007–2008). Presently 12 species of *Cryptocoryne* are recognized from Sarawak.

When several species of *Cryptocoryne* inhabit the same and adjacent stream and river systems, it is not unlikely that hybridization will occur. Natural hybrids between species of *Cryptocoryne* are not uncommon; they have previously been reported from Sri Lanka (Jacobsen 1981, 1987), Thailand (Jacobsen 1980; T. Idei unpub-

lished), Lao P. D. R. (T. Idei unpublished), Peninsular Malaysia (Othman & al. 2009), Singapore (Bastmeijer & Kiew 2001) and Kalimantan (Jacobsen & al. 2002; Bastmeijer & al. 2013). A molecular study of *C. xpurpurea* Ridl. (*C. cordata* Griff. × *C. griffithii* Schott) in Peninsular Malaysia, Sarawak and Kalimantan, presented by Ipor & al. (2010), showed that it was possible to distinguish between hybrid populations. The study also showed that it was possible to separate the Peninsular Malaysian populations (with diploid *C. cordata* var. *cordata* as one of the parents) from the Bornean populations (with tetraploid *C. cordata* var. *grabowskii* (Engl.) N. Jacobsen as one of the parents).

1 Department of Plant Science & Environmental Ecology, Faculty of Resource Science & Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia.

2 Section of Organismal Biology, Department of Plant- and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark; *e-mail: nika@plen.ku.dk (author for correspondence).

Results

During a *Cryptocoryne* field trip in 2004, I.B.I. brought us to a site west of Bau (some 55 km west of Kuching), in the eastern part of the Batang Kayan catchment area. In small forest-ditch pools we saw a green-leaved *Cryptocoryne* population which according to I.B.I.'s information produced a spathe with a purple limb. The plant was taken into cultivation as accession *NJS 04-07* and indeed soon proved to have a purple limb, thereby resembling *C. xpurpurea* nothovar. *purpurea* of Peninsular Malaysia (*C. cordata* var. *cordata* × *C. griffithii*, $2n = 34$), and the South Kalimantan counterpart *C. xpurpurea* nothovar. *borneoensis* N. Jacobsen & al. (*C. cordata* var. *grabowskii* × *C. griffithii*, $2n = 51$; Jacobsen & al. 2002; Othman & al. 2009).

Ipor & al. (2008), in their treatment of the ecology and distribution of the *Cryptocoryne* species in Sarawak, pictured and assigned the Batang Kayan plant with uncertain status under *C. xpurpurea* nothovar. *borneoensis*.

Although the Batang Kayan plant resembled the hybrid *Cryptocoryne xpurpurea*, there was a problem: one of the parental species of this hybrid, *C. griffithii*, has only been reported from the southern part of Kalimantan, Peninsular Malaysia and some of the nearby Riau islands. This fact left some uncertainty as to the Batang Kayan plant belonging to *C. xpurpurea*.

The other parental taxon of *Cryptocoryne xpurpurea* nothovar. *borneoensis*, viz. *C. cordata* var. *grabowskii*, is widespread on Borneo, including Sarawak, and has been found in Sungai Stunggang, also in the Batang Kayan basin (less than 20 km north of the Batang Kayan locality), as well as in the adjacent Sungai Sarawak basin toward the east.

Another parent candidate with a purple spathe limb within the region would be *Cryptocoryne ferruginea* Engl. var. *ferruginea*, which has been found some 20 km



Fig. 1. *Cryptocoryne xbatangkayanensis* – A: habitat of type locality; B: habitat close up. – Both photographs taken on 29 Nov 2004 by N. Jacobsen.

east of the Batang Kayan locality at Sungai Salalang (north of Bau) within the neighbouring Sungai Sarawak catchment basin.

The relatively small Sungai Salalang and its tributaries lies toward the eastern border the Batang Kayan basin some 15 km east of the Batang Kayan locality; this rather short distance to the hybrid locality makes a hybridization event possible, especially since the intervening area has not been investigated for *Cryptocoryne* populations. Moreover, even though at the present time the two catchment areas have separate runoffs, they had combined runoffs during the last glaciation and until about 13000 BP (Voris 2000; Sathiamurthy & Voris 2006).



Fig. 2. *Cryptocoryne xbatangkayanensis* ($2n = 85$) – A: limb of spathe; B: opened kettle showing female and male flowers. – Photographs by N. Jacobsen.

Sarawak *Cryptocoryne* species have been included in crossing experiments performed over the years at the University of Copenhagen. One of the successful combinations is CyCy 447, i.e. *C. ferruginea* var. *ferruginea* (female) \times *C. cordata* var. *grabowskii* (male), $2n = 51$, reported here [*C. ferruginea* var. *ferruginea*: Malaysia, Sarawak, Samarahan Div., Balai Ringin, Waser SK 05-02, $2n = 34 \times$ *C. cordata* var. *grabowskii*: Malaysia, Sarawak, Sri Aman Div., Sungai Semawak, IBI 05-1, $2n = 68$]. The hybrid CyCy 447 has a longer spathe limb, but otherwise the colour and surface of the spathe limb matches rather well the Batang Kayan plants.

The Batang Kayan population has proved to have a chromosome number of $2n = 85$. This is not a straightforward number, although an explanation could be that it evolved from a hybridization between an unreduced gamete from *Cryptocoryne cordata* var. *grabowskii* (68 chromosomes) and a normal reduced gamete from *C. ferruginea* var. *ferruginea* (17 chromosomes), that is $68 + 17 = 85$.

Usually hybrids are somewhat intermediate between the parents, and if the parents have different chromosome numbers, e.g. diploids and tetraploids, the parent that contributes the most chromosomes gives rise to a genetic dosage effect in the resulting morphology of the hybrid. This phenomenon is well known, and has been shown in hybridization experiments in, e.g., *Hordeum* L. (barley) (Bothmer & Jacobsen 1986) and *Crocus* L. (Ørgaard & al. 1995). The morphological differences between the Batang Kayan population ($2n = 85$) and the artificial hybrid CyCy 447 ($2n = 51$) can therefore be

explained by genome dosage effects.

Unpublished results of an AFLP study of the Malaysian cordate-leaved species of *Cryptocoryne* also show that the Batang Kayan plants are genetically closest to CyCy 447, and secondly to *C. cordata* var. *grabowskii*. Additionally, the Batang Kayan plants group with neither *C. cordata* var. *cordata*, nor *C. griffithii*, nor *C. xpurpurea*.

We conclude that the Kampong Stungkor Baru population in the Batang Kayan basin represents an interspecific hybrid between *Cryptocoryne cordata* var. *grabowskii* and *C. ferruginea* var. *ferruginea* (alphabetically ordered).

Cryptocoryne xbatangkayanensis Ipor, Ørgaard & N. Jacobsen, **nothosp. nov.** – Fig. 1 & 2.

Holotype: Malaysia, Sarawak, Kuching Div., Batang Kayan basin, near Kampong Stungkor Baru, I. B. Ipor & N. Jacobsen NJS 04-07 (SAR; isotypes: B, C, L).

Description — Leaves 15–25 cm long; blade ovate, 3–10 cm long, 2–6 cm wide, surface smooth to somewhat irregular, lower surface \pm purplish, upper surface bright green to somewhat purplish mottled, sometimes brownish on midvein, base cuneate to \pm cordate. Spathe 8–20 cm long; kettle 1.5–2 cm long, inside purple in upper part, spotted into tube; tube whitish, 5–12 cm long; limb bright shining red-purple (fading with age), ovate, 2–6 cm long, slightly rugose with scattered regular small bumps, apex acuminate; collar absent, collar zone dark red to reddish, more whitish toward tube opening, broad. Spadix with 5–7 female flowers, white to brownish in upper part; stigmas elliptic; olfactory bodies dark yellowish, rounded; male flowers 40–60; appendix whitish to slightly purple spotted; pollen fertility 0–5%. Infructescence unknown.

Chromosome number — $2n = 85$, reported here.

Distribution — Only known from the type locality area.

Ecology — Formerly secondary forest and partly abandoned rubber farm. Muddy bottom of a small drainage ditch after the establishment of a diversion stream. Presently most parts of the habitat have been cleared for oil palm planting.

Conservation status — The area where *Cryptocoryne xbatangkayanensis* is found is within a partly abandoned rubber plantation in secondary forest, and it is foreseeable that the area will be cleared in the near future, although small populations in the vicinity, surprisingly enough, seem to be self-sustainable in oil palm planting areas.

Acknowledgements

Karen Rysbjerg Munk skilfully prepared the slides for the chromosome counting. The Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, kindly provided greenhouse facilities for the cultivation of material. The authors are grateful to the Ministry of Higher Learning Malaysia and University Malaysia Sarawak for financial support (grant: NRGs/1088/2013(2)) and to the Department of Forestry Sarawak for providing the research permit. The authors express their appreciation to James Abai and Prof. Cheksum Tawan for their assistance. Dr Josef Bogner and an anonymous reviewer are also thanked for their useful and constructive comments on an earlier draft of this paper.

References

- Bastmeijer J. D. & Kiew R. 2001: A New *Cryptocoryne* hybrid (*Araceae*) from the Bukit Timah Nature Reserve, Singapore. – *Gard. Bull. Singapore* **53**: 9–17.
- Bastmeijer J. D., Kishi H., Takahashi N., Wongso S. & Jacobsen N. 2013: Eine neue Varietät der *Cryptocoryne ferruginea* Engl. von Sekadau, West Kalimantan, Indonesien. – *Aqua Pl.* **38**(3): 84–93.
- Bothmer R. von & Jacobsen N. 1986: Interspecific crosses in *Hordeum* (*Poaceae*). – *Pl. Syst. Evol.* **153**: 49–64.

Engler A. 1879: *Araceae*, specialmente Bornensi e Papuane raccolte da O. Beccari. – *Bull. R. Soc. Tosc. Ort.* **4**: 295–302 (with an introduction by O. Beccari on pp. 265–266).

- Ipor I. B., Tawan C. S., Abai J., Saupi N. & Meekiong K. 2007–2008: Oekologie und Verbreitung der *Cryptocoryne*-Arten (*Araceae*) in Sarawak (Malaysia) Teil 1–4. – *Aqua Pl.* **32**(3): 101–106; **32**(4): 131–140; **33**(1): 22–27; **33**(2): 48–52.
- Ipor I. B., Tawan C. S., Jacobsen N., Wongso S., Idei T., Ariff N. H. M. & Roslan H. A. 2010: Genotyping natural hybrids of *Cryptocoryne purpurea* in Borneo and Peninsular Malaysia. – *J. Trop. Biol. Conserv.* **7**: 81–86.



Fig. 3. Artificial hybrid: CyCy 447 ($2n = 51$): *Cryptocoryne ferruginea* var. *ferruginea* × *C. cordata* var. *grabowskii* – A: plant in cultivation; B: limb of spathe; C: opened kettle showing female and male flowers. – Photographs by N. Jacobsen.

Jacobsen N. 1980: The *Cryptocoryne albida* group of mainland Asia (*Araceae*). – Misc. Pap. Landbouwhoogeschool **19**: 183–204.

Jacobsen N. 1981: *Cryptocoryne undulata* Wendt und Bemerkungen zu anderen Arten – ein Nachtrag. – Aqua Pl. **6**(4): 92–94.

Jacobsen N. 1985: The *Cryptocoryne* (*Araceae*) of Borneo. – Nordic J. Bot. **5**: 31–50.

Jacobsen N. 1987: *Cryptocoryne* in: Dassanayake M. D. & Fosberg F. R. (ed.). A revised handbook to the flora of Ceylon **VI**: 85–99.

Jacobsen N. 2002: Der *Cryptocoryne cordata* Griffith-Komplex (*Araceae*) in Malesien. – Aqua Pl. **27**(4): 150–151.

Jacobsen N., Bastmeijer J. D. & Sasaki Y. 2002: *Cryptocoryne xpurpurea* Ridley nothovar. *borneoensis* N. Jacobsen, Bastmeijer & Y. Sasaki (*Araceae*), eine neue Hybridvarietät aus Kalimantan. – Aqua Pl. **27**(4): 152–154.

Ørgaard M., Jacobsen N. & Heslop-Harrison J. S. 1995: The hybrid origin of two cultivars of *Crocus* (*Iridaceae*) analysed by molecular cytogenetics including genomic southern and in situ hybridization. – Ann. Bot. **76**: 253–262.

Othman A. S., Jacobsen N. & Mansor M. 2009: *Cryptocoryne* of Peninsular Malaysia. – Pulau Pinang: Penerbit University Sains Malaysia.

Sathiamurthy E. & Voris H. K. 2006: Maps of Holocene sea level transgression and submerged lakes on the Sunda Shelf. – Nat. Hist. J. Chulalongkorn Univ. Suppl. **2**: 1–43.

Schulze J. 1971: *Cryptocorynen* aus Sarawak I–IV. – Aquarien Terrarien Z. **24**: 230–233, 267–270, 303–306, 336–339.

Voris H. K. 2000: Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. – J. Biogeogr. **27**: 1153–1167.



Fig 4. Parents of CyCy 447: A & B: *Cryptocoryne ferruginea* var. *ferruginea*, Malaysia, Sarawak, Sri Aman Div., Balai Ringin, Waser SK 05-02, $2n = 34$; C & D: *C. cordata* var. *grabowskii*, Malaysia, Sarawak, Sri Aman Div., Sg. Semawak, IBI 05-1, $2n = 68$. – Photographs by N. Jacobsen.