



Contents of Willdenowia 52

Source: Willdenowia, 52(3) : 436

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.52.52311>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Contents of Willdenowia 52

| | |
|--|---------------|
| Anderberg A. A. & Bengtson A.: Taxonomic novelties in the <i>Asteraceae</i> – <i>Inuleae</i> with the description of a new genus, <i>Galgera</i> separate from <i>Laggera</i> | 373 |
| Bécquer E. R., Bochorny T., Gavrutenko M. & Michelangeli F. A.: A revision of the “basal-axile placentation clade” of <i>Miconieae</i> , the newly erected <i>Miconia</i> sect. <i>Liogieria</i> (<i>Melastomataceae</i> : <i>Miconieae</i>) from the Greater Antilles | 387 |
| Burgt X. M. van der, Haba P. M., Magassouba S. & Veranso-Libalah M. C.: <i>Benna alternifolia</i> (<i>Melastomataceae</i> : <i>Sonerileae</i>), a new herbaceous genus and species from Guinea, West Africa | 25 |
| Chatelain C., Uotila P., Benhouhou S., Mombrial F., Mesbah M., Baa S. & Benghanem A. N.: <i>Chenopodium hoggarensis</i> (<i>Amaranthaceae</i>), a new species from Algeria and Chad | 75 |
| Dillenberger M. S. & Kadereit J. W.: The distinction between <i>Sagina apetala</i> and <i>S. micropetala</i> (<i>Caryophyllaceae</i> : <i>Sagineae</i>), their phylogenetic relationships, and a note on the coastal origin of some widespread ruderals | 5 |
| Fernandez-Hilario R., Rojas Gonzáles R. del P., Villanueva-Espinoza R., Lajo L., Wong Sato A. A., Paredes-Burneo D., Pillaca-Huacre L., Michelangeli F. A. & Goldenberg R.: Nine new species and a new country record for <i>Meriania</i> (<i>Melastomataceae</i>) from Peru | 39 |
| Firat M., Özüdoğru B. & Yıldırım H.: A new bellflower, <i>Campanula dersimensis</i> (<i>Campanulaceae</i>), from E Anatolia, Turkey | 167 |
| Fischer E. & Lobin W.: The genus <i>Isoetes</i> (<i>Isoetaceae</i>) in Central Africa (Democratic Republic of the Congo, Rwanda, Burundi) with the description of three new species | 315 |
| González-Zamora P., Aquino D., Mohl J. & Sánchez D.: A new endemic species of <i>Mammillaria</i> (<i>Cactaceae</i>) from San Luis Potosí, Mexico | 359 |
| Güzel M. E., Kilian N., Sennikov A. N., Coşkunçelebi K., Makbul S. & Gültepe M.: <i>Caucasoseris</i> , a new genus of subtribe <i>Chondrillinae</i> (<i>Asteraceae</i> : <i>Cichorieae</i>) for the enigmatic <i>Prenanthes abietina</i> | 103 |
| Kaehler M. & Lohmann L. G.: Taxonomic revisions in <i>Fridericia</i> (<i>Bignoniaceae</i> , <i>Bignoniaceae</i>) II: the “Neomac-fadya” clade | 247 |
| Oberprieler C., Töpfer A., Dorfner M., Stock M. & Vogt R.: An updated subtribal classification of <i>Compositae</i> tribe <i>Anthemideae</i> based on extended phylogenetic reconstructions | 117 |
| Poulsen A. D., Pomoso P. & Magun T.: Three species of <i>Etilingera</i> (<i>Zingiberaceae</i>) recollected in the footsteps of Rudolf Schlechter in Papua New Guinea | 153 |
| Raab-Straube E. von & Raus Th. (ed.): Euro+Med-Checklist Notulae, 15 | 273 |
| Raus Th.: Taxonomic, nomenclatural and floristic review of <i>Amaranthaceae</i> of Greece and neighbouring countries | 335 |
| Schmidt A. R., Korall P., Krings M., Weststrand S., Bergschneider L., Sadowski E.-M., Bechteler J., Rikkinen J. & Regalado L.: <i>Selaginella</i> in Cretaceous amber from Myanmar | 179 |
| Schwerdtfeger M.: Book review: Dörken V. M., Edwards D., Ladd P. G. & Parsons R. F., The four dimensions of terrestrial plants: reproduction, structure, evolution and ecology | 313 |
| Stoyanov S., Marinov Y., Apostolova-Stoyanova N., Randelović V. & Vukojičić S.: Taxonomic review of <i>Scrophularia</i> sect. <i>Tomiphyllum</i> in Bulgaria and Serbia: the case of <i>Scrophularia bulgarica</i> | 303 |
| Troia A.: The unnoticed northward expansion of <i>Najas marina</i> subsp. <i>armata</i> (<i>Hydrocharitaceae</i>) in the Mediterranean area: an effect of climate change? | 91 |
| Vogt R. & Gottschlich G.: Ferdinand Tessoroff (1879–1924) and his <i>Hieracium</i> collection | 83 |
| Book review | 303 |
| Indexes to new names and combinations appearing in Willdenowia 52 | 151, 301, 433 |
| Indexes to typifications of names in Willdenowia 52 | 152, 302, 434 |
| Reviewers of manuscripts submitted for publication during 2021 | 435 |
| Contents of Willdenowia 52 | 436 |