

Land Stewardship and Protection of an Endangered Plant Species in an Insular World Biosphere Reserve

Author: Medina, Félix M.

Source: Natural Areas Journal, 41(3) : 209-212

Published By: Natural Areas Association

URL: <https://doi.org/10.3375/043.041.0307>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Land Stewardship and Protection of an Endangered Plant Species in an Insular World Biosphere Reserve

Félix M. Medina^{1,2}

¹Servicio de Medio Ambiente, Cabildo Insular de La Palma, Avenida Los Indianos 20, 38700 Santa Cruz de La Palma, Canary Islands, Spain; felix.medina@cablapalma.es

²Island Ecology and Evolution Research Group (IPNA-CSIC), Astrofísico Francisco Sánchez 3, 38206 La Laguna, Tenerife, Canary Islands, Spain

Associate Editor: Donatella Cogoni

ABSTRACT

Lotus eremiticus is an endemic species from La Palma World Biosphere Reserve. It has a small distribution range, low population size, and is threatened by introduced herbivores. Since these threats have not been removed from the protected area, they were excluded by building a fence. The land where the species grows is private property, so an agreement with the landowner was reached to permit measures to favor its recovery. During 2008–2019, as a result of this agreement and the conservation efforts, a large population increase occurred, from the initial 5 individuals to the 30 plants that are currently distributed at the original site. Furthermore, these measures allow the species to maintain stable population dynamics, meaning that this endangered species is itself capable of recovery if the threat is removed. This is a good example of how land stewardship is an effective tool to conserve endangered species.

Index terms: invasive herbivores; landowners; *Lotus eremiticus*; oceanic islands; recovery plan; stewardship; threatened species

Biosphere Reserves are recognized areas of representative environments that have been internationally designated within the framework of UNESCO's MAB Programme for their value to conservation, through providing the scientific knowledge, skills, and values to support sustainable development (Bridgewater and Cresswell 1998). La Palma (one of the Canary Islands, Spain) was entirely declared a World Biosphere Reserve in 2002 since it hosts a rich biodiversity at regional and national scales. More than 5434 terrestrial species (1056 of them endemic) have been listed in its territory (Martín et al. 2005a), including at least 879 vascular plant species (20% of them endemic to the island).

Habitat destruction and especially the introduction of invasive species are the most serious threats to the conservation of native species in the Canary Islands (Whittaker and Fernández-Palacios 2007). Introduced herbivorous mammals are the most important threat to the conservation of the endemic and threatened plant species (García-Casanova et al. 2001) and some of them such as goats (*Capra hircus*) and European rabbits (*Oryctolagus cuniculus*) are considered among the worst invasive species worldwide (Lowe et al. 2000). These destructive species, as well as domesticated (*Ovis aries*) and Barbary sheep (*Ammotragus lervia*) (Nogales et al. 2006) are currently affecting native plant species in La Palma World Biosphere Reserve (see Garzón-Machado et al. 2010; Irl et al. 2012).

A total of 366 species are considered as threatened in the Canary archipelago (Martín et al. 2005b), and 22 of them (15 plants) are present on La Palma (Martín et al. 2005a). One of these is *Lotus eremiticus* A. Santos (Fabaceae), a creeping perennial and hermaphroditic chamaephyte, with little development, thin branches, and small leaflets 1 cm long (Santos 1983) (Figure 1). Flowering is showy and abundant in the months of April and May. It shows self-compatibility; fruiting,

which occurs July–August, is scarce but produces fertile seeds. More aspects of its reproductive biology are unknown, such as its pollinators, although the role of birds and reptiles as pollinators of the *Lotus* genus has been described (Ojeda et al. 2012; Siverio and Rodríguez-Rodríguez 2012). *L. eremiticus* is an endemic plant of La Palma and it is threatened by introduced herbivores (goats and rabbits), rats (*Rattus rattus*), and its low population and scarce distribution, formerly only five plants in one location (Medina and Hernández Martín 2015). For this reason, the species was included in the Spanish List of Endangered Species as in danger of extinction (IUCN category: Critically Endangered; Martín-Cáceres et al. 2011), and a recovery plan was approved by the Canary Islands Government and started in 2006. As a high-priority measure, this plan included increasing the current distribution of the species, as well as its number of individuals and populations, through reintroduction of plants obtained from seeds and cuttings collected from the natural population. Following the recommendations of the recovery plan, this action would be achieved by promoting legal action to acquire the properties where this endangered species is distributed. A second important objective, and taking into account that there are no other human impacts on the location besides uncontrolled livestock, was to reduce or eliminate herbivore pressure, thus improving the environmental conditions in the species' natural habitat.

Lotus eremiticus is located in only one place, Roque de la Viña, Don Pedro, Garafía (UTM: 28R 218720.35 N; 3193075.39 W; 300 m a.s.l.; Figure 2), inside a protected area classed as a strict nature reserve within Natura 2000 Net (IUCN category I). Vegetation is mainly composed of *Rhamnus crenulata*–*Junipero canariensis* and *Myrica fayae*–*Ericetum arboreae* associations with rocky elements of *Greenovia*–*Aeonietea*. The main conservation



Figure 1.—Detail of the endemic endangered species *Lotus eremiticus*.

objective of this protected area is to maintain the population of the species through reducing damage from introduced herbivores. Livestock grazing is prohibited in this protected area, but it was previously a traditional activity for centuries. Consequently, large numbers of free-ranging or feral goats and sheep are still present in the area, affecting this threatened species as well as other native plants.

Control or eradication of feral livestock is feasible on islands and an important tool for plant and environment conservation (Campbell and Donlan 2005). However, culling animal populations is an important source of conflict with animal rights associations that wish to prevent any animal suffering (Hutchins and Wemmer 1986). This ethical issue frequently hinders the approval of eradication campaigns by stakeholders and decision makers (see Lohr et al. 2014). As a result of this, no management plan or decision had been considered by the Island Council (Cabildo) to remove livestock from this protected area. So, to conserve this endangered plant species, the only possible solution was to install an enclosure fence to prevent the effects of herbivores. This has proved to be an effective measure in avoiding the extinction of endemic plant species (Garzón-Machado et al. 2010; Cogoni et al. 2013; Fenu et al. 2016), at least while control or eradication programs are being applied (Fenu et al. 2020).

Despite the location of *L. eremiticus* being inside a protected area, it is on private property. Current law allows the Island Council to implement conservation schemes based on the management plan of this protected area. However, to avoid conflicts with landowners, it was important to achieve their support for the conservation issues and measures involved in protecting this endangered species (Vickerman 1999; Olive and Raymond 2010). Nevertheless, in our case, the owners wanted to receive something in return because they had rented their properties to farmers and herdsmen as a place to obtain forage for their livestock. Furthermore, several years ago the landowners offered the Canary Government the chance to purchase the area, but a limited budget did not permit this, which is a common cause of failure in land acquisition (Wilcove and Chen



Figure 2.—Roque de la Viña, locus classicus of *Lotus eremiticus* in La Palma Island Biosphere Reserve.

1998; Kamal et al. 2015). Taking into account the cost of buying this property, other ways to involve landowners in implementing management plans should be considered (Farmer et al. 2017). A range of incentives have been proposed to convince landowners to collaborate in land conservation and endangered species (Wilcove and Chen 1998; Vickerman 1999; Shogren et al. 2003; Kabii and Horwitz 2006; Kamal et al. 2015), including broad legal mechanisms such as conservation easements, covenants, or development agreements (Kabii and Horwitz 2006; Rissman et al. 2007; Ernst and Wallace 2008). However, there are few reports where stewardship options have been mentioned as a management option to conserve specific endangered species. This is probably due to landowners usually being unaware of endangered species on their properties (Olive and McCune 2017). Some specific projects have been carried out in the case of the Florida panther (*Felis concolor coryi*; Maehr 1990; Kreye and Pienaar 2015), the crested caracara (*Caracara cheriway*; Morrison and Humphrey 2001), or the Japanese murrelet (*Synthliboramphus wumizusume*; Hyde 2019).

From among the possible incentives, a rental agreement with the landowner was finally selected as the best solution to allow us to take appropriate decisions to protect this endangered plant species. Fortunately, the location of *L. eremiticus* is in a single property of about 12 ha belonging to just one person, which enormously facilitated the agreement. After several meetings with the owner and the participation of a conservationist neighbor who helped us to contact him, the lease agreement was signed in 2008 for a minimum of 5 y with the option to extend this or purchase. During 2014, the first lease was extended because the required budget for purchase was not approved. Unfortunately, the owner died that year and signing the second agreement was delayed until 2016 as his heirs resolved all the legal issues. However, they allowed us to continue with our conservation work on the property without an official agreement. Currently, the contract is valid until 2022 with a cost of 2200 Euros (approx. \$2445 US) per year, meaning a total expenditure from 2008 to 2019 of 23,500 Euros (\$26,119 US).

The agreement allows us to manage the property and fence the area surrounding where *L. eremiticus* is distributed, reaching an

approximate surface area of 0.55 ha. Once the impact of livestock was eliminated, we had to control the presence of rabbits and rats. To do this, three Tomahawk-type live traps were installed to capture the rabbits. After 4 wk without trapping any rabbits, and checking that there were no rabbit remains or signs inside, the fence traps were removed, and new damage by this species has not been subsequently observed. To minimize rat impact, which was periodically observed during the driest seasons in the area (summer), rat bait with second-generation anticoagulants (Difenacoum 0.005%) was employed, poisoning being one of the most common methods used to control this invasive species on islands (Duron et al. 2017).

As a result of these conservation measures, in the period between 2010 (completion of fence) and 2019 (end of first phase of the recovery plan), the natural population of *L. eremiticus* increased to 30 individuals, growing from the natural soil seedbank. In this same period, natural population dynamics were observed: a total of 89 plants had grown from the seedbank, but 62 died, probably due to the drought during those years. Another aim of the recovery plan was to reinforce the population. To do so, the total initial population (five individuals in only one location) was fortnight monitored by analyzing plant size and reproductive stage. This allows us to collect plant material (cuts and seeds), which were treated in greenhouse before their reintroduction in the same place and using substrate from the original location. Planting took place after rainfall in spring or autumn and providing water for a week during the next year to ensure the plants had taken root. Initial genetic analysis showed the species has a very low level of genetic variation (Pérez-Vargas et al. 2021), which allows us to maintain the original genetic set of the population along the recovery process. This process was carried out during this project on several occasions with a total of 228 cuttings obtained from the natural population. At the end of the recovery program (January 2019), a total of 77 *L. eremiticus* plants were counted inside the fence. Of these, 25 had grown directly from the soil seedbank, 28 were grown in the nursery greenhouse from seeds collected from naturally growing plants and reintroduced by transplanting them into the area, and 24 were reintroduced from cuttings grown in the natural area. An important successful result of this project was the natural regeneration of the *L. eremiticus* population originating from the reintroduced cuttings. Two new plants were still growing after 9 y, pointing to the capacity of this species to naturally regenerate its population if the main threat disappears.

New potential suitable sites for this species were defined throughout predictive analysis applying GIS (Geographical Information Systems) tools (Bermejo-Domínguez et al. 2007), which allow us to increase species distribution. The implementation of this agreement and the results obtained allowed us to establish another new land stewardship with the town council of Garafia, the municipality where *L. eremiticus* is found. Once they knew of the work done to favor recovery of this species and the suitable new place, the council decided to provide us a public property where a new population was introduced by the recovery plan team. In May 2017, 40 plants were introduced by local people involved in this project. After an initial 95% success, to date only 26 plants are still in the area. Plant care and

attention has been provided by local people and environmental workers, always under technical supervision of the recovery plan technicians, as another example of how land stewardship helps to conserve native and endangered species (Ernst and Wallace 2008). Another example of the involvement of the local residents and neighbors in the conservation of this species took place in December 2018, when a landslide caused serious damage to the fence, allowing several goats to enter. A local farmer particularly fond of the plant and who wishes to conserve it (Ernst and Wallace 2008) because his aunt was the person who discovered this species in the 1950s, acted rapidly to remove three goats from inside the plot and partially repair the fence.

As an example of land stewardship, this agreement has been considered a fundamental tool during the project to preserve this endangered species: “*Lotus eremiticus* in La Palma World Biosphere Reserve.” Moreover, it has been very useful to protect its habitat as well as other native and protected plant species present in the same area, although this project was not foreseen or run as a multispecies recovery plan (Langpap and Kerkvliet 2012). Talking to local residents, neighbors, and landowners provides important information for making appropriate management decisions (Henderson et al. 2014). Promoting environmental conservation through educational programs is also considered a basic instrument to engage landowners and other citizens, to gain their support in stewardship activities (Farmer et al. 2017).

ACKNOWLEDGMENTS

Our thanks to Néstor Rodríguez Pérez and his heirs for their willingness to reach the necessary agreement to adequately manage this endangered species. Vicente García López helped us contact the landowners and continues to be engaged in the project. Guillermo Hernández Martín and Segundo Lorenzo Aguiar participated in all the programs of this recovery scheme. We are also grateful to all the staff of the Island and Town Councils (Cabildo Insular de La Palma, Ayuntamiento de la Villa de Garafia) for their work and support in conserving this species.

LITERATURE CITED

- Bermejo-Domínguez, J.A., P.L. Pérez de Paz, and M.J. del Arco Aguiar. 2007. Aplicación de los sistemas de información geográfica (SIG) en el análisis predictivo de flora en peligro de extinción: *Lotus eremiticus*. Santos (Isla de La Palma – Islas Canarias). Revista de Estudios Generales de la Isla de La Palma 3:513–541.
- Bridgewater, P.B., and I.D. Cresswell. 1998. The reality of the world network of Biosphere Reserves: Its relevance for the implementation of the Convention on Biological Diversity. Pp. 1–6 in Proceedings of a Workshop at the 1996 IUCN World Conservation Congress, Montreal, Canada. Biosphere Reserves – Myth or Reality? IUCN, Gland, Switzerland and Cambridge, UK.
- Campbell, K., and J. Donlan. 2005. Feral goat eradications on islands. Conservation Biology 19:1362–1374.
- Cogoni, D., G. Fenu, E. Concas, and G. Bacchetta. 2013. The effectiveness of plant conservation measures: The *Dianthus morisianus* reintroduction. Oryx 47:203–206.
- Duron, Q., A.B. Shiels, and E. Vidal. 2017. Control of invasive rats on islands and priorities for future action. Conservation Biology 31:761–771.

- Ernst, T., and G.N. Wallace. 2008. Characteristics, motivations, and management actions of landowners engaged in private land conservation in Larimer County Colorado. *Natural Areas Journal* 28:109–120.
- Farmer, J.R., Z. Ma, M. Drescher, E.G. Knackmuhs, and S.L. Dickinson. 2017. Private landowners, voluntary conservation programs, and implementation of conservation friendly land management practices. *Conservation Letters* 10:58–66.
- Fenu, G., D. Cogoni, and G. Bacchetta. 2016. The role of fencing in the success of threatened plant species translocation. *Plant Ecology* 217:207–217.
- Fenu, G., G. Bacchetta, C.S. Christodoulou, D. Cogoni, C. Fournaraki, G. del Galdo Gian Prieto, P. Gotsiou, A. Kyratzis, C. Piazza, M. Vicens, and B. de Montmollin. 2020. A common approach to the conservation of threatened island vascular plants: First results in the Mediterranean Basin. *Diversity* 12:157.
- García Casanova, J., J.L. Rodríguez Luengo, and C. Rodríguez Piñero. 2001. Especies amenazadas. Pp. 167–172 in J.M. Fernández-Palacios and J.L. Martín Esquivel, eds., *Naturaleza de las Islas Canarias. Ecología y Conservación*. Publicaciones Turquesa, S.L. Santa Cruz de Tenerife, Spain.
- Garzón-Machado, V., J.M. González-Mancebo, A. Palomares-Martínez, A. Acevedo-Rodríguez, J.M. Fernández-Palacios, M. del-Arco-Aguilar, and P.L. Pérez de Paz. 2010. Strong negative effect of alien herbivores on endemic legumes of the Canary pine forest. *Biological Conservation* 143:2685–2694.
- Henderson, A.E., M. Reed, and S.K. Davis. 2014. Voluntary stewardship and the Canadian Species at Risk Act: Exploring rancher willingness to support species at risk in the Canadian prairies. *Human Dimensions of Wildlife* 19:17–32.
- Hutchins, M., and C. Wemmer. 1986. Wildlife conservation and animal rights: Are they compatible? Pp. 111–137 in M.W. Fox and L.D. Mickley, eds., *Advances in Animal Welfare Science 1986/87*. Humane Society of the United States, Washington, DC.
- Hyde, C. 2019. The mark of the Japanese murrelet (*Synthliboramphus wumizusume*): A study of song and stewardship in Japan's inland sea. Senior Thesis, Pomona College, Claremont, CA.
- Irl, S.D.H., M. Steinbauer, W. Babel, C. Beierkuhnlein, G. Blume-Werry, J. Messinger, A. Palomares Martínez, S. Strohmeier, and A. Jentsch. 2012. An 11-yr enclosure experiment in a high-elevation island ecosystem: Introduced herbivore impact on shrub species richness, seedling recruitment and population dynamics. *Journal of Vegetation Science* 23:1114–1125.
- Kabii, T., and P. Horwitz. 2006. A review of landholder motivations and determinants for participation in conservation covenanting programmes. *Environmental Conservation* 33:11–20.
- Kamal, S., M. Grodzińska-Jurczak, and G. Brown. 2015. Conservation on private land: A review of global strategies with a proposed classification system. *Journal of Environmental Planning and Management* 58:576–597.
- Kreye, M.M., and E.F. Pienaar. 2015. A critical review of efforts to protect Florida panther habitat on private lands. *Land Use Policy* 48:428–436.
- Langpap, C., and J. Kerkvliet. 2012. Endangered species conservation on private land: Assessing the effectiveness of habitat conservation plans. *Journal of Environmental Economics and Management* 64:1–15.
- Lohr, C.A., C.A. Lepczyk, and E.D. Johnson. 2014. The islands are different: Human perception of game species in Hawaii. *Environmental Management* 54:814–827.
- Lowe, S., M. Browne, S. Boudjelas, and M. De Poorter. 2000. 100 of the world's worst invasive alien species – A selection from the Global Invasive Species Database. *Aliens* 12:1–12.
- Maehr, D.S. 1990. The Florida panther and private lands. *Conservation Biology* 4:167–170.
- Martín, J.L., M. Marrero, N. Zurita, M. Arechavaleta, and I. Izquierdo. 2005a. Biodiversidad en gráficas. Especies silvestres de las Islas Canarias. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias.
- Martín, J.L., S. Fajardo, M.A. Cabrera, M. Arechavaleta, A. Aguiar, S. Martín, and M. Naranjo. 2005b. Evaluación 2004 de especies amenazadas de Canarias. Especies en peligro de extinción, sensibles a la alteración de su hábitat y vulnerables. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias.
- Martín Cáceres, K., R. Mesa Coello, and A. Santos Guerra. 2011. *Lotus eremiticus*. The IUCN Red List of Threatened Species 2011: e.T165218A5991682. Accessed 29 Apr 2020 from <<https://dx.doi.org/10.2305/IUCN.UK.2011-1.RLTS.T165218A5991682.en>>.
- Medina, F.M., and G. Hernández Martín. 2015. Conservación de dos especies amenazadas de la flora endémica de la Isla de La Palma. *Conservación Vegetal* 19:1–3.
- Morrison, J.L., and S.R. Humphrey. 2001. Conservation value of private lands for crested caracaras in Florida. *Conservation Biology* 15:675–684.
- Nogales, M., J.L. Rodríguez-Luengo, and P. Marrero. 2006. Ecological effects and distribution of invasive non-native mammals on the Canary Islands. *Mammal Review* 36:49–65.
- Ojeda, I., A. Santos-Guerra, R. Jaén-Molina, F. Oliva-Tejera, J. Caujapé-Castells, and Q. Cronk. 2012. The origin of bird pollination in Macaronesian *Lotus* (Loteae, Leguminosae). *Molecular Phylogenetics and Evolution* 62:306–318.
- Olive, A., and J.L. McCune. 2017. Wonder, ignorance, and resistance: Landowners and the stewardship of endangered species. *Journal of Rural Studies* 49:13–22.
- Olive, A., and L. Raymond. 2010. Reconciling norm conflict in endangered species conservation on private land. *Natural Resource Journal* 50:431–454.
- Pérez-Vargas, I., A.M. Portero Álvarez, P.L. Pérez de Paz, and J.A. Pérez. 2021. Retrotransposon-based molecular markers as a tool in delimiting species in section *Ryncholotus*, a recent radiation group of Macaronesian *Lotus*. *Systematics and Biodiversity* 19:110–120.
- Rissman, A.R., L. Lozier, T. Comendant, P. Kareiva, J.M. Kiesecker, M.R. Shaw, and A.M. Merenlender. 2007. Conservation easements: Biodiversity protection and private use. *Conservation Biology* 21:709–718.
- Santos, A. 1983. Vegetación y flora de La Palma. Editorial Interinsular Canaria. Santa Cruz de Tenerife.
- Shogren, J.F., G.M. Parkhurst, and C. Settle. 2003. Integrating economics and ecology to protect nature on private lands: Models, methods, and mindsets. *Environmental Science and Policy* 6:233–242.
- Siverio, F., and M.C. Rodríguez-Rodríguez. 2012. *Gallotia galloti* (Canary lizard). Nectarivory. *Herpetological Review* 43:333–334.
- Vickerman, S. 1999. A state model for implementing stewardship incentives to conserve biodiversity and endangered species. *Science of the Total Environment* 240:41–50.
- Whittaker, R.J., and J.M. Fernández-Palacios. 2007. *Island Biogeography – Ecology, Evolution, and Conservation*. 2nd ed. Oxford University Press, Oxford, UK.
- Wilcove, D.S., and L.Y. Chen. 1998. Management costs for endangered species. *Conservation Biology* 12:1405–1407.