

Nesting Ecology of the Collared Scops Owl *Otus lettia* in Thailand

Authors: Leadprathom, Kumron, Chimchome, Vijak, and Bumrungsri, Sara

Source: *Ardea*, 97(4) : 457-461

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.097.0409>

BioOne Complete ([complete.BioOne.org](https://complete.bioone.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Nesting ecology of the Collared Scops Owl *Otus lettia* in Thailand

Kumron Leadprathom^{1,*}, Vijak Chimchome¹ & Sara Bumrungsri²

Leadprathom K., Chimchome V. & Bumrungsri S. 2009. Nesting ecology of the Collared Scops Owl *Otus lettia* in Thailand. In: Johnson D.H., Van Nieuwenhuyse D. & Duncan J.R. (eds) Proc. Fourth World Owl Conf. Oct–Nov 2007, Groningen, The Netherlands. Ardea 97(4): 457–461.


Information on the Collared Scops Owl *Otus lettia* in Thailand is extremely limited and this study examined its nesting ecology. From August 2006 – September 2007, 23 nests were found and observed in Chanthaburi Province. The breeding season for the Collared Scops Owl started in early January and continued until the end of April. The majority of nest sites were a tree cavity with the opening facing skyward. The average nest height was 3.4 m (SD 1.7) from the ground; average size of the nest entrance was 16 (SD 6.1) cm x 28 (SD 17.4) cm; cavity dimensions were 17 (SD 6.1) cm wide, 24 (SD 8.9) cm long, and depth was 24 (SD 28.5) cm. The average crown cover in the nest area was 43%. The majority of the land used around the nest site was agricultural land. The number of eggs laid per nest was 1–4, with a majority of the nests containing three eggs, and the hatching rate was 60%. The average weight of hatched chicks was 13 (SD 1.6) g and the fledging weight was 88 (SD 7.6) g. The nesting success was 52% (12 of 23 nests), the survival rate (to fledging) was 65% and mortality rate was inversely related to chick age.

Key words: Collared Scops Owl, *Otus lettia*, nest, eggs, nesting season, chick survival, fledging success, Thailand

¹Faculty of Forestry Kasetsart University, Bangkok, Thailand;

²Faculty of Science, Prince of Songkhla University, Songkhla, Thailand;

*corresponding author (kumron57@gmail.com)



INTRODUCTION

The Collared Scops Owl *Otus lettia* is a very common resident in Thailand (Lekagul & Round 1991), but information on its ecology is extremely limited. It is a small owl (19–25 cm), distributed in East Asia (India, China and Nepal) and southeast Asia (Albania, Ceylon, Malaysia, Indonesia, Hainan, Taiwan, Great Sunda Islands, Philippines and Thailand) (Howard & Moore 1980, Hekstra 1984, Khobkhet 1999). The Collared Scops Owl is a nocturnal bird that is seldom observed in daytime. Its diet is mainly insects and other small animals, such as beetles, grasshoppers, lizards, earthworms, mice, bats, small birds and frogs. Their preferred habitats are dry evergreen forest, evergreen forest, hill evergreen forest, grassland and fruit orchard. The breeding season is between February and April. Typically, three to five eggs are laid in natural hollows or woodpecker holes in a tree trunk or dead stump and

most nesting sites are 2–5 m in height. Incubation starts with the first egg laid, and chicks in the same nest always show disparity in size (Khobkhet 1999). The young are semi-altricial at hatching, with their eyes closed and their bodies covered with natal down (Loahajinda 1985). This study describes the nesting behaviour and other basic information of this owl.

METHODS

Study area

Chanthaburi Province was our study area as this area is covered in a variety of land use types, such as fruit orchards, Para rubber plantation and patches of forest. Chanthaburi Province is 6338 km² in size and is located in the eastern part of Thailand (Fig. 1), c. 245 km from Bangkok. It has a tropical climate influenced by the southwest monsoon. The amount of rainfall in 2005



Figure 1. Location of Chanthaburi Province in Thailand.

was 2870 mm, the rainy season runs from May to September, and rainfall decreases from October to April (Meteorological Department 2006). The primary land-use in Chanthaburi Province is agriculture, followed by forests, water sources, urbanization and housing. Orchard farming is the most common agricultural activity in the province (Land Development Department 2004: OAE, 1995–2002).

A poster program was utilized to help us communicate with local citizens to find nests and roosts of Collared Scops Owls. The posters were distributed to communities, schools, local administrators and displayed at community stores between January and June 2007. All nests found were located by GPS and mapped. After we found a nest, the nest tree species, nest height and nest dimensions (width, length, and depth) were recorded. Plant composition and structure in a circular vegetation plot of 10-m radius around the nest was also determined. The crown cover of trees (diameter at breast height: DBH > 4.5 cm) in each plot was estimated using the formula for an ellipse: area = $0.25\pi \times d1 \times d2$, where d1 is the largest crown diameter and d2 is the diameter perpendicular to d1. We characterized land use around nests sites, using plot radii of

200 m, 305 m, and 1000 m. The first two plot sizes were based on the extent of the core area and home range of the Eurasian Scops Owls *Otus scops* in altered and semi-arid landscapes, respectively (Martinez *et al.* 2007). The 1000-m radius plot was based on 50% of the home range of the Spotted Owl *Strix occidentalis* in the southern Oregon Coast Ranges (Carey *et al.* 1990). Land use within these three plots was evaluated using the Arc GIS program, based upon the land-use map from the Land Development Department (2004).

Adult, nesting owls were trapped, weighed and measurements taken of the tarsus, bill and wingspan. Adults and chicks were banded with unique combinations of coloured plastic rings. Egg dimensions were measured and the chicks were marked when hatched on their talon. After hatching, each chick was measured (tarsus, bill and wingspan) and weighed every 2–3 days. After the chicks fledged, they were identified by their rings. Measurements are given as mean \pm SD unless otherwise noted.

Point Biserial Correlations (Pearson correlation coefficients) between crown cover, land-use type and nest characteristics was used to assess nesting success. Regression analysis and stepwise regression were calculated to derive the equations necessary to predict chick age.

RESULTS

Nest site characteristics

Twenty-three owl nests were found between early February and the end of April 2007, in agricultural areas. Eighteen of the nests were in trees and five were on the ground (Table 1). Nest dimensions are given in Table 2. Most owl nests (74%) opened on top (skyward).

Table 1. The location and characteristics of 23 Collared Scops Owl nests in Chanthaburi Province, Thailand.

Nest Location	Nest characteristics	Number of nests	
In trees	In dead tree stumps	2	
	In live trees	In the tree trunk	8
		On epiphytic ferns	4
		On dead branches	3
		In the tree fork	1
On the ground	In dead stumps	2	
	In cement tubes	2	
	Under tree fork	1	

The average crown cover at the nest location was $136.1 \pm 40.1 \text{ m}^2$ or 43% of the total 10-m radius area around the owl's nests, while the average crown cover above the nest was $132.0 \pm 40.3 \text{ m}^2$ or 42% of total area. The average nest tree's crown cover was $29.7 \pm 17.6 \text{ m}^2$, or 9% of total area. No correlation was found between the percentage of crown cover and nest success ($P > 0.05$).

Within a 200-m radius of the owl's nest, 93.9% of the area consisted of agricultural land, 3.4% of water sources and 2.8% of urbanization and housing (Table 3). Proportional land use was similar at distances of 305 m and 1000 m. There was no correlation between the percentage of each type of land use (agricultural area, the water sources, urbanization and housing area and forested area) and nest success ($P > 0.05$).

Eggs and breeding biology

The average dimensions of the eggs were $28.85 \pm 1.10 \times 33.65 \pm 1.39 \text{ mm}$ ($n = 41$). Clutch sizes ranged from 1–4 eggs, with the majority of nests having 3 eggs (63%, $n = 23$). The hatching rate was 60% (27 of 45 eggs). The incubation period was 22–29 days ($n = 3$) and the nestling period was 18–26 days ($n = 6$). The interval time between the first attempt that failed with chicks (at 7–8 days old) and the next attempt was 15 days ($n = 1$). Breeding pairs left their chicks alone dur-

ing the daytime after they were 16.0 ± 4.1 days old. The average body weight, and bill, tarsus and wingspan length of 12 adults (sex unknown) were $141.3 \pm 17.8 \text{ g}$, $15.0 \pm 1.8 \text{ mm}$, $32.5 \pm 2.1 \text{ mm}$ and $55.8 \pm 2.4 \text{ cm}$, respectively.

Chick growth and development

Newly hatched chicks were semi-altricial, their eyes were still closed with natal down, and the egg tooth was still apparent. Details of chick development are as follows: *Age 6–10 days*, the remiges and semi-plumes began to grow. The eyes opened on average at 6 ± 2.8 days and the egg tooth disappeared on average by 8.5 ± 2.4 days. *Age 11–15 days*, the bristles around the bill began to grow and were obvious by 20 days old. *Age 16–20 days*, the rectrices began to grow, while the remings and semi-plumes had become fully developed. During this period some of the chicks fledged. *Age 21–25 days*, fully feathered and the chicks fledged from the nest.

The overall daily growth rate for body weight, bill, tarsus, wingspan was 3.0 g, 0.2 mm, 0.7 mm and 1.2 cm, respectively. The highest daily rate of increase in body weight, 4.5 g/d, was when the chick was 6–7 days old. The chick attained maximum body weight when 24–25 days old, but the body weight had decreased by 1–2 g at 1–2 day before fledging. The highest daily rate

Table 2. The dimensions of Collared Scops Owl nest trees and hollows in Chanthaburi Province, Thailand.

	Nest tree ($n = 18$)			Nest hollow ($n = 23$)				
	DBH ^a (cm)	Height (m)	Nest height (m)	Entrance		Cavity		
				Width (cm)	Length (cm)	Width (cm)	Length (cm)	Depth (cm)
Mean \pm SD	41.9 ± 14.5	9.3 ± 5.3	3.4 ± 1.7	16.4 ± 6.1	28.3 ± 17.4	17.5 ± 6.1	23.7 ± 8.9	24.3 ± 28.5
Range	25.1–83.7	1.4–21.0	1.2–6.1	6.0–34.0	14.0–83.0	11.0–35.0	13.0–51.0	0–137.0

^aDiameter at breast height.

Table 3. The percentage of the land use types within different areas around Collared Scops Owl nests in Chanthaburi Province, Thailand.

Area around nest (m)	Percentage of the land use types					
	Field or mixed field crop ^b	Forest	River, canal, reservoir, lake, pond ^a	Urbanization and housing area	Mixed orchard, Para rubber ^b	Transplanted paddy field ^b
200	6.66	0.00	3.36	2.77	87.21	0.00
305	8.52	0.00	2.93	2.56	85.99	0.00
1000	13.12	1.58	2.55	3.69	78.18	0.88

^aWater sources. ^bAgricultural area.

of increase in bill length was 0.4 mm/d when 3–6 days old. The highest daily rate of increase in tarsus length was 1.2 mm/d when 1–8 days old. The highest daily rate of increase in wingspan was 1.81 cm/d when 12–13 days old. All rate of increase, bill, tarsus and wingspan, declined in that order from hatching to fledging. Comparative data on the averages of body weight, and bill, tarsus and wingspan length among hatchlings, fledglings and their parents (adults) are shown in Table 4.

By using stepwise regression analysis we predicted the age of the owl chick between hatching and fledging using measurements, and we found that the wingspan was suitable for predicting the age of owl chicks in this period:

$$\text{Age (day)} = [0.635 \times \text{Wingspan (cm)}] - 2.490$$

(Adjusted $R^2 = 0.989$, $P < 0.05$)

Breeding success

Breeding success, where the outcome was known, is as follows: the hatching success was 60% (27 of 45 eggs); fledging rate was 65.2% (30 of 46 young). Overall nest success (at least one young surviving to fledging) was 52.2% (12 of 23 nesting attempts). A total of 30 chicks fledged from 12 nests (2.5 ± 0.9 fledgling per successful nest), with 54.5% of pairs rearing one or more young (12 of 23 owl nests, 22 pairs, one pair used a different nest after the first attempt failed). Chick mortality rate was inversely related to chick age: when chicks were 5 days old, mortality rate was 38% and declined to 31% at 6–10 days old, 25% at 11–15 days old, and 6% at 16–20 days old.

DISCUSSION

In Chantaburi Province, Collared Scops Owl laid their eggs in the dry season between January and early March. The chicks hatched from mid-February to late-March and fledged from mid-March to late-April, which was by the beginning of the wet season. The breeding

season of Collared Scops Owl in this study (late-January to late-April) was similar to that of other Scops Owls in southeast Asia, starting in the late dry season and ending in the rainy season, such as Andaman Scops Owl *Otus bali* (mid-February – mid-April), Sunda Scops Owl *Otus lempiji* (February–April), and Philippine Scops Owl *Otus megalotis* (January–May) (König *et al.* 1999).

According to the egg and clutch sizes in this study, the egg dimensions of the Collared Scops Owl were about the same size as eggs of the Sunda Scops Owl (28.8 × 33.5 mm) but bigger than those of the Indian Scops Owl (27.0 × 33.0 mm). The clutch size was similar to the Indian Scops Owl clutch (3–4 eggs), but larger than Sunda Scops Owl clutch (2 eggs, rarely 3; König *et al.* 1999). Regarding clutch size and breeding season of Collared Scops Owl, Hekstra (1984) indicated that clutch size varied from north to south across its range, laying four or five eggs in the north in the spring but only three or four eggs in the south where it breeds during the period of maximum rainfall.

Although most owl nests were located in agricultural areas (fruit orchards and Para rubber plantations), it should not be concluded that Collared Scops Owls actually prefer to nest in these areas. The methods of nest searching and nest location used public relations, and as most people were active around their own agricultural land, most nests were found in those areas. Future studies should emphasize nest searchers in forested areas, such as Community Forests, National Parks, Forest Parks and Wildlife Sanctuaries.

In Australia, suitable tree hollows occurred most often in the tree canopy and least often in the main trunk (Harper *et al.* 2004), and in dead wood more than in live wood. In another Australian study by Whitford (2002), owls preferred hollows in the tree trunk rather than nesting on branches. Interestingly, in our study, the Collared Scops Owl used epiphytes, the ground and cement tubes as nest cavities, which suggests that nest sites were in short supply and that the provision of nest boxes would likely improve conditions for the owl. Most of the nest sites in this study were

Table 4. Average body weight, and bill, tarsus and wingspan among Collared Scops Owl hatchlings, fledglings and adults (parents). Mean \pm SD and sample size (in parentheses) are given.

Stage	Weight (g)	Bill (mm)	Tarsus (mm)	Wingspan (cm)
Hatchling	12.8 \pm 1.6 (18)	6.3 \pm 0.5 (14)	11.4 \pm 0.9 (14)	7.7 \pm 0.5 (14)
Fledgling	88.4 \pm 7.6 (32)	12.2 \pm 0.7 (32)	30.1 \pm 2.2 (32)	38.8 \pm 3.8 (32)
Adult	141.3 \pm 17.8 (12)	15.0 \pm 1.8 (10)	32.5 \pm 2.1 (10)	55.8 \pm 2.4 (10)

open on top, allowing rainwater to easily flood the nest, with nest drainage likely a significant factor in chick survival. We also found that one nest cavity of Collared Scops Owl was used by Asian Barred Owlets *Glaucidium cuculoides* after the Collared Scops Owls had left, indicating that two species of owl can share the same nest and habitat but at different times.

Conservation-oriented research is of paramount importance in Thailand. Many people still think that the owl is an evil bird. Most Thai people believe that when they hear or see an owl it is a sign of bad news. Because of this belief, most people try to eradicate owls from their land by shooting, stealing eggs, using pesticide to kill owls, or trapping owls near the nest. In addition, changing land use is another significant threat, such as destruction of owl habitat by urbanization or deforestation.

ACKNOWLEDGEMENTS

This work was supported by the TRF/BIOTEC Special Program for Biodiversity Research and Training grant BRT T_349010. We would like to thank the Forest Research Center, Kasetsart University, Thailand and World Owl Conference 2007 committee for financial supporting to attend at the World Owl Conference 2007 and thank to Dr. Alan Kemp for his review and valuable suggestions.

REFERENCES

- Carey A.B., Reid J.A. & Horton S.P. 1990. Spotted Owl Home Range and Habitat use in Southern Oregon Coast Ranges. *J. Wildl. Manage.* 54: 11–17.
- Harper J.M., McCarthy M.A., van der Ree R. & Fox J.C. 2004. Overcoming bias in ground-based surveys of hollow-bearing trees using double-sampling. *For. Ecol. Manage.* 190: 291–300.
- Hekstra G.P. 1984. Scops and Screech Owls. In: Burton J.A. (ed.) *Owls of the world*. Eurobook Limited, UK, pp. 86–107.

- Howard R. & Moore A. 1980. *A Complete checklist of the birds of the world*. Chaucer Press, Great Britain.
- Khobkhet O. 1999. *Birds of Thailand II*. Sarakadee, Bangkok. (In Thai)
- König, C., Weick F. & Becking J.H. 1999. *Owls: A guide to the owls of the world*. Pica Press, Sussex.
- Land Development Department. 2004. Supporting system to plan on economics crop AgZone 2.2. Land Development Department (Computer file), Bangkok, Thailand.
- Lekagul B. & Round D. 1991. *A guide to the birds of Thailand*. Saha Karn Bhaet, Bangkok.
- Loahajinda V. 1985. *Ornithology I*. Burapasasn, Bangkok. (In Thai)
- Martinez J.A., Zuberogoitia I., Martinez J.E., Zabala J. & Calvo J.F. 2007. Patterns of territory settlement by Eurasian scops-owls (*Otus scops*) in altered semi-arid landscapes. *J. Arid Environ.* 69: 400–409.
- Meteorological Department. 2006. Monthly statistic of rainfall and temperature. Meteorological Department, Bangkok. (In Thai)
- Whitford K.R. 2002. Hollows in jarrah (*Eucalyptus marginata*) and marri (*Corymbia calophylla*) trees I. Hollow sizes, tree attributes and ages. *For. Ecol. Manage.* 160: 201–214.

SAMENVATTING

In dit onderzoek in de provincie Chanthaburi, Thailand, werd informatie verzameld over de broedbiologie van de Gekraagde Dwergooruil *Otus lettia*. Tussen augustus 2006 en september 2007 werden 23 nesten van deze soort gevonden. De nesten lagen overwegend in landbouwgebieden. De bedekking door boomkruinen rond het nest was gemiddeld 43%. Het broedseizoen begon vroeg in januari en duurde tot eind april. De meeste nesten waren in een boomholte gemaakt met de opening naar boven gericht. De nesten waren op gemiddeld 3,4 m hoogte gebouwd. De nestopening was gemiddeld 16 × 28 cm groot, de nestholte gemiddeld 17 cm hoog en 24 cm breed en diep. De legselgrootte was 1–4 en de meeste nesten bevatten 3 eieren. Van de eieren kwam 60% uit. Bij het uitkomen wogen de jongen gemiddeld 13 g, bij het uitvliegen gemiddeld 88 g. Het nestsucces was 52%. De overleving van de jongen in de nestfase was gemiddeld 65%, waarbij de overlevingskans met de leeftijd toenam.

ARDEA

TIJDSCHRIFT DER NEDERLANDSE ORNITHOLOGISCHE UNIE (NOU)

ARDEA is the scientific journal of the Netherlands Ornithologists' Union (NOU), published bi-annually in spring and autumn. Next to the regular issues, special issues are produced frequently. The NOU was founded in 1901 as a non-profit ornithological society, composed of persons interested in field ornithology, ecology and biology of birds. All members of the NOU receive ARDEA and LIMOSA and are invited to attend scientific meetings held two or three times per year.

NETHERLANDS ORNITHOLOGISTS' UNION (NOU)

Chairman – J.M. Tinbergen, Animal Ecology Group, University of Groningen, P.O. Box 14, 9750 AA Haren, The Netherlands

Secretary – P.J. van den Hout, Royal Netherlands Institute for Sea Research (NIOZ), P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands (hout@nioz.nl)

Treasurer – E.C. Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl)

Further board members – E. Boerma, G.J. Gerritsen, J. Komdeur, J. Ouweland, G.L. Ouweneel, J.J. de Vries

Membership NOU – The 2010 membership fee for persons with a postal address in The Netherlands is €42 (or €25 for persons <25 years old at the end of the year). Family members (€9 per year) do not receive journals. Foreign membership amounts to €54 (Europe), or €65 (rest of the world). Payments to ING-bank account 285522 in the name of Nederlandse Ornithologische Unie, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (BIC: INGBNL2A and IBAN: NL36INGB0000285522). Payment by creditcard is possible. Correspondence concerning membership, payment alternatives and change of address should be sent to: Erwin de Visser, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (nou ledenadmin@gmail.com).

Research grants – The NOU supports ornithological research and scientific publications through its Huib Kluijver Fund and the 'Stichting Vogeltrekstation'. Applications for grants can be addressed to the NOU Secretary. Donations to either fund are welcomed by the NOU treasurer.

Internet – www.nou.nu

ARDEA

Editors of ARDEA – Rob G. Bijlsma, Wapse (Editor in chief); Christiaan Both, Groningen; Niels J. Dingemans, Groningen; Dik Heg, Bern; Ken Kraaijeveld, Leiden; Kees van Oers, Heteren; Jouke Prop, Ezinge (Technical editor); Julia Stahl, Oldenburg; B. Irene Tieleman, Groningen; Yvonne I. Verkuil, Groningen

Dissertation reviews – Popko Wiersma, Groningen

Editorial address – Jouke Prop, Allersmaweg 56, 9891 TD Ezinge, The Netherlands (ardea.nou@planet.nl)

Internet – www.ardeajournal.nl. The website offers free downloads of all papers published in Ardea and forerunners from 1904 onwards. The most recent publications are available only to subscribers to Ardea and members of the NOU.

Subscription ARDEA – Separate subscription to ARDEA is possible. The 2010 subscription rates are €36 (The Netherlands), €42 (Europe), and €50 (rest of the world). Institutional subscription rates are €53, €69, and €78, respectively). Papers that were published more than five years ago can be freely downloaded as pdf by anyone through ARDEA's website. More recent papers are available only to members of the NOU and subscribers of ARDEA-online. Receiving a hard-copy with additional access to ARDEA-online costs €55 (The Netherlands and Europe), €70 (rest of the world), or €110 (institutions). Subscriptions to ARDEA-online (without receiving a hard copy) cost €40 (individuals worldwide), or €85 (institutions). Payments to ING-bank account 125347, in the name of Nederlandse Ornithologische Unie, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (BIC: INGBNL2A and IBAN: NL16INGB0000125347). Correspondence concerning subscription, change of address, and orders for back volumes to: Ekko Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl).

World Owl Conference Special

Editors – David H. Johnson, Dries Van Nieuwenhuysse and James R. Duncan, in cooperation with Jouke Prop and Rob G. Bijlsma

Technical editor – Jouke Prop

Dutch summaries – Arie L. Spaans, Dries Van Nieuwenhuysse, Jouke Prop, Rob G. Bijlsma, or authors

Graphs and layout – Dick Visser

Drawings – Jos Zwarts

Cover photos - Serge Sorbi

front – Snowy Owl

back – Snowy Owl, Great Grey Owl and young Tengmalm's Owl

Production – Hein Bloem, Johan de Jong and Arnold van den Burg

© Nederlandse Ornithologische Unie (NOU), 2009

Printed by Van Denderen, Groningen, The Netherlands, December 2009