

Studying Earthworms (Annelida: Oligochaeta) in South Africa

Authors: Nxele, T.C., Lamani, S., Measey, G.J., Armstrong, A.J., Plisko, J.D., et al.

Source: African Invertebrates, 56(3) : 779-806

Published By: KwaZulu-Natal Museum

URL: <https://doi.org/10.5733/afin.056.0319>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Studying earthworms (Annelida: Oligochaeta) in South Africa

T.C. Nxele^{1*}, S. Lamani^{2,3}, G.J. Measey², A.J. Armstrong⁴, J.D. Plisko^{1,5},
S. Willows-Munro⁵, C. Janion-Scheepers^{2,3,6} and J.R.U. Wilson^{2,3}

¹KwaZulu-Natal Museum, Private Bag 9070, Pietermaritzburg, 3200 South Africa;
tnxele@nmsa.org.za; danutaplisko@gmail.com

²Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private Bag X1, Matieland, 7602 South Africa; cjanion@gmail.com; john.wilson2@gmail.com;
john@measey.com; slamani@sun.ac.za

³South African National Biodiversity Institute, Kirstenbosch Research Centre,
Claremont, 7735 South Africa;

⁴Ezemvelo KZN Wildlife, P.O. Box 13053, Cascades, 3202 South Africa;
Adrian.Armstrong@kznwildlife.com

⁵University of KwaZulu-Natal, School of Life Sciences, P.O. Box X01, Scottsville, 3209 South Africa;
willows-munro@ukzn.ac.za

⁶School of Biological Sciences, Monash University, Victoria, 3800 Australia;

*Corresponding author

ABSTRACT

Earthworms are an important component of southern African invertebrate diversity, due both to their influential roles in soil ecosystems, and the relatively large number of species. As of 2010, there were 282 indigenous earthworm species (most endemic) known to South Africa belonging to three families: Microchaetidae, Tritogeniidae and Acanthodrilidae. In addition, 44 introduced species from six families had been recorded. However, earthworms are rarely included in environmental monitoring or conservation programmes—partly because sampling and species identification are difficult and many sampling methods are destructive and/or toxic. In this paper we review the earthworm sampling techniques most commonly used by screening data from a digitised literature collection on South African earthworms and on-line global searches. By examining a case study sampling of three vegetation types, this paper highlights taxonomic challenges and the effort required to properly curate specimens. The study provides recommendations for future sampling and highlights some key priorities for future work on the group.

From the literature review in early 2012, it is clear that collection techniques are often insufficiently recorded in published work. A total of 10 938 publications from the period 1950 to 2012 were found from the literature search and digitised collection and from these only 32 papers recorded the sampling methodology (mainly hand sorting) for South African research, pointing to the need to adopt standard sampling and reporting protocols. We also tested two of the most popular methodologies in the field. Sampling was conducted in January and February 2012 at four sites, with 24 plots at each site (12 digging and 12 using mustard extraction). A total of 2 094 earthworms collected could be assigned a species name, with introduced species predominating at both disturbed and natural sites. It took a team of three to five people digging and hand collecting all earthworm specimens encountered in a plot of 50 cm × 50 cm × 20 cm deep around 45 to 60 minutes. However, much more time was spent curating and identifying samples. While we recommend following the ISO (ISO11268-3, ISO23611-1) protocol for collecting introduced taxa, to get a complete inventory of South African earthworms a range of sampling techniques will be required; in particular, a large 1 m × 1 m × 20 cm plot is required for many large bodied native taxa, and the collection of giant earthworms will require different approaches. The identification of specimens requires skills that are scarce in the country and so there is an urgent need for training and funding for fundamental work on earthworm taxonomy. An atlas project could serve as a focal point for future research. In providing some general recommendations based on the long and fruitful history of research on earthworms in South Africa, we are optimistic that a better understanding of the group will help us to both improve our usage of natural resources and provide insights into this vitally important edaphic group.

KEYWORDS: Afrotropical Region, South Africa, megadrile, earthworms, sampling techniques, biodiversity, sampling effort, taxonomy, literature database.

INTRODUCTION

Pressures such as climate change, pollution and urbanisation mean that biodiversity research is becoming increasingly important if we are to manage ecosystem functioning

<http://africaninvertebrates.org>

urn:lsid:zoobank.org:pub:0658A5B4-8B5B-4455-A412-E66D770C87CB

and services such as soil fertility, clean water and waste processing (Naeem *et al.* 1994; Slotow & Hamer 2000; Crouch & Smith 2011; Hamer 2010). However, global change drivers and their synergistic effects (Millennium Ecosystem Assessment 2005) might cause extinctions of species not even known to science (Essl *et al.* 2013; Costello *et al.* 2013). One aspect that has often been overlooked is the significant contribution of soil fauna to soil health, below ground diversity, and more broadly to ecosystem functioning (Wardle 2002, Wardle *et al.* 2004, Louw *et al.* 2014).

South Africa contains one of the most diverse temperate faunas in the world (Steenkamp & Smith 2006), and has a strong history of biodiversity research (Crouch & Smith 2011). Despite this, knowledge of the soil fauna of South Africa is scattered and the fauna poorly understood (Janion-Scheepers *et al.* in prep.). Most research has focused on taxonomically well-known groups such as ants (Parr *et al.* 2005, Botes *et al.* 2006, Braschler *et al.* 2012), beetles (Davis 1997, Davis *et al.* 1999, Botes *et al.* 2007) and spiders (Dippenaar-Schoeman & Craemer 2000), with only a few recent efforts on groups such as Collembola (Janion *et al.* 2011, 2015), mites (Ermilov & Hugo-Coetzee 2012) and nematodes (Borgonie *et al.* 2011, Knoetze *et al.* 2006). Lack of current research is certainly driven by the logistics of sampling soil fauna. Sampling is often difficult and time consuming. Moreover, the lack of standardisation in sampling protocols has hampered comparisons of diversity between soil fauna studies.

The important role of earthworms as soil processors and ecosystem engineers has been widely demonstrated (Edwards & Bohlen 1996; Lavelle *et al.* 2006), and the role they play in soil fertility and health was even recognised by Darwin (1881). The activities of earthworms in the soil are vital for a healthy, fully functional system (Butt & Grigoropoulou 2010). These activities can substantially change the physical and chemical characteristics of the soil environment either directly or indirectly, with consequences for the entire soil food web, nutrient distribution and even vertebrate and understory plant communities (Edwards & Bohlen 1996). However, many ecological studies group all earthworms together, despite important functional differences between taxa reflecting a great diversity and wide range of adaptations to environmental conditions (Coleman *et al.* 2004). If the role of earthworms in ecosystems is to be quantified, a precise and accurate estimation of their diversity, abundance and biomass is needed (Valckx *et al.* 2011).

Earthworm sampling methods can be broadly divided into active, ethological (behavioural) and passive collection methods (Bartlett *et al.* 2010; Valckx *et al.* 2011). The main method of active sampling is by digging up a portion of soil (ideally of known volume) and sorting through the soil by hand to collect all earthworms and cocoons. Alternatively, earthworms can be expelled from the soil due to their behavioural response to certain stimuli by a range of chemical expellants or vermifuges (Coleman *et al.* 2004). It is also common to use both digging and hand sorting as well as chemical expellants (Baretta *et al.* 2007). Finally, indirect methods involve looking for signs of earthworms or waiting until they emerge on the surface (e.g. dispersal of some species after heavy rains).

Different chemical expellants or irritants can expel earthworms from the soil. Commonly used expellants are formaldehyde (Raw 1959; ISO 11268-3, 1999; Eichinger *et al.* 2007), commercial hot mustard (Gunn 1992; East & Knight 1998; Chan & Munro 2001; Lawrence & Bowers 2002; Eisenhauer *et al.* 2008), allyl isothiocyanate (AITC) (Zarborski 2003), household detergents such as washing-up liquid (East & Knight 1998) and potassium permanganate (Evans & Guild 1947, Reinecke & Ryke 1972). Other

ethological methods used are the octet or electrical method (Čoja *et al.* 2008), heat extraction (Čoja *et al.* 2008), mechanical vibration and the use of pitfall traps (Callaham *et al.* 2003). Not all of these methods are DNA compatible and certain collection techniques can limit downstream molecular projects. When sampling earthworms formalin is often used as an expellant. However, it is not possible to recover high-quality DNA from tissue that has been exposed to formalin (Moelans *et al.* 2011), and as such the use of formalin is discouraged when tissues are to be used in molecular studies. Several authors have suggested that mustard be used as an alternative to formalin as an expellant (Gunn 1992; East & Knight 1998; Chan & Munro 2001; Lawrence & Bowers 2002; Eisenhauer *et al.* 2008; Pelosi *et al.* 2009; Valckx *et al.* 2011). Gunn (1992) found that the use of mustard was more successful than formalin, while Lawrence and Bowers (2002) found that mustard was environmentally friendly and a more efficient alternative to formalin across soil and habitat types. The use of mustard powder or prepared mustard presents some problems for standardising earthworm sampling, as the composition of the mustard, how it is applied, and the local soil conditions can vary (Zarborski 2003).

The International Organisation of Standardisation (ISO) set a standard for the sampling of soil invertebrates to address the need to standardise the sampling of terrestrial soil invertebrates (ISO 23611-1, 2006). In part 1 of the ISO document a standard is set for the hand sorting and formalin extraction of earthworms (ISO 23611-1, 2006). The isolation and hand sorting of earthworms in a soil sample of a certain area (0.25 m²) or volume (25 × 25 × 20 cm) is suggested. This method is unfortunately a very laborious procedure and is difficult when the soil has high clay content or if there is a dense root mat present, and when physical disturbance of the study site is not acceptable, meaning digging is not an option (Bouche & Garner 1984; Gunn 1992; Eisenhauer *et al.* 2007).

Although digging is the most reliable method, a true and complete account of earthworm abundance will require a range of sampling techniques (Lawrence & Bowers 2002), as the relative efficiency of different sampling methods can vary with site characteristics, season and earthworm species (Callaham & Hendrix 1997). Indeed, most studies encourage the use of more than one sampling method (Coleman *et al.* 2004). By using a combination of active, behavioural, and passive techniques (e.g. digging, formalin extraction, and cast surveys), the accuracy of estimates of earthworm populations can be significantly increased (Bouche & Gardner 1984).

Ecologists and conservationists rely on taxonomists for information regarding species identifications (Chang *et al.* 2009). Earthworm identification is, however, not straightforward. The use of morphological characters is time consuming, labour intensive and requires trained specialists (Bartlett *et al.* 2010). A high level of cryptic diversity occurs (Rougerie *et al.* 2009), and the identification of juveniles and cocoons to species level cannot be made by traditional taxonomic means alone.

Earthworm research in South Africa has been conducted since the nineteenth century (Beddard 1895). As of 2010, there were 282 described earthworm species indigenous to South Africa belonging to three families: Microchaetidae, Tritogeniidae and Acanthodrilidae (specifically to the sub-family Acanthodrilinae) (Plisko 2010). However, large areas of South Africa remain to be surveyed, and most of the groups that have been well sampled require taxonomic revision.

The reference and research collection of earthworms housed in the KwaZulu-Natal Museum (Pietermaritzburg, KwaZulu-Natal) has 138 types (Plisko 2006, 2007, 2008). All types and most specimens were fixed in formalin and so they currently cannot be

used in molecular studies. In terms of introduced fauna, 50 species had been recorded by Plisko (2010) but the number has been corrected by Plisko & Nxele (2015) to 44 due to some of the species possibly being indigenous. Many of these species are widespread and abundant, with introduced species dominating disturbed agricultural sites (Ljungström 1972; Visser & Reinecke 1977; Plisko 2000; Dlamini *et al.* 2001; Haynes *et al.* 2003; Plisko 2010). While the extent to which introduced earthworms have colonised natural sites is poorly known, results from Dlinza Forest Nature Reserve in Eshowe and from a recent survey of Queen Elizabeth Park, in Pietermaritzburg, highlighted that even at less disturbed sites introduced taxa can be more common than native taxa (Plisko 2000; Nxele 2012). It remains to be seen whether introduced earthworms are having important economic or ecological impacts in South Africa, but introduced earthworms have had major detrimental effects in several locations around the world (Hendrix *et al.* 2008), particularly in areas where earthworms were not previously present (e.g. islands, or areas post-glaciation).

The aim of this paper is to review the literature on techniques commonly used for earthworm sampling, conduct some preliminary sampling with the two most widely used methods that are compatible with genetic techniques, provide guidelines and recommendations for sampling, and highlight future research focus areas.

MATERIAL AND METHODS

Review of sampling methods

The methods used to sample earthworms were investigated by reviewing sampling techniques globally and locally. A targeted literature search was conducted using ISI Web of Science™. The search was conducted in 2012 by firstly determining the total amount of available published data on “earthworms” from 1950 to 2012 by searching for the term “earthworm*” in the title, topic or abstract. Secondly, a targeted search was done to establish the number of available research papers discussing or reviewing different earthworm sampling techniques, using the following term: “(earthworm* OR oligochaeta*) AND (sampling method* OR hand sorting* OR formalin* OR allyl isothiocyanate* OR mustard*)”. Additional data from the publications were also collected to evaluate recent and past sampling techniques, the use of different terminology when describing surveys, the collection of biogeographical data, the preservation of samples, and the use of different taxonomic identification tools. A digital literature collection in the KwaZulu-Natal Museum was also searched (donated by A. J. Reinecke and J. D. Plisko). This collection contains articles of relevance on South African earthworms, with a focus on taxonomy, ecology, vermiculture, morphology, physiology, neurology, genetics and ecotoxicology. Some papers in the collection had already been recovered in the web search and therefore were not included; only the additional papers not recovered by the web search were added.

Pilot sampling

Earthworms were sampled using digging and hand sorting at four sites in KwaZulu-Natal, South Africa, during January and February 2012 (i.e. immediately following summer rains when the collection of earthworms is most effective (Plisko 2002a; Fig. 1; Table 1). Two of the sites were in natural vegetation in protected areas (a mistbelt forest and a grassland) and two in disturbed areas (National Botanical Gardens—



Fig. 1. Earthworm sampling. (A) The most common method is for plots to be dug up, the soil sorted by hand, and earthworms removed. (B) An alternative method is to apply chemical irritants (e.g. mustard solution) to a plot and the earthworms collected once they emerge from the soil. (C) Sorting, cleaning and processing earthworms in the field is a major component of sampling. Decisions as to the size of plot used, how deep the plots are dug, and which chemicals are used in which concentrations, as well as the timing of the sampling, all need to be adjusted depending on the goals of sampling. For large individuals (and anecic species) different sampling methods are needed (see text).

Pietermaritzburg, and a fallow agricultural field). At the agricultural site, sampling was conducted both in ploughed cultivated areas and on contour ridges between the cultivated areas (the latter had kikuyu grass, weeds and tall reeds). At each of the four study sites,

TABLE 1
Localities sampled during the pilot survey.

| Site | Habitat type | Management type | Lat / Long |
|--|-----------------------------------|--|-----------------------|
| Queen Elizabeth Park | Grassland | Protected nature reserve | 29.57148°S 30.32235°E |
| Cedara | Agriculture | Tilled, fallow, previously planted with soya beans | 29.54673°S 30.27778°E |
| Karkloof | Mistbelt forest | Protected nature reserve | 29.30393°S 30.22873°E |
| National Botanical Gardens — Pietermaritzburg | Planted gardens and compost heaps | Landscaped gardens | 29.60668°S 30.34832°E |

12 sampling plots of 50 cm × 50 cm (after Keith & Schmidt 2013) randomly spaced at intervals not closer than 5 m from each other were selected, and soil to a depth of 20 cm dug out using a metal frame as a guide. The soil was placed onto a plastic sheet and hand sorted for earthworms. Where possible, leaf litter was collected and also searched for earthworms prior to the digging.

In addition, 12 more plots of the same size were sampled using a chemical irritant extraction technique. Mustard powder was prepared according to the protocol from ISO 23611-1 (ISO 2006). One hundred grams of dry mustard powder (Colman's) was dissolved in 1 litre of water. In the field, the mixture was diluted with 9 litres of water in 10 litre watering cans, to give a concentration of 10 gL⁻¹. Leaf litter was removed from each plot and then the mixture poured onto the plot. The earthworms that appeared on top of the soil were collected for about 30 minutes.

Sampled earthworms were washed twice in clean tap water and afterwards narcotised in 20% ethanol. The narcotised earthworms were then weighed (wet mass) to the nearest milligram (mg) using a field balance (Highland portable precision range field balance). Earthworms were then preserved in absolute ethanol, clearly marked with labels and transported to the KZN Museum laboratory for identification. Earthworms were identified to species level using a Wild (Heerbrugg) microscope and descriptions and keys from the published literature (Ljungström 1972; Gates 1972; Plisko 1992; Csuzdi 2010; Blakemore 2011). The collected material was deposited in the KZN Museum.

Sample-based rarefaction curves were plotted and the ACE estimator employed to examine the degree of sampling efficiency and to determine the predicted number of species in each site, using the computer programme EstimateS V9.1.0 (Colwell 2013). All unidentified earthworms (juveniles, damaged adults and other specimens that could not be identified to species level) were removed from the dataset prior to rarefaction and estimation. Sampling was considered adequate if the estimator converged at the highest observed (identified) value (Longino *et al.* 2002).

Future directions for earthworm research in South Africa

To determine future priorities, we held a workshop on earthworm research in November 2011 and again in January 2015. Using various horizon scanning exercises in agriculture, ecology and environmental sciences as a background (e.g. Sutherland *et al.* 2013), we developed various key priorities for earthworm research.

RESULTS

Review of sampling methods

A total of 10 938 publications were found using the search keyword “earthworm*”. Of these papers, only 480 (4.4%) were directly linked to earthworm sampling by using the search keywords: “(earthworm* OR oligochaeta*) AND (sampling method* OR hand sorting* OR formalin* OR allyl isothiocyanate* OR mustard*)”. Two important issues were observed. First, a vast array of different concepts and terminologies were found to describe earthworm sampling. Different authors use different terms to describe the same procedure or concept, and to avoid confusion authors need to use well defined terms (De Zorzi *et al.* 2005). One method is to develop standard protocols (e.g. Supplementary Material 1). Second, when reviewing the sampling protocol it was found that although thorough descriptions of sampling techniques were given, preservation techniques, determination of biomass and taxonomic identification were poorly documented and often not described at all. The search of the database found 32 papers that described sampling methods for South African studies, of these 13 were already recovered from the web literature search. All the publications that contained South African data were recorded in Table 2. The publications containing data from outside South Africa were recorded in Table 3. Most South African studies used random digging and hand sorting in different habitats such as forests and grasslands.

Pilot sampling

Of the earthworms collected using the two sampling methods at the four sampling sites, a total of 2 094 earthworms were assigned to 16 species belonging to five families (Table 4). Some earthworms (196) could not be identified as they were unknown juveniles or damaged specimens and were not used in the results. The species accumulation curves (Supplementary material Fig. 2 A–D) indicate that most of the earthworm species at a site were sampled where species richness was relatively low. At sites where earthworm richness was higher, undersampling of species was indicated.

In the grassland of Queen Elizabeth Park a total of 217 earthworms were collected through digging and hand sorting; no earthworms were collected when mustard was used. Only three species (*Dichogaster* sp., *Amyntas corticis* (Kinberg, 1867) and *Amyntas rodericensis* (Grube, 1879)) were found, all introduced to South Africa. Previous studies in this park collected six species in the grassland, including the indigenous *Tritogenia howickiana* (Michaelsen, 1913) and nine species when including more vegetation types in the park (Nxele 2012). In Cedara, no earthworms were found from the plots that were in the middle of the cultivated field either by digging or mustard, but 85 earthworms from five different species (*Octolasion lacteum* (Örley, 1881), *Amyntas aeruginosus* Kinberg, 1867, *Amyntas corticis*, *Amyntas gracilis* (Kinberg, 1867) and *Amyntas minimus* (Horst, 1893)) were collected from plots that were in the grass contour between the fields. All the sampled earthworms were introduced species. In addition, unknown juveniles were also collected, but could not be assigned to any species.

When using the digging and hand sorting method in the Karkloof Forest a total of 766 earthworms and nine species were collected. Eight of these species (*Pontoscolex corethrurus* (Müller, 1857), *Aporrectodea rosea* (Savigny, 1826), *Dendrobaena octaedra* (Savigny, 1826), *Dendrobaena* sp., *Dendrodrilus* sp., *Amyntas minimus*, *Amyntas* sp., unidentified lumbricid species) were introduced species, and only one indigenous specimen (*Tritogenia* sp.) was found. When using the mustard method, 485

TABLE 2

Studies that sampled earthworms in South Africa (1895–2012) combined from the literature collection database in the KwaZulu-Natal Museum and literature review. N/A = no dominant species or number of species and ecological category were mentioned.

| Reference | Focus | Method | Location | Dominant species and ecological category |
|----------------------------------|---|--|--|---|
| Dlamini <i>et al.</i> 2001 | Quantitative & qualitative sampling of exotic earthworms on sugarcane estates | Digging and handsorting | Eshowe-Empangeni, KZN | <i>Pontoscolex corethrurus</i> ; Endogeic |
| Haynes <i>et al.</i> 2003 | Quantitative & qualitative sampling of earthworms in agricultural land in KZN | Digging and handsorting | KZN Midlands | <i>Amyntas diffringens</i> ; Endogeic |
| Horn <i>et al.</i> 2007 | Diversity of earthworms in forests in Limpopo | Searching and digging | Soutpansberg and Northern Drakensberg, Limpopo | <i>Amyntas diffringens</i> ; Endogeic |
| James & Davidson 2012 | Taxonomic studies | Digging and handsorting | KZN | N/A |
| Jamieson 1967 | Taxonomy | Digging and handsorting | Various sites | <i>Eukeria saltensis</i> ; Endogeic |
| Ljungström 1972 | Taxonomic studies | Digging and handsorting | Various sites | <i>Pontoscolex corethrurus</i> ; Endogeic |
| Michaelsen 1913 | Taxonomic studies | Digging and handsorting | KZN | N/A |
| Nxele 2012 | Taxonomy & Biodiversity assessment | Digging and handsorting | Queen Elizabeth Park, Pietermaritzburg, KZN | <i>Amyntas rodericensis</i> ; Epigeic |
| Pickford 1937 | Taxonomy | Digging | Various sites | <i>Eodriloides arundinis</i> ; Epigeic |
| Plisko 1991–1993, 1996a, b; 1997 | Taxonomic studies | Digging and handsorting | Various sites | N/A |
| Plisko 2000–2004, 2006a, b 2007 | Taxonomic studies | Digging and handsorting | Various sites | N/A |
| Reinecke & Ackerman 1977 | Taxonomic studies | Digging and handsorting | Northern Kruger National Park, Limpopo | N/A |
| Reinecke & Ljungström 1969 | Ecological studies - introduced species | Digging and handsorting | Potchefstroom | <i>Eisenia rosea</i> ; Endogeic |
| Reinecke & Ryke 1970 | Ecophysiological study of lumbricids & microchaetids | Digging and handsorting | Potchefstroom | <i>Proandricus modestus</i> ; Endogeic |
| Reinecke & Ryke 1972 | Ecology, sampling methods | Digging and handsorting, formalin, KMnO ₄ | Potchefstroom | <i>Proandricus modestus</i> ; Endogeic |

TABLE 2 (continued)

Studies that sampled earthworms in South Africa (1895–2012) combined from the literature collection database in the KwaZulu-Natal Museum and literature review. N/A = no dominant species or number of species and ecological category were mentioned.

| Reference | Focus | Method | Location | Dominant species and ecological category |
|------------------------|---|-------------------------|--|--|
| Reynolds 1993 | Taxonomic studies | Digging and handsorting | Swaziland | N/A |
| Uys 2006 | Biodiversity assessment | Digging and handsorting | South Africa: Drakensberg, KZN | N/A |
| Visser & Reinecke 1977 | Quantitative & qualitative sampling in irrigation area - Population densities | Digging and handsorting | South Africa: Potchefstroom, North West Province | <i>Eisenia rosea</i> ; Endogeic |
| Zicsi & Plisko 1991 | Taxonomy | Digging and handsorting | South Africa: KZN | N/A |

earthworms were collected from seven species (*Pontoscolex corethrus*, *Aporrectodea rosea*, *Dendrobaena octaedra*, *Dendrobaena* sp., *Amyntas minimus*, *Amyntas* sp., and unidentified lumbricid species) all of which were also collected using the digging and hand sorting method. When using the digging and hand sorting method in the Botanical gardens a total of 275 earthworms from 11 species were collected. When using the mustard method 266 earthworms from 12 species were collected, including *Tritogenia* sp (Table 4). The endogeic *P. corethrus* was dominant in QEP, the epigeic *A. rodericensis* was dominant in Cedara, in the Karkloof Forest the endogeic *O. lacteum* was collected in high numbers using both methods though digging collected almost double the number of individuals than the mustard, and the endogeic *A. minimus* was dominant in the Botanical Gardens.

The earthworms collected by digging and hand sorting weighed on average 0.358 g per worm (n=294), slightly heavier than the earthworms collected using mustard extraction (average of 0.294 g each, n=273).

Estimates of the sampling effort are shown in Table 5. Four people completed the pilot study in 10 days with eight working hours in each day. This time included travelling, setting up the field workstation, sampling and travelling with equipment from plot to plot, and packing up and cleaning after each day. As field assistants became more skilled over the course of the study, the time taken to complete a task decreased. When using the digging and hand sorting method it took an average of 40 minutes for one person to sample a plot (50 × 50 × 20 cm) with an average of 44 earthworms per plot. The time taken to collect earthworms with the mustard technique averaged 13 minutes per plot (50 × 50 cm) with an average of 35 earthworms collected per plot. Standardising on 100 earthworms, preserving the earthworms (including washing and narcotising them) and the determination of the biomass took an average of 26 and 42 minutes, respectively. The time taken to identify each earthworm varied greatly depending on (a) the experience of the taxonomist and (b) the identity of the worm. It took three months to complete the classification for all the earthworms in this study, an average of two hours per specimen.

TABLE 3

Sampling methods used in earthworm studies from other countries from the literature collection database in the KwaZulu-Natal Museum.

| Reference | Focus | Method | Location |
|--------------------------------|---------------------|--|--------------|
| Beyer <i>et al.</i> 1980 | Ecotoxicology | Digging and hand sorting | USA |
| Beyer & Cromartie 1987 | Ecotoxicology | Digging and hand sorting | USA |
| Boyd 1957 | Ecology | Active searching, KMnO ₄ , Digging and hand sorting | Scotland |
| Butt 1991 | Vermiculture | Formalin | UK |
| Butt <i>et al.</i> 2006 | Ecology | Digging and hand sorting | England |
| Chaudhuri & Bhattacharjee 1999 | Ecology | Digging and hand sorting | India |
| Cook <i>et al.</i> 1980 | Ecotoxicology | Formalin | Nigeria |
| Cuendet 1983 | Ecology | Formalin, Digging and hand sorting | Switzerland |
| Damoff & Reynolds, 2004 | Taxonomy | Digging and hand sorting | USA |
| Daniel <i>et al.</i> 1992 | Sampling techniques | Formalin, Chloroacetophenone, Digging and hand sorting | Switzerland |
| Edwards & Lofty 1982 | Vermiculture | Formalin | UK |
| Enckell & Rundgren 1988 | Ecology | Formalin | Faroe Island |
| Enckell <i>et al.</i> 1986 | Ecology | Formalin | Faroe Island |
| Evans & Guild 1947 | Ecology | Vermifuge | UK |
| Fragoso & Reynolds 1997 | Taxonomy | Digging and hand sorting, Active searching | Mexico |
| Guild 1951 | Ecology | KMnO ₄ | Scotland |
| Guild 1952a | Ecology | KMnO ₄ | UK |
| Guild 1952b | Ecology | KMnO ₄ | Scotland |
| Guild 1957 | Ecology | Digging and hand sorting | Scotland |
| Haimi & Boucelham 1991 | Ecology | Active searching | Finland |
| Hendriksen 1990 | Ecology | Formalin | Denmark |
| Hopp 1947 | Ecology | Soil corer | USA |
| Joannes & Kretzschmar 1983 | Physiology | Formalin | France |
| Laverack 1960 | Physiology | Digging and hand sorting | England |
| Laverack 1961 | Physiology | Formalin | UK |
| Martin 1980 | Ecology | Digging and hand sorting | New Zealand |
| Mischis <i>et al.</i> 1997 | Taxonomy | Digging and hand sorting, Active searching | Argentina |
| Moreno <i>et al.</i> 1982 | Taxonomy | Formalin, Digging | Spain |
| Murchie 1958 | Ecology | KMnO ₄ , Digging and hand sorting | USA |
| Nelson & Satchell 1962 | Sampling techniques | Digging and hand sorting | England |

TABLE 3 (*continued*)

Sampling methods used in earthworm studies from other countries from the literature collection database in KwaZulu-Natal Museum.

| Reference | Focus | Method | Location |
|-------------------------------|---------------------|---|----------------------|
| Nordström & Rundgren 1972 | Sampling techniques | Formalin, Formalin + detergent, KMnO ₄ , Digging and hand sorting | Sweden |
| Nordström & Rundgren 1973 | Ecological | Formalin | Sweden |
| Nuutinen & Haukka 1990 | Ecological | Formalin | Finland |
| Omedeo & Martinucci 1987 | Taxonomy | Digging and hand sorting, Active searching | Italy & Algeria |
| Pavliček <i>et al.</i> 1996 | Biodiversity | Formalin | Israel |
| Pop 1987 | Ecology | Formalin | Carpathians |
| Pop & Postolache 1987 | Ecology | Formalin | Romania |
| Raw 1962 | Ecology | Formalin | UK |
| Reeves & Reynolds 2004 | Taxonomy | Digging and hand sorting | Cayman Islands |
| Reynolds 1972 | Ecology | Digging and hand sorting | USA |
| Reynolds <i>et al.</i> 1995 | Taxonomy | Digging and hand sorting | Belize |
| Reynolds & Hanel 2005 | Taxonomy | Digging and hand sorting | South Atlantic ocean |
| Reynoldson 1955 | Ecology | KMnO ₄ , Digging and hand sorting | Wales |
| Reynoldson <i>et al.</i> 1955 | Ecology | Active searching | UK |
| Satchell 1970 | Ecology | KMnO ₄ , Formalin, Digging and hand sorting, Heat extraction, Floating | UK |
| Scheu 1987 | Ecology | Active searching | Canada |
| Scheu 1993 | Ecology | Active searching | Canada |
| Schwert 1977 | Taxonomy | Digging and hand sorting | Canada |
| Sims 1967 | Taxonomy | Formalin | Gambia |
| Tomlin & Gore 1974 | Ecotoxicology | Formalin | Canada |

Future directions for earthworm research in South Africa

The key priorities for research identified during the workshop on earthworms are summarised in Box 1. These included foundational biodiversity goals such as preparing an atlas, taxonomic keys, field guides and a DNA barcode library. Other goals included increasing outreach efforts and public interest (e.g. through a roadshow). The latter are considered important in terms of engaging with stakeholders that already have an interest in earthworms as well as providing exposure to taxonomic experts, thereby assisting with the potential enrolment of future taxonomists. There are many urgent needs, but it is clear that these goals must be underpinned by foundational biodiversity survey and description work, which in turn requires capacity building in taxonomy in South Africa.

TABLE 4
Species abundance for each sampled site.

| Family | Species | Status | Queen Elizabeth Park (natural grassland) | | Cedara (Agricultural field) | | Karkloof (natural forest) | | Botanic Gardens (transformed) | | Total |
|-----------------|--------------------------------|----------------------|--|------------|-----------------------------|------------|---------------------------|------------|-------------------------------|------------|-------------|
| | | | Digging | Mustard | Digging | Mustard | Digging | Mustard | Digging | Mustard | |
| Acanthodrilidae | <i>Acanthodrilidae</i> sp. | ? | 0 | | 0 | | 0 | 0 | 0 | 5 | 5 |
| Acanthodrilidae | <i>Dichogaster</i> sp. | Introduced | 4 | | 0 | | 0 | 0 | 2 | 0 | 6 |
| Pontoscolicidae | <i>Pontoscolex corethrurus</i> | Introduced; Endogeic | 212 | | 0 | | 0 | 0 | 2 | 0 | 214 |
| Lumbricidae | <i>Aporrectodea rosea</i> | Introduced; Endogeic | 0 | | 0 | | 48 | 7 | 4 | 1 | 60 |
| Lumbricidae | <i>Dendrobaena octaedra</i> | Introduced; Epigeic | 0 | | 0 | | 7 | 0 | 0 | 0 | 7 |
| Lumbricidae | <i>Dendrobaena</i> sp. | Introduced | 0 | | 0 | | 64 | 68 | 0 | 1 | 133 |
| Lumbricidae | <i>Dendrodrilus</i> sp. | Introduced | 0 | | 0 | | 1 | 0 | 0 | 0 | 1 |
| Lumbricidae | Lumbricidae sp. | Introduced | 0 | | 0 | | 159 | 143 | 7 | 19 | 328 |
| Lumbricidae | <i>Octolasion lacteum</i> | Introduced; Endogeic | 0 | | 0 | | 474 | 258 | 11 | 14 | 757 |
| Megascolecidae | <i>Amyntas aeruginosus</i> | Introduced; Epigeic | 0 | | 1 | | 0 | 0 | 30 | 15 | 46 |
| Megascolecidae | <i>A. corticis</i> | Introduced; Epigeic | 0 | | 22 | | 9 | 7 | 5 | 4 | 47 |
| Megascolecidae | <i>A. gracilis</i> | Introduced; Epigeic | 0 | | 2 | | 0 | 0 | 9 | 1 | 12 |
| Megascolecidae | <i>A. minimus</i> | Introduced; Endogeic | 0 | | 14 | | 0 | 0 | 100 | 74 | 188 |
| Megascolecidae | <i>A. rodericensis</i> | Introduced; Epigeic | 1 | | 46 | | 0 | 0 | 29 | 32 | 108 |
| Megascolecidae | <i>Amyntas</i> sp. | Introduced | 0 | | 0 | | 3 | 2 | 76 | 99 | 180 |
| Tritogenidae | <i>Tritogenia</i> sp. | Native; Endogeic | 0 | | 0 | | 1 | 0 | 0 | 1 | 2 |
| | TOTAL | | 217 | N/A | 85 | N/A | 766 | 485 | 275 | 266 | 2094 |

DISCUSSION

A wide variety of techniques are used for the sampling of earthworms in the field (Bouché & Gardner 1984, Druce *et al.* 2004; Čoja *et al.* 2008), reflecting in part the different research questions asked, but also because earthworm species have a variety of ecological strategies, patterns of behaviour and habitat preferences (Lee 1985; Blakemore 2002). Epigeic and endogeic species can often be collected by hand; whereas anecic species might require chemical irritants to expel the earthworms from the soil. In our sampling, epigeic earthworms dominated in Cedara and endogeic earthworms dominated in the Botanical Gardens. In addition, earthworm populations are usually spatially aggregated (Valckx *et al.* 2011). This complicates sampling efforts and attempts to estimate population sizes, particularly as the factors that determine this patchy spatial distribution are not yet sufficiently understood (Valckx *et al.* 2009). Thus, sampling techniques need to account for the spatial distribution of earthworms in order to get an accurate estimate of species richness and abundance. At the indigenous forest in Karkloof, a greater abundance of earthworms was obtained by digging in comparison to the mustard solution method, whereas almost equal abundances were obtained using the two methods in the transformed habitat of the Botanical Gardens. Despite the existence of international standards (ISO 23611-1 and ISO 11268-3), many studies are not comparable due to the lack of standardisation of sampling protocols. While there has been a substantial and long history of earthworm research in South Africa, for various purposes, most sampling in South Africa has been via digging and hand sorting (Table 2).

In this study we used hand digging with manual sorting and mustard extraction. Only one indigenous earthworm was collected during the whole study (*Tritogenia* sp.: Table 4).

TABLE 5
Table indicating the average sampling effort and costs per task or activity.*

| Task/ activity | Effort/ costs |
|--|--|
| Logistics (excluding accommodation for 1 person): | |
| Travel to all sites | R1 261 (R3.30/km for 382 km) |
| Car rental | R1 886 |
| Stipends | R2 250 (3 field assistants, R150/per person/day) |
| Air ticket | R1 573 |
| Food & tolls | R2 364 |
| Digging | Average of 55–60 seconds per worm |
| Mustard | Average of 21–30 seconds per worm |
| Preservation | Average of 15.5 seconds per worm |
| Determination of biomass | Average of 21–29 seconds per worm |
| Worm sampling and preservation | R100 per plot — sampling 30 earthworms per plot |
| Identification | 2hr/worm, R80/hr |

* To measure the amount of time, costs and effort needed per sample, detailed notes were kept on each of the different activities or tasks. These included: digging and collection of earthworms by hand, using the mustard solution collection technique and collecting earthworms, processing of samples (killing, labelling, adding to preservative, etc.), taking of soil samples and other environmental notes and measurements, preparation of plate for barcoding (including photographing specimens), weighing of specimens and identification of specimens. A detailed budget was kept for the entire project in order to estimate the logistical costs, the cost per sampling plot and the cost per soil sample. The values are as per South African Rand, January 2015

While this might be expected for disturbed areas such as gardens and agricultural areas, it was disconcerting to see a similar pattern in protected areas. This is likely the result of the human activities adjacent to and even in the protected areas. Similarly, Nxele (2012) collected five species of earthworm in grassland in Queen Elizabeth Park by digging 1 m × 1 m quadrats to a depth of 0.5 m: the indigenous *Tritogenia howickiana* and the non-indigenous *Pontoscolex corethrurus*, *Amyntas gracilis*, *Amyntas minimus*, and *Amyntas rodericensis*. These results are different from results obtained by Baretta *et al.* (2007) who also used digging and handsorting plus a chemical expellent, dilute Formol (0.5%). These authors found four native species and one exotic species when all methods were combined; however the most abundant was the exotic *Amyntas corticis*. The natural forests have high earthworm diversity because of high food availability and variability (Fragoso *et al.* 2006). In Karkloof forest, nine species were found and this is the second highest richness of all sampled sites. A native species was collected in this forest, similar to other studies in terms of diversity but completely different in terms of the number of natives found. For example, Huerta *et al.* (2007) studied the diversity of earthworms in heavily deforested Tabasco in Mexico. They sampled tropical rain forest and plantations of mango (*Mangifera indica*) and cacao (*Cacao theobroma*). These authors found 14 native species and five exotics, the tropical rain forest having the greater earthworm diversity.

The method of digging and hand-sorting soil from a 25 × 25 cm quadrat to 20 cm depth seems to be more suitable for sampling epigeic and endogeic earthworm species, probably as a result of their much higher densities in the upper soil layers. As our survey showed, no anecic earthworms were collected with our quadrat size. The size of each plot is a trade-off between collecting individuals and work. Based on available data and experience, we suggest following the ISO standard (i.e. a sample plot size of 50 cm × 50 cm and 20 cm deep) for smaller sized earthworms in South Africa, and digging larger and deeper quadrats for quantitatively sampling larger sized earthworms (i.e. plots of at least 1 m × 1 m × 0.5 m) (Nxele 2012).

Twelve soil samples appeared sufficient to sample most species of non-indigenous earthworms at species-poor sites, but not at species-rich sites as observed in the species accumulative curves (Supplementary material 2). The number of samples per site must be large enough to represent communities at the site adequately, but not so large that they are too time-consuming to process (Stark *et al.* 2001). From the literature review, the number of samples varied from five or six samples (Chaudhuri & Bhattacharjee 1999; Uys 2006) to 46 samples (Cuendet 1983) depending on the research purpose. Our literature survey indicates that given the high spatial and temporal variability in earthworm communities and variability introduced by different sampling techniques, giving a specific guidance on the sample size is difficult, and will likely be quite context specific.

Additional modifications are required for very large bodied taxa, as they were not collected with our quadrat size in our survey. First, one can dig even larger areas (e.g. approximately 2 m × 2 m × 0.5 m deep for earthworms up to around 100 cm long). Second, sampling of larger earthworms, especially giant earthworms, can be accomplished by searching for fresh casts (indicated by active deposition of fresh mud above ground), in a defined larger area, quickly topping the casts and grabbing the earthworm by the tail end, after which the tail end will anastomose and can immediately be preserved in 95 % ethanol. The species might then be identified using DNA barcoding, and the density

per unit area estimated. However, it should be noted that specimens can't be kept for morphological examination with this technique, and as such it is of limited use. Finally, some sampling will need to be opportunistic (e.g. sampling of giant earthworms that cross roads after rains). Owing to the sporadic nature of overland movement, a network of volunteers in various parts of the survey region might be able to track the weather conditions and collect specimens when conditions are right. The specimens would be useful for taxonomic study and a future atlas project.

Equipment requirements for earthworm sampling techniques and protocols are relatively modest (Supplementary Material 1). However, it takes a lot of time to sample and process samples. Time, resources and expensive equipment are limiting factors to zoological research globally (Bartlett *et al.* 2010; Butt & Grigoropoulou 2010, Uys *et al.* 2010). For these reasons the design inventories, sampling methods and planning and monitoring needs to be as effective and efficient as possible (Uys *et al.* 2010). Two people take 20 minutes to sample a plot of 50 × 50 × 20 cm by digging, and 13 minutes using mustard solution because of the waiting time involved. Mustard may be a useful and quicker method to sample alien earthworm species in leaf litter and topsoil. When using mustard solution, there is less labour involved compared to digging and hand sorting and using mustard solution does not affect the DNA of the specimens, which makes it suitable for molecular analyses. However, the results appear to be variable, and it is hard to control the amount that penetrates into the soil.

Setting a standard for the determination of earthworm biomass would make it possible to compare different datasets, but the biomass of an individual species or of a population as a whole can be used to estimate production and productivity of an ecosystem only when comparing ecosystems of the same or similar structure and species composition. The biomass of earthworm populations varies greatly and in most soils it exceeds the biomass of all other soil-inhabiting invertebrates (Edwards 1994). One practical issue is that the gut contents are a substantial proportion of an earthworm's biomass, and earthworms lose about 10–20% of their mass during fixation (Lee 1985), thus the point at which individuals are weighed must be made clear.

The major limitation to sampling is the ability to identify all the specimens collected to species level. With few taxonomists trained to identify South African earthworms (South Africa has one retired taxonomist and a full-time junior taxonomist at the KwaZulu-Natal Museum), additional capacity and new methods to facilitate species discovery and description are necessary. Earthworms are well suited to combining DNA-based analysis with morphological taxonomy (Hogg & Hebert 2003; Rougerie *et al.* 2009; Vernooy *et al.* 2010), an approach that has been used successfully in other parts of the world (Boyer *et al.* 2011, Porco *et al.* 2013). This approach involves training taxonomists in morphological identification, ensuring collections are developed and linked to DNA samples, cataloguing all relevant information and making it accessible.

The need to enlarge South Africa's capacity to answer applied and basic research questions on the roles and impacts of earthworms in the functioning of natural ecosystems was unanimously agreed upon during a workshop on soil organisms at the XVII Congress of the Entomological Society of Southern Africa (July 2011), and separately by the CAPE Invasive Alien Animal Working Group (May 2011) (Louw *et al.* 2014).

DNA bar-coding and development of user-friendly identification keys are needed to facilitate identification to species level. Given that DNA barcoding has been shown to be

extremely useful in the delimitation of difficult to identify species (Pop *et al.* 2003), the collection of samples in a DNA-compatible way should be included in future sampling protocols. In addition, juvenile specimens can be barcoded to verify their identifications (Richard *et al.* 2010). The earthworm barcoding library is growing on the Barcode of Life Data System (BOLD), also revealing interesting cryptic diversity (James & Davidson 2012, Porco *et al.* 2013, Decaëns *et al.* 2013).

The way ahead

Documenting the diversity of invertebrate groups should be a priority (Box 1) as invertebrate surveys, collections, and taxonomic descriptions should form an important part of setting conservation priorities (Essl *et al.* 2013). South Africa has achieved major success with atlas projects which have led to red-listing, volunteer networks and conservation actions (Harrison *et al.* 1997; Minter *et al.* 2004; Mecenero *et al.* 2013; Bates *et al.* 2014). A South African earthworm Atlas and Conservation Assessment would allow indigenous species of earthworms to be considered for conservation action through conservation plans, Environmental Impact Assessments (EIAs), and the like. Although South Africa is seen as a leader in the area of conservation planning (Balmford 2003), the number of EIAs or Biodiversity Assessments that have considered earthworms are few (McGeoch *et al.* 2011). Owing to the importance of earthworms in soil ecosystems, and their clear diversity importance in South Africa, this deficiency should be rectified. As a prelude to the Atlas project, we plan to run sampling and identification workshops in various provinces. The Atlas project would include both indigenous and non-indigenous species.

Data on diversity and distribution of earthworms have mostly been collected by taxonomists, building up large reference collections of study material (e.g. KZN Museum). These collections adequately serve the purpose of taxonomists but have limited use for environmental management, planning and conservation. Quantified, consolidated surveys and data for earthworms are patchy and scarce. Large amounts of material are collected but resources in terms of capacity and funds required to process the samples are limited. Although the plethora of different methodologies of earthworm sampling in use around the world could be a cause for inconsistencies in comparing different sets of data, the focus should be on efficiency, consistency, ease of use and practicability. When acquiring data during field trips, sampling trials needs to be maximised and optimised: i.e. gathering of soil data and biogeographic information, and measures of the effectiveness of the sampling techniques, as well as sampling all taxonomic groups present. These would be key sampling needs during surveys if questions on biodiversity, abundance and distributions are to be addressed. Surveys should be well planned and executed, and quantitative data collected, which in return will assist in obtaining diversity and distribution patterns for earthworms.

In addition to the integration of sampling techniques to extract earthworms from the soil, taxonomic identification and compiling a DNA barcode reference library, information is needed to assist in understanding the abiotic conditions that determine the spatial distribution of earthworms at the local scale. An assessment of soil quality that includes biological, chemical and physical properties can provide valuable information for evaluation of the sustainability of land management practices (Doran & Parkin 1994), the introduction of earthworms in areas for soil rehabilitation (Fründ *et al.* 2004), and

Box 1: Key priorities for work on megadrile earthworms in South Africa

- Compile an Atlas & Red List assessment.
- Develop a field guide to roughly separate specimens which will include pictures of live earthworms.
- Map and model species distributions based on climatic conditions, soils and land use.
- Engender public interest through outreach and participation to create awareness, expose and increase public interest on the subject.
- Develop a DNA barcode reference library.
- Construct a dated molecular phylogeny of indigenous South African species to enable a determination of key biogeographic breaks and explore evolutionary history.
- Develop a standardised protocol for sampling, collecting and curating earthworms (See Supplementary Material 1).
- Train future expertise for the identification of earthworm species in South Africa. This will be in the form of the training of students and farmers, annual “maintenance” training courses and the provision of identification services.
- Assess the risks of future introductions to South Africa and around South Africa via the various dispersal pathways.
- Determine the status and invasive potential of introduced species.
- Research the value of ecosystem services provided by earthworms under South African conditions, and the potential of species as bio-indicators of soil health.

the management of areas susceptible to earthworm invasions (Gundale 2002, Gundale *et al.* 2005).

To build our ecological and taxonomic knowledge of soil fauna in South Africa, studies of soil organisms need more coordination (Louw *et al.* 2014). In order to do this, one needs to move away from traditional *ad hoc* sampling to more coordinated, standardised sampling for comparison with other studies. The development of a survey plan should be interdisciplinary (involvement of soil scientists and biologists), thereby maximising sampling data and minimising effort, and thus will save time and money and assist with knowledge and information generation across disciplines and taxonomic groups. Also, increasing sampling intensity (effort) will increase detection probability.

CONCLUSION

We consider that the ISO sampling methodology augmented with more qualitative sampling is suitable for sampling most non-indigenous species of earthworm in South Africa. Sampling of indigenous species apparently requires a larger quadrat size or the use of other methods. Sampling and preservation techniques need to allow DNA-based studies to assist, complement and inform traditional morphology-based taxonomy. However, if we are to understand, conserve and manage earthworms we require more foundational knowledge. This will not happen without capacity-building and substantial resources.

ACKNOWLEDGEMENTS

Liezie Hartman provided assistance with the field-work and commented on various drafts of the manuscript. We thank M. Mostovski for useful comments on an earlier draft of the manuscript. This project was funded by the South African Working for Water Programme of the Department of Environmental Affairs and the National Research Foundation and Department of Science and Technology under the SABI Programme (grant # 79005), with support from the DST-NRF Centre of Excellence for Invasion Biology. We would

like to thank Ezemvelo KZN Wildlife for collection permits, and Mr Konanani, senior horticulturalist at the Pietermaritzburg Botanical Gardens and Mr Skhakhane, farm manager at Cedara Agricultural College for access to the properties. Field assistants (L. Bambalele, S. Mkhize and A. Malamlela) are acknowledged for their technical help during sampling.

REFERENCES

- BALMFORD, A. 2003. Conservation planning in the real world: South Africa shows the way. *Trends in Ecology and Evolution* **18**: 435–438.
- BARETTA, D., BROWN, G.G., JAMES, S.W. & CARDOSO, E.J.B.N. 2007. Earthworm populations sampled using collection methods in Atlantic forests with *Araucaria angustifolia*. *Science and Agriculture (Piracicaba, Brazil)* **64** (4): 384–392.
- BARTLETT, M.D., BRIONES, M.J.I., NELSON, R., SCHMIDT, O., SPURGEON, D. & CREAMER, R.E. 2010. A critical review of current methods in earthworm ecology: From individuals to populations. *European Journal of Soil Biology* **46**: 67–73.
- BATES, M.F., BRANCH, W.R., BAUER, A.M., BURGER, M., MARAIS, J., ALEXANDER, G.J., & DE VILLIERS, M.S., (eds). 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. *Suricata* **1**. South African National Biodiversity Institute, Pretoria.
- BEDDARD, F.E. 1895. Preliminary account on new species of earthworms belonging to the Hamburg Museum. *Proceedings of the Zoological Society of London* **63**: 210–248.
- BEYER, W.N., & GISH, C.D. 1980. Persistence in earthworms and potential hazards to birds of soil applied DDT, dieldrin and heptachlor. *Journal of Applied Ecology* **17**: 295–307.
- BEYER, W.N., & CROMARTIE, E.J. 1987. A survey of Pb, Cu, Zn, Cd, Cr, As, and Se in earthworms and soil from diverse sites. *Environmental Monitoring and Assessment* **8** (1): 27–36.
- BLAKEMORE, R.J. 2002. *Cosmopolitan earthworms – an eco-taxonomic guide to the peregrine species of world families*. Monograph. CD ROM.
- 2011. *Cosmopolitan Earthworms— an Eco-Taxonomic Guide to the Peregrine Species of the World*. Fourth Edition, VermEcology Solution, Yokohama, Japan.
- BORGONIE, G., GARCIA-MOYANO, A., LITTHAUER, D., BERT, W., BESTER, A., VAN HEERDEN, E., MOLLER, C., ERASMUS, M. & ONSTOTT, T.C. 2011. Nematoda from the terrestrial deep subsurface of South Africa. *Nature* **474**: 79–82.
- BOTES, A., MCGEOCH, M.A., ROBERTSON, H.G., VAN NIEKERK, A., DAVIDS, H.P. & CHOWN, S.L. 2006. Ants, altitude and change in the northern Cape Floristic Region. *Journal of Biogeography* **33**: 71–90.
- BOTES, A., MCGEOCH, M.A. & CHOWN, S.L. 2007. Ground-dwelling beetle assemblages in the northern Cape Floristic Region: Patterns, correlates and implications. *Austral Ecology* **32** (2): 210–224.
- BOUCHÉ, M.B & GARDNER, R.H. 1984. Earthworm functions VIII – Population estimation techniques. *Revue d'écologie et de Biologie du Sol* **21**: 37–63.
- BOYD, J.M. 1957. The Lumbricidae of a dune-machair soil gradient in Tiree, Argyll. *Journal of Natural History* **10** (112): 274–282.
- BOYER, S., BLAKEMORE, R.J., WRATTEN, S.D. 2011. An integrative taxonomic approach to the identification of three new New Zealand endemic earthworm species (Acanthodrilidae, Octochaetidae: Oligochaeta). *Zootaxa* **2994**: 21–32.
- BRASCHLER B., CHOWN S.L. & GASTON K.J. 2012 The Fynbos and Succulent Karoo Biomes do not have exceptional local ant richness. *PLoS ONE* **7** (3): 31463. doi:10.1371.
- BUTT, K.R. 1991. The effects of temperature on the intensive production of *Lumbricus terrestris* (Oligochaeta: Lumbricidae). *Pedobiologia* **35** (4): 257–264.
- BUTT, KR., LOWE, CN. & DUNCANSON, P. 2006. Earthworms of an urban cemetery. *8th International Symposium on Earthworm Ecology*, Krakow, Poland.
- BUTT, K.R. & GROGOROPOULOU, N. 2010. Field investigations of *Lumbricus terrestris* spatial distribution and dispersal through monitoring of manipulated, enclosed plots. *Soil Biology and Biochemistry* **42**: 40–47.
- CALLAHAM, M.A. JR & HENDRIX, P.F. 1997. Relative abundance and seasonal activity of earthworms (Lumbricidae and megascolecidae) as determined by hand-sorting and formalin extraction in forest soils on the Southern Appalachian Piedmont. *Soil Biology and Biochemistry* **29**: 317–321.
- CALLAHAM JR., M.A., HENDRIX, P.F., PHILLIPS, R.J. 2003. Occurrence of an exotic earthworm (*Amyntas agrestis*) in undisturbed soils of the southern Appalachian Mountains, USA. *Pedobiologia* **47**: 466–470.
- CHANG, C., ROUGERIE, R. CHEN, J. 2009. Identifying earthworms through DNA barcodes: Pitfalls and promise. *Pedobiologia* **52**: 171–180.
- CHAN, K. & MUNRO, K. 2001. Evaluating mustard extracts for earthworm sampling. *Pedobiologia* **45**: 272–278.

- CHAUDHURI, P.S. & BHATTACHARJEE, G. 1999. Earthworm resources of Tripura. *Proceedings – National Academy of Sciences India Section B* **69** (2): 159–170.
- CHAZDON, R. L., COLWELL, R. K., DENSLow, J. S. & GUARIGUATA, M. R. 1998. Statistical methods for estimating species richness of woody regeneration in primary and secondary rain forests of NE Costa Rica. In: Dallmeier, F. & Comiskey, J. A., eds., *Forest biodiversity research, monitoring and modeling: Conceptual background and Old World case studies*. Paris: Parthenon Publishing, pp. 285–309.
- ČOJA, T., ZEHEtNER, K., BRUCKNER, A., WATZINGER, A & MEYER, E. 2008. Efficacy and side effects of five sampling methods for soil earthworms (Annelida, Lumbricidae). *Ecotoxicology and Environmental Safety* **71**: 552–565.
- COLEMAN D.C., CROSSLEY D.A.J. & HENDRIX P.F. 2004. *Fundamentals of Soil Ecology*. Amsterdam, The Netherlands; Boston, MA, USA: Elsevier Academic Press.
- COLWELL, R.K. 2013. EstimateS 9.1.0: Statistical estimation of species richness and shared species from samples. (<http://viceroy.eeb.uconn.edu/estimates/>; accessed 29/12/2015).
- COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., PERFECT, T.J. & YEADON, R. 1980. Effects of cultivation and DDT on earthworm activity in a forest soil in the sub-humid tropics. *Journal of Applied Ecology* **17**: 21–29.
- COSTELLO, M.J., MAY, R.M. & STORK, N.E. 2013. Can we name Earth's species before they go extinct? *Science* **339**: 413–416.
- CROUCH, N.R. & SMITH, G.F. 2011. Informing and influencing the interface between biodiversity science and biodiversity policy in South Africa. *Botanical Journal of the Linnean Society* **166**: 301–309.
- CUENDET, G. 1983. Predation on earthworms by the Black-headed gull (*Larus ridibundus* L.). In: *Earthworm Ecology* (pp. 415–424). Springer Netherlands.
- CSUZDI, C. 2010. A monograph of the Paleotropical Benhamiinae earthworms. (Annelida: Oligochaeta, Acanthodrilidae). In: Csudi Cs. & Mahunka S., eds, *Pedozoologica Hungarica, Taxonomic, zoogeographic and faunistic studies on soil animals, No 6*. Budapest: Hungarian Natural History Museum.
- DAMOFF, G.A. & REYNOLDS, J.W. 2004. Additional earthworms (Oligochaeta: Lumbricidae) from Colorado, USA. *Megadrilogica* **10** (5): 29–37.
- DANIEL, O., JAGER, P., CUENDET, G. & BIERI, M. 1992. Sampling of *Lumbricus terrestris* (Oligochaeta: Lumbricidae) *Pedobiologia* **36**: 231–220.
- DARWIN, C. 1881. The formation of vegetable mould through the action of earthworms with some observations of their habits. John Murray, UK, 326 pp.
- DAVIS, A. 1997. Climatic and biogeographical associations of southern African dung beetles (Coleoptera: Scarabaeidae s. str.). *African Journal of Ecology* **35** (1): 10–38.
- DAVIS, A.V.L., SCHOLTZ, C.H. & CHOWN, S.L. 1999. Species turnover, community boundaries, and biogeographical composition of dung beetle assemblages across an altitudinal gradient in South Africa *Journal of Biogeography* **26**: 1039–1055.
- DECAËNS T., PORCO D., ROUGERIE R., BROWN G.G. & JAMES S.W. 2013. Potential of DNA barcoding for earthworm research in taxonomy and ecology. *Applied Soil Ecology* **65**: 35–42.
- DE ZORZI, P., BARBIZZI, S., BELLI, M., CICERI, G., FAJGELJ, A., MOORE, D., SANSONE, U. & VAN DER PERK, M. 2005. Terminology in soil sampling (IUPAC Recommendations 2005). *Pure and Applied Chemistry* **77**: 827–841.
- DIPPENAAR-SCHOEMAN, A.S., & CRAEMER, C. 2000. The South African National Survey of Arachnida. *Plant Protection News* **56**: 11–12.
- DLAMINI, T.C. HAYNES, R.J. & VAN ANTWERPEN, R. 2001. Exotic earthworm species dominant in solid on sugarcane estates in the Eshowe area of the North Coast of KwaZulu-Natal. *Proceedings of the South African Sugar Technologists' Association* **75**: 217–221.
- DORAN, J.W. & T.B. PARKIN. 1994. Defining and assessing soil quality. In: J.W. Doran et al., eds, *Defining Soil Quality for a Sustainable Environment*, pp 3–21, SSSA Special Publication No. 35, Soil Science Society of America and American Society of Agronomy, Madison, WI.
- DRUCE, D., HAMER, M. & SLOtOW, R. 2004. Sampling strategies for millipedes (Diplopoda), centipedes (Chilopoda) and scorpions (Scorpionida) in savanna habitats. *African Zoology* **39**: 293–304.
- EAST, D. & KNIGHT, D. 1998. Sampling soil earthworm populations using household detergent and mustard. *Journal of Biological Education* **32**: 201–206.
- EDWARDS, C.A., & LOFTY, J.R. 1982. Nitrogenous fertilizers and earthworm populations in agricultural soils. *Soil Biology and Biochemistry* **14** (5): 515–521.
- EDWARDS, C.A. 1994. *Earthworm Ecology – Second edition*. London: CRC Press.
- EDWARDS, C.A. & BOHLEN, P.J. 1996. *Biology and Ecology of earthworms*. London: Chapman & Hall.
- EICHINGER, E., BRUCKNER, A. & STEMMER, M. 2007. Earthworm expulsion by formalin has severe and lasting side effects on soil biota and plants. *Ecotoxicology and Environmental Safety* **67**: 260–266.

- EISENHAUER, N., PARTSCH, S., PARKINSON, D. & SCHEU, S. 2007. Invasion of a deciduous forest by earthworms: changes in soil chemistry, microflora, microarthropods and vegetation. *Soil Biology and Biochemistry* **39** (5): 1099–1110.
- EISENHAUER, N. & SCHEU, S. 2008. Invasibility of experimental grassland communities: the role of earthworms, plant functional group identity and seed size. *Oikos* **117** (7): 1026–1036.
- EISENHAUER, N., STRAUBE, D. & SCHEU, S. 2008. Efficiency of two widespread non-destructive extraction methods under dry soil conditions for different ecological earthworm groups. *European Journal of Soil Biology* **44**: 141–145.
- ENCKELL, P.H., BENGSTON, S.A., DOUWES, P., NIKLASSON, M., STILLE, B., & WIMAN, B. 1986. The dispersal pattern of an anthropochorous species: Genetic variation in populations of *Lumbricus terrestris* L. (Lumbricidae) in the Faroe Islands. *Hereditas*, **104** (2): 253–261.
- ENCKELL, P.H. & RUNDGREN, S. 1988. Anthropochorous earthworms (Lumbricidae) as indicators of abandoned settlements in the Faroe Islands. *Journal of Archaeological Science* **15** (4): 439–451.
- ERMILOV, S.G. & HUGO-COETZEE, A. 2012. Oribatid mites of the genus *Eremulus* Berlese, 1908 (Acari: Oribatida: Eremulidae) from South Africa. *African Invertebrates* **53** (2): 559–569.
- ESSL, F., MOSER, D., DIRNBÖCK, T., DULLINGER, S., MILASOWSKY, N., WINTER, M., & RABITSCH, W. 2013. Native, alien, endemic, threatened, and extinct species diversity in European countries. *Biological Conservation* **164**: 90–97.
- EVANS, A.C. & GUILD, W.J. 1947. Studies on the relationships between earthworms and soil fertility: I. Biological studies in the field. *Annals of Applied Biology* **34**: 307–330.
- FRAGOSO, C. & REYNOLDS, J.W. 1997. On some earthworms from central and southeastern Mexican mountains, including two new species of the genus *Dichogaster* (Dichogastrini). *Megadriologica* **7** (2): 9–19.
- FRAGOSO, C., LAVELLE, P., BLANCHART, E., SENAPATI, B.K., JIMÉNEZ, J.J., MARTÍNEZ, M.A., DECAËNS, T. & TONDOH, J. 2006. Earthworm communities of tropical agroecosystems: Origin, structure and influence of management practices. In: Lavelle, P.; Brussaard, L.; Hendrix, P., eds, *Earthworm management in tropical agroecosystems*. Wallingford: CAB International, p.27–55.
- FRÜND, H., EGBERT, E. & DUMBECK, G. 2004. Spatial distribution of earthworms [Lumbricidae] in recultivated soils of the Rhenish lignite-mining area, Germany. *Journal of Plant Nutrition and Soil Science* **167**: 494–502.
- GATES, G.E. 1972. Burmese earthworms: an introduction to the systematics and biology of megadrile oligochaetes with special references to Southeast Asia. *Transactions of the American Philosophical Society* **62** (7): 1–326.
- GRUBE, E. 1879. Annelida. *Philosophical Transactions of the Royal Society of London* **168**: 554–556.
- GUGIC, D., NASSIRI, M., NADJI, M., MORALES, A. & VINCEK, V. 2007. Novel tissue preservative and tissue fixative for comparative pathology and animal research. *Journal of Experimental Animal Science* **43**: 271–281.
- GUILD, W.J.M. 1951. The lumbricidae in upland areas. I. Population variation in a small glen. *Journal of Natural History* **4** (43): 653–657.
- 1952a. XXXII.—The Lumbricidae in upland areas. II. Population variation on hill pasture. *Journal of Natural History*, **5** (51): 286–292.
- 1952b. Variation in earthworm numbers within field populations. *The Journal of Animal Ecology* **21**: 169–181.
- 1957. Occurrence of populations of *Alloobophora limicola* (Lumbricidae), in SW Scotland. *Journal of Natural History* **10** (117): 665–668.
- GUNDALE, M.J. 2002. The influence of exotic earthworms on soil organic horizon and the rare fern *Botrychium mormo*. *Conservation Biology* **16**: 1555–1561.
- GUNDALE, M.J., JOLLY, W.M. & DELUCA, T.H. 2005. Susceptibility of a northern hardwood forest to exotic earthworm invasion. *Conservation Biology* **19**: 1075–1083.
- GUNN, A. 1992. The use of mustard to estimate earthworm populations. *Pedobiologia* **36**: 65–67.
- HAIMI, J. & BOUCELHAM, M. 1991. Influence of a litter feeding earthworm. *Pedobiologia* **35**: 247–251.
- HAMER, M. 2010. African Invertebrates in the International Year of Biodiversity. *African Invertebrates* **51**: 223–230.
- HAYNES, R.J., DOMINY, C.S. & GRAHAM, M.H. 2003. Effect of agricultural land use on soil organic matter status and the composition of earthworm communities in KwaZulu-Natal, South Africa. *Agriculture, Ecosystems and Environment* **95**: 453–464.
- HENDRIX, P.F., CALLAHAN, M.A., DRAKE, J.M., HUANG, C., JAMES, S.W., SNYDER, B.A. & ZHANG, W. 2008. Pandora's Box contained bait: The global problem of introduced earthworms. *The Annual Review of Ecology, Evolution and Systematics* **39**: 593–613.
- HENDRIKSEN, N.B. 1990. Leaf litter selection by detritivore and geophagous earthworms. *Biology and Fertility of Soils* **10** (1): 17–21.

- HOGG, I.D. & HEBERT, P.D.N. 2004. Biological identification of springtails (Hexapoda: Collembola) from the Canadian Arctic, using mitochondrial DNA barcodes. *Canadian Journal of Zoology* **82** (5): 749–754.
- HOPP, H. 1947. The Ecology of Earthworms in Cropland. *Soil Science Society of America* **12**: 503–507.
- HORN, J.L., PLISKO, J.D. & HAMER, M.L. 2007. The leaf-litter earthworm fauna (Annelida: Oligochaeta) of forests in Limpopo Province, South Africa: diversity, communities and conservation. *African Zoology* **42**: 172–179.
- HORST, R. 1893. Earthworms from the Malaya Archipelago. In: Weber, M., ed., *Zoologische Ergebnisse einer Reise in Niederländisch Ost-Indien*. Vol. 3. Leiden: E.J. Brill, pp. 28–77.
- HUERTA, E., RODRIGUEZ-OLAN, J., EVIA-CASTILLO, I., MONTEJO-MENESES, E., DE LA CRUZ-MONDRAGÓN, M., GARCIA-HERNANDEZ, R. & URIBE, S. 2007. Earthworms and soil properties in Tabasco, Mexico. *European Journal of Soil Biology* **43**: S190–S195.
- INTERNATIONAL ORGANIZATION FOR STANDARDIZATION (ISO). 1999. Soil quality: Guidance on the determination of effects in field situations Part 3. ISO 11268-3. Geneva, Switzerland.
- INTERNATIONAL ORGANIZATION FOR STANDARDIZATION (ISO). 2006. Soil quality – Sampling of Soil Invertebrates Part 1: Hand-sorting and formalin Extraction of Earthworms. ISO 23611-1. Geneva, Switzerland.
- JANION, C., BEDOS, A., BENGTTSSON, J., DEHARVENG, L., JANSEN VAN VUUREN, B., LEINAAS, H.P., LIU, A., MALMSTRÖM, A., PORCO, D. & CHOWN, S.L. 2011. Springtail diversity in South Africa: research letter. *South African Journal of Science* **107** (11 & 12): 1–7.
- JANION-SCHEEPERS, C., DEHARVENG, L., BEDOS, A. & CHOWN, S.L. 2015. Updated list of Collembola species currently recorded from South Africa. *ZooKeys* **503**: 55–88.
- JAMES, S.W., PORCO, D., DECAËNS, T., RICHARD, B., ROUGERIE, R. & ERSÉUS, C. 2010. DNA Barcoding reveals cryptic diversity in *Lumbricus terrestris* L., 1758 (Clitellata): Resurrection of *L. Herculeus* (Savigny, 1826). *PLoS ONE* **5**: e15629.
- JAMES, S.W. & DAVIDSON, S.K. 2012. Molecular phylogeny of earthworms (Annelida: Crassicitellata) based on 28S, 18S and 16S gene sequences. *Invertebrate Systematics* **26**: 213–229.
- JAMIESON, B.G.M. 1967. New records of Onerodrilidae and Lumbricidae (Oligochaeta) from South African Rivers. *Zoologica Africana* **3** (1): 59–86.
- JOANNES, H., & KRETZSCHMAR, A. 1983. Model of gut transit in earthworms *Nicodrilus* spp. *Revue d'écologie et de biologie du sol*, **20** (3): 349–366.
- KEITH, A.M. & SCHMIDT, O. 2013. First record of the earthworm *Proselodrilus amplisetosus* (Oligochaeta: Lumbricidae) outside continental Europe. *Irish Naturalists Journal* **32** (1): 26–28.
- KINBERG, J.G.H. 1867. *Annulata nova. Öfversigt af Kongliga Vetenskaps-akademiens Förhandlingar* **23**: 97–100.
- KNOETZE, R., MALAN, A.P. & MOUTON, C. 2006. Differentiation of South African potato cyst nematodes (PCN) by analysis of the rDNA internal transcribed spacer region. *African Plant Protection* **12**: 103–110.
- LAVELLE, P., DECAËNS, T., AUBERT, M., BAROT, S., BLOUIN, M., BUREAU, F., MARGERIE, P., MORA, P. & ROSSI, J.P. 2006. Soil invertebrates and ecosystem services. *European Journal of Soil Biology* **42**: S3–S15.
- LAVERACK, M.S. 1960. Tactile and chemical perception in earthworms—I. Responses to touch, sodium chloride, quinine and sugars. *Comparative Biochemistry and Physiology*, **1** (2): 155–163.
- 1961. The effect of temperature changes on the spontaneous nervous activity of the isolated nerve cord of *Lumbricus terrestris*. *Comparative Biochemistry and Physiology* **3** (2): 136–140.
- LAWRENCE, A.P. & BOWERS, M.A. 2002. A test of the “hot” mustard extraction method of sampling earthworms. *Soil Biology and Biochemistry* **34**: 549–552.
- LEHMANN, U. & KREIPE, H. 2001. Real-time PCR analysis of DNA and RNA extracted from formalin-fixed and paraffin-embedded biopsies. *Methods* **25**: 409–418.
- LJUNGSTRÖM, P. 1972. Introduced earthworms of South Africa. On their taxonomy, distribution, history of introduction and on the extermination of endemic earthworms. *Zool. Jb. Syst. Bd.* **99**: 1–81.
- LONGINO, J.T., CODDINGTON, J. & COLWELL, R.K. 2002. The ant fauna of a tropical rain forest: estimating species richness three different ways. *Ecology* **83** (3): 689–702.
- LOUW, S., WILSON, J.R.U., JANION, C., VELDTMAN, R., DAVIES, S.J. & ADDISON, M. 2014. The unknown underworld: Understanding soil health in South Africa. *South African Journal of Science* **110**: Art. #a0064.
- MARTIN, N.A. 1980. Earthworm (Oligochaeta: Lumbricidae) populations and late summer pasture renovation. *New Zealand Journal of Agricultural Research* **23** (3): 417–419.
- MCGEOCH, M.A., SITHOLE, H., SAMWAYS, M.J., SIMAIKA, J.P., PRYKE, J.S., PICKER, M., UYS, C., ARMSTRONG, A., DIPPENAAR-SCHOEMAN, A.S., ENGELBRECHT, I.A., BRASCHLER, B., HAMER, M. 2011. Conservation and monitoring of invertebrates in terrestrial protected areas. *Koedoe* **52**: 1–13. Art.#1000

- MECENERO, S., BALL, J.B., EDGE, D.A., HAMER, M.L., HENNING, G.A., KRÜGER, M., PRINGLE, E.L., TERBLANCHE, R.F. & WILLIAMS, M.C., (eds). 2013. *Conservation assessment of butterflies of South Africa, Lesotho and Swaziland: red list and atlas*. Safronics (Pty) Ltd, Johannesburg & Animal Demography Unit, Cape Town.
- MICHAELSEN, W. 1913. Report upon the Oligochaeta in the South African Museum at Cape Town. *Annals of the South African Museum* **13**: 43–62.
- MINTER, L.R., BURGER, M., HARRISON, J.A., BRAAK, H.H., BISHOP, P.J. & KLOEPFER, D., (eds). 2004. Atlas and Red Data Book of the Frogs of South Africa, Lesotho and Swaziland. SI/MAB Series #9. Smithsonian Institution, Washington, DC.
- MISCHIS, C.C., HERRERA, J.A.D. & FREGUGLIA, M.A. 1997. Earthworm community distribution (Annelida, Oligochaeta) along an altitudinal gradient on altas cumbres road (Cordoba, Argentina). *Megadrilogica* **6** (11): 97–104.
- MOELANS, C.B., OOSTENRIJK, D., MOONS, M.J. & VAN DIEST, P.J. 2011. Formaldehyde substitute fixatives: effects on nucleic acid preservation. *Clinical Pathology* **64**: 960–967.
- MORENO, A.G., JESUS, J.B., & DIAZ COSIN, D.J. 1982. Contribution to the knowledge of Spanish Lumbricidae: II. *Dendrobaena alvaradoi* n. sp. *Italian Journal of Zoology* **49** (1–2): 21–24.
- MÜLLER, F. 1857. *Lumbricus corethrurus*, Bürstenschwanz. *Archiv für Naturgeschichte* **23**: 113–116.
- MURCHIE, W.R. 1958. Biology of the oligochaete *Eisenia rosea* (Savigny) in an upland forest soil of southern Michigan. *American Midland Naturalist* **60**: 113–131.
- NELSON, J.M. & SATCHELL, J.E. 1962. The extraction of Lumbricidae from soil with special reference to the hand-sorting method. In: Murphy, P.W., ed., *Progress in Soil Zoology*. London: Butterworths, London, pp. 294–299.
- NAEEM, S., THOMPSON, L.J., LAWLER, S. P., LAWTON, J. H., & WOODFIN, R. M. 1994. Declining biodiversity can alter the performance of ecosystems. *Nature* **368** (6473): 734–737.
- NORDSTRÖM, S. & RUNDGREN, S. 1972. Methods of sampling lumbricids. *Oikos* **23**: 344–352.
- 1973. Associations of lumbricids in Southern Sweden. *Pedobiologia* **13**: 301–326.
- NUUTINEN, V. & HAUKKA, J. 1990. Conventional and organic cropping systems at Suitia. VII: Earthworms. *Journal of Agricultural Science in Finland* **62** (4): 357–367.
- NXELE, T.C. 2012. The megadrile fauna (Annelida: Oligochaeta) of Queen Elizabeth Park, South Africa: species composition and distribution within different vegetation types. *African Invertebrates* **53**: 543–558.
- OMODEO, P. & MARTINUCCI, G.B. 1987. Comparison of the earthworm fauna of some oak forests in Italy and Algeria. In: P. Bonvicini Pagliai, & P. Omodeo, eds, *On earthworms. Selected symposia and monographs UZI*, Vol. 2, pp. 225–234.
- ÖRLEY, L. 1881. A magyarországi Oligochaeták faunája. I. Terricolae. *Mathematikai és természettudományi Közlemények* **16**: 562–611.
- PARR, C.L., SINCLAIR, B.J., ANDERSEN, A.N., GASTON, K.J. & CHOWN, S.L. 2005. Constraint and competition assemblages: cross-continental and modelling approach for ants. *American Naturalist* **165**: 481–494.
- PAVLÍČEK, T., CSUZDI, C., SMOOHA, G., BEILES, A. & NEVO, E. 1996. Biodiversity and microhabitat distribution of earthworms at “Evolution Canyon,” a Mediterranean microsite, Mount Carmel, Israel. *Israel Journal of Zoology* **42** (4): 449–454.
- PELOSI, C., BERTRAND, M., ČAPOWIEZ, Y., BOLZARD, H. & ROGER-ESTRADE, J. 2009. Earthworm collection from agricultural fields: Comparisons of selected expellants in presence/absence of hand-sorting. *European Journal of Soil Biology* **45**: 176–183.
- PICKFORD, G.E. 1937. A monograph of the Acanthodriline earthworms of South Africa. Cambridge: W Heffer & Sons Ltd.
- PLISKO, J.D. 1991. Two new species of *Microchaetus* Rapp, 1849 from the Eastern Cape Province of South Africa (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **32**: 293–303.
- 1992. The Microchaetidae of Natal, with descriptions of new species of *Microchaetus* Rapp and *Tritogenia* Kinberg, and the new genus *Proandricus* (Oligochaeta). *Annals of the Natal Museum* **33**: 337–378.
- 1993a. Eight new species *Proandricus* Plisko, 1992 from South Africa (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **34** (2): 199–220.
- 1993b. Four new species, and new records, of *Microchaetus* Rapp, 1849 from the Cape Province, South Africa, and Transkei (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **34** (2): 221–237.
- 1996a. *Michalakus*, a remarkable new genus of microchaetid earthworm from South Africa (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **37**: 287–293.
- 1996b. Six new earthworm species of the southern African genus *Proandricus* Plisko, 1992 (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **37**: 295–307.
- 1997. New species of genus *Tritogenia* Kinberg, 1867 from southern Africa (Oligochaeta: Microchaetidae). *Annals of the Natal Museum* **38**: 241–281.

- 2000. The role of nature reserves in the protection of the terrestrial earthworm fauna (Oligochaeta), based on the material from Dlinza Forest Nature Reserve (Kwazulu-Natal, South Africa). *Lammergeyer* **46**: 75–80.
- 2001. Notes on the occurrence of the introduced earthworm *Pontoscolex corethrurus* (Müller, 1857) in South Africa (Oligochaeta: Glossoscolecidae). *African Invertebrates* **42**: 323–334.
- 2002a. Nine new earthworm species of *Proandricus* Plisko, 1992 from South Africa and Lesotho (Oligochaeta: Microchaetidae). *African Invertebrates* **43**: 183–203.
- 2002b. Three new earthworm species of *Microchaetus* Rapp, 1849, and new data on two earlier known species of this genus (Oligochaeta: Microchaetidae). *African Invertebrates* **43**: 205–214.
- 2003. Eleven new South African earthworms (Oligochaeta: Microchaetidae) with new information on some known species, and an inventory of the microchaetids of KwaZulu-Natal. *African Invertebrates* **44** (2): 279–325.
- 2004. Review of the balantine genus *Udeina* Michaelsen, 1910 with descriptions of six new species in South Africa (Oligochaeta: Acanthodrilidae, Acanthodrilinae). *African Invertebrates* **45**: 287–313.
- 2006. A systematic reassessment of the genus *Microchaetus* Rapp, 1849: its amended definition, reinstatement of *Geogenia* Kinberg, 1867, and erection of a new genus *Kazimierzus* (Oligochaeta: Microchaetidae). *African Invertebrates* **47**: 31–56.
- 2007. New species of South African acanthodriline earthworms of the genera *Eodriloides* and *Chilota*, with a redescription of *Chilota quindecimus* (Oligochaeta: Acanthodrilidae). *African Invertebrates* **48**: 33–40.
- 2010. Megadrile earthworm taxa introduced to South African soils (Oligochaeta: Acanthodrilidae, Eudrilidae, Glossoscolecidae, Lumbricidae, Megascolecidae, Ocnodrilidae). *African Invertebrates* **51**: 289–312.
- PLISKO, J.D. & NXELE, T.C. 2015. An annotated key separating foreign earthworm species from the indigenous South African taxa (Oligochaeta: Acanthodrilidae, Eudrilidae, Glossoscolecidae, Lumbricidae, Megascolecidae, Microchaetidae, Ocnodrilidae and Tritogeniidae). *African Invertebrates* **56** (3): 663–708.
- PLISKO, J.D. & ZICSI, A. 1991. Über neue *Tritogenia*-Arten aus Süd-Afrika (Oligochaeta: Microchaetidae). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* **88**: 111–123.
- POP, V.V. 1987. Density and biomass of earthworm synusia in forest ecosystems of the Romanian Carpathians. In *On earthworms. Selected Symposia and Monographs*. *UZI* **2**: 183–190.
- POP, V.V. & POSTOLACHE, T. 1987. Giant earthworms build up vermic mountain rendzinas. *On Earthworms*. *Mucchi, Modena* 141–150.
- POP, A.A., WINK, M. & POP, V.V. 2003. Use of 18S, 16S rDNA and cytochrome c oxidase sequences in earthworm taxonomy (Oligochaeta, Lumbricidae). *Pedobiologia* **47**: 428–433.
- PORCO D., DECAËNS T., DEHARVENG L., JAMES S.W., SKARZYNSKI D., ERSÉUS C., BUTT K., RICHARD B. & HEBERT P. 2013. Biological invasions in soil: DNA barcoding as a monitoring tool in a multiple taxa survey targeting European earthworms and collembolans in North America. *Biological Invasions* **13**: 899–910.
- RAW, F. 1959. Estimating earthworm populations by using formalin. *Nature* **14**: 1661–1662.
- RAW, F. 1962. Studies of earthworm populations in orchards. *Annals of Applied Biology* **50** (3): 389–404.
- REID, W.V., MOONEY, H. A., CROPPER, A., CAPISTRANO, D., CARPENTER, S.R., CHOPRA, K., DASGUPTA, P., DIETZ, T., DURAIAPPAH, A.K., HASSAN, R., KASPERSON, R., LEEMANS, R., MAY, R.M., MCMICHAEL, T. (A.J.), PINGALI, P., SAMPER, C., SCHOLLES, R., WATSON, R.T., ZAKRI, A.H., SHIDONG, Z., ASH, N.J., BENNETT, E., KUMAR, P., LEE, M.J., RAUDSEPP-HEARNE, C., SIMONS, H., THONELL, J. & ZUREK, M.B. 2005. Millennium Ecosystem Assessment. Ecosystems and Human Well-being: Synthesis. Washington, DC: Island Press.
- REEVES, W.K. & REYNOLDS, J.W. 2004. Earthworms (Oligochaeta: Glossoscolecidae, Megascolecidae and Octochaetidae) and terrestrial polychaetes (Polychaeta: Nereididae) of Grand Cayman Island, with notes on their natural enemies. *Megadrilologica* **10** (6): 39–46.
- REINECKE, A.J. & ACKERMAN, D. 1977. New Earthworm species (Octochaetinae) from the North-Eastern Transvaal, South Africa. *Wetenskaplike bydraes van die PU vir CHO* **90**: 1–12.
- REINECKE, A.J. & LJUNGSTRÖM, P.O. 1969. An ecological study of earthworms from banks of the Mooi River in Potchefstroom South Africa. *Pedobiologia* **9** (1–2): 106–111.
- REINECKE, A.J. & RYKE, P.A. 1970. On the water relations of *microchaetus modestus* Mich. (Microchaetidae, Oligochaeta). *Wetenskaplike bydraes van die Potchefstroomse Univeriteit vir C.H.O (B)* **15**: 1–21.
- REINECKE, A.J. & RYKE, P.A.J. 1972. 'n Onderzoek na die ekologie van *Microchaetus modestus* (Oligochaeta), 'n Suid-Afrikaanse endemiese spesie. *Wetenskaplike bydraes van die PU vir CHO, Reeks B: Natuurwetenskappe* **42**: 1–42.
- REYNOLDS, J.W. 1972. Relationship of earthworm (Oligochaeta: Acanthodrilidae and Lumbricidae) distribution and biomass in six heterogeneous woodlot sites in Tippecanoe County, Indiana. *Tennessee Academy of Science* **47** (2): 63–67.

- 1993. On some Earthworms from Swaziland (Oligochaeta: Glossoscolecidae, Megascolidae, Microchaetidae and Octochaetidae) *Megadrilologica* **5** (1): 1–8.
- REYNOLDS, J.W., FRAGOSO, C.G. & GUERRA, C.A. 1995. New earthworm records from Belize, C.A. (Oligochaeta: Acanthodrilidae, Eudrilidae, Glossoscolecidae, Megascolecidae, Ocnodrilidae and Octochaetidae). *Megadrilologica* **6** (7): 63–70.
- REYNOLDS, J.W. & HANEL, C. 2005. The earthworms (Oligochaeta: Lumbricidae) of Tristan da Cunha and Nightingale Islands, South Atlantic Ocean. *Megadrilologica* **10** (7): 47–56.
- REYNOLDS, T.B. 1955. Observations on the earthworms of north Wales. *The North Western Naturalist* **3**: 291–304.
- REYNOLDS, T.B., O'CONNOR, F.B. & KELLY, W.A. 1955. Observations on the earthworms of Bardsey. *Bardsey Observatory Report* **9**.
- RICHARD, B., DECAËNS T., ROUGERIE R., JAMES S., PORCO D., HEBERT P. 2010 Re-integrating earthworm juveniles in soil diversity studies: species identification through DNA barcoding. *Molecular Ecology Resources* **10**: 606–614.
- ROUGERIE, R., DECAËNS. T., DEHARVENG, L., PORCO, D., JAMES, S.W., CHANG, C., RICHARD, B., POTAPOV, M., SUHARDJONO, Y. & HEBERT, P.D.N. 2009. DNA barcodes for soil animal taxonomy. *Pesquisa Agropecuária Brasileira* **44**: 789–801.
- SATCHELL, J.E. 1970. Measuring population and energy flow in earthworms. In: Phillipson, J. ed., *Methods of study in soil ecology*. Paris: UNESCO, pp. 261–267.
- SAVIGNY, J.C. 1826. [La multiplicité des espèces de ver de terre]. In: Cuvier, G: Analyse des Travaux de l'Académie royale des Sciences, pendant l'année 1821, partie physique. *Mémoires de l'Académie des Sciences de l'Institut de France (Physique)*, Paris **5**: 176–184.
- SCHEU, S. 1987. The role of substrate feeding earthworms (Lumbricidae) for bioturbation in a beechwood soil. *Oecologia* **72** (2): 192–196.
- SCHEU, S. 1993. The earthworm (Lumbricidae) distribution in Alberta (Canada). *Magadrilologica* **4** (11): 175–185.
- SCHWERT, D.P. 1977. The first North American record of *Aporrectodea icterica* (Savigny, 1826) (Oligochaeta, Lumbricidae), with observations on the colonization of exotic earthworm species in Canada. *Canadian Journal of Zoology* **55** (1): 245–248.
- SIMS, R.W. 1967. Earthworms (Acanthodrilidae and Eudrilidae: Oligochaeta) from Gambia.
- SLOTOW, R & HAMER, M. 2000. Biodiversity research in South Africa: comments on current trends and methods. *South African Journal of Science* **96**: 222–224.
- STARK, J.D., BOOTHROYD, I.K.G., HARDING, J.S., MAXTED, J.R & SCARSBROOK, M.R. 2001. Protocols for sampling macro invertebrates in wadeable streams. *New Zealand Macro invertebrate Working Group Report No. 1 – 57*.
- STEENKAMP, Y. & SMITH, G.F. 2006. Introduction to A checklist of South African plants. In: Germishuizen, G., Meyer, N.L., Steenkamp, Y. & Keith, M., eds. *A checklist of South African plants*. Southern African Botanical Diversity Network Report 41. Pretoria: SABONET, pp. iv–ix.
- SUTHERLAND, W.J., FRECKLETON, R.P., GODFRAY, H.C.J., BEISSINGER, S.R., BENTON, T., CAMERON, D.D. CARMEL, Y., COOMES, D.A., COULSON, T., EMMERSON, M.C., HAILS, R.S., HAYS, G.C., HODGSON, D.J., HUTCHINGS, M.J., JOHNSON, D., JONES, J.P.G., KEELING, M.J., KOKKO, H., KUNIN, W.E., LAMBIN, X., LEWIS, O.T., MALHI, Y., MIESZKOWSKA, N., MILNER-GULLAND, E.J., NORRIS, K., PHILLIMORE, A.B., PURVES, D.W., REID, J.M., REUMAN, D.C., THOMPSON, K., TRAVIS, J.M.J., TURNBULL, L.A., WARDLE, D.A. & WIEGAND, T. 2013. Identification of 100 fundamental ecological questions. *Journal of Ecology* **101**: 58–67.
- THAKURIA, D., SCHMIDT, O., LILIENSIEK, A., EGAN, D. & DOOHAN, F.M. 2009. Field preservation and DNA extraction methods for intestinal microbial diversity analysis in earthworms. *Journal of Microbiological Methods* **76**: 226–233.
- TOMLIN, A.D. & GORE, F.L. 1974. Effects of six insecticides and a fungicide on the numbers and biomass of earthworms in pasture. *Bulletin of environmental contamination and toxicology* **12** (4): 487–492.
- UYS, C.J. 2006. Invertebrate diversity in Afrotemperate forests: spatial and seasonal changes and implications for conservation. MSc Thesis. Pietermaritzburg: University of KwaZulu-Natal.
- UYS, C., HAMER, M. & SLOTOW, R. 2010. Step process for selecting and testing surrogates and indicators of afro temperate forest invertebrate diversity. *PLoS ONE* **5**: e11000.
- VALCKX, J., COCKX, L., WAUTERS, J., VAN MEIRVENNE, M., GOVERS, G., HERMY, M. & MUYS, B. 2009. Within-field spatial distribution of earthworm populations related to species interactions and soil apparent electrical conductivity. *Applied Soil Ecology* **41**: 315–328.
- VALCKX, A., GOVERS, G., HERMY, M. & MUYS, B. 2011. Biology of earthworms. *Soil Biology* **24**: 19–38.
- VERNOOY, R., HARIBABU, E., MULLER, M.R., VOGEL, J.H., HEBERT, P.D.N., SCHINDEL, D.E., SHIMURA, J. & SINGER, G.A.C. (2010) Barcoding life to conserve biological diversity: Beyond the taxonomic imperative. *PLoS Biology* **8**: e1000417.

- VISSER, F.A. & REINECKE, A.J. 1977. The earthworms of the Mooi River irrigation area in Potchefstroom, South Africa (Oligochaeta: Lumbricidae, Acanthodrilidae, Microchaetidae and Ocnodrilidae). *Publicaciones del Centro Pirenaico de Biología Experimental* **9**: 95–108.
- WARDLE, D.A. 2002. Communities and ecosystems: linking the aboveground and belowground components (Vol. 34). Princeton: Princeton University Press.
- WARDLE, D.A., BARDGETT, R.D., KLIRONOMOS, J.N., SETÄLÄ, H., VAN DER PUTTEN, W.H., & WALL, D.H. 2004. Ecological linkages between aboveground and belowground biota. *Science* **304** (5677): 1629–1633.
- ZARBORSKI, E.R. 2003. Allyl isothiocyanate: an alternative chemical expellant for sampling earthworms. *Applied Soil Ecology* **22**: 87–95.

SUPPLEMENTARY MATERIAL 1: A PROPOSED PROTOCOL FOR SAMPLING EARTHWORMS

Equipment list

- 2 × measuring tape (5 m).
- String or rope.
- markers.
- distilled water (amount depends on how many plots needed; for 60 g of mustard, 1L of water is needed).
- mustard powder (60 g per 1 L of water).
- watering can.
- cleansing tray.
- GPS.
- sorting trays.
- collecting vials (*ca.* 10 per plot).
- spade.
- fork.
- pick axe.
- ethanol (*ca.* 0.5 L per plot).

Site information

Take and database photos of each sampling site.

Make notes of the:

- Site type (grassland, forest etc.).
- Topography.
- Date.
- Time.
- Weather conditions.
- Rainfall.
- Humidity (Lutron humidity meter).
- Ambient air temperature (Lutron air temperature meter).
- Vegetation type.
- % of bare soil.
- Condition of the site e.g. burnt or not.

Plot

- Place plot frame on selected plot.
- Photograph area.

- Search plot for casts (physical count).
- Collect casts (if needed)– label and store.

Soil sampling

- Make note of litter component of soil (depth) in the plot.
- Soil samples should be taken from inside the same plot once the earthworm sample has been removed.
- Ideally one sample per plot should be taken but due to the analysis cost, a minimum of 5 samples should be taken from a site. If the sampling site is against a slope, 5 soil samples should be taken at the top of the slope, 5 in the middle and 5 at the bottom.
- Remove plot vegetation by cutting the grass and any other vegetation to ground
- Measure soil temperature with a digital soil thermometer.
- Soil temperature to be taken at:
 - Just under the soil surface.
 - At 10 cm.
 - Between 20–30 cm.
 - At 50 cm.
- Take soil density measurements with Kingtest DCP (as prescribed by product manual) to a depth of 50 cm.
- Take soil sample with 75 mm soil auger to a depth of 50 cm.
- Take soil moisture measurements with the Lutron Soil Moisture meter from the soil sample in the soil auger.
- Moisture measurements to be taken at:
 - Just under the soil surface.
 - 0–10 cm.
 - 10–20 cm.
 - 50 cm.
- Label and store the soil sample in a paper bag for further analysis.

Plot size

- According to ISO 23611-1 plot size has been established at 50 × 50 cm. Larger plots (at least 1 × 1 m) are required for indigenous species.
- A pre-made steel frame is used to demarcate the plot area. It can be pressed into the soil if possible but should not be hammered into the soil which will cause disturbance.

Number of plots

- From our study, 12 randomly selected plots more than 5 m apart were mostly sufficient, but this will be highly context specific.

Digging and hand sorting

- Search through the litter component of the soil (2–5 cm).
- Collect earthworms, mark and record them correctly (epigeic or litter species).
- Dig out the soil in the area of the plot frame to the standard depth.
- Place the soil on a plastic sheet and search for earthworms.
- Move the sorted soil to one area for replacement into the hole.

Mustard extraction

- Remove the epigeic (topsoil) species – living and feeding on soil surface.
- Dissolve 60 g of dry mustard powder in 1 liter of distilled water. (ISO 11268-3).
- On site: Mix the mustard emulsion into 9 litres of water = 6.67 g/L.
- Pour the mixture onto the test site in 2 or 3 portions according to seepage capacity.
- The total duration of the extraction may be 30 mins.
- Allow the mustard solution to deeply seep into the ground without damming up in the plot area.
- Collect all emerging earthworms for approximately 30 mins.

Preservation of earthworms after collection

Although formalin is widely used as a fixative for biological specimens, nucleic acids extracted from formalin-preserved specimens are not available for downstream DNA use (Lehmann and Kreipe 2001, Gugic *et al.* 2007; James *et al.* 2010); therefore fixation in formalin is not recommended for molecular studies. Absolute ethanol is an excellent agent for field preservation of DNA within earthworm samples (Thakuria *et al.* 2008) and is recommended for future use in earthworm sampling in South Africa.

- Earthworms to be kept in 2 groups: (a) epigeic/litter species and (b) soil/endogeic & anecic species.
- Earthworms are washed in tap water.
- 20% ethanol is then used to kill them.
- 4% formalin is used to fix the specimens and after 24 hours transferred to 70% ethanol. Earthworms to be used for molecular studies are not fixed with formalin but are transferred straight to 100% ethanol.
- Specimens are identified in the lab.

Labelling

Important information is written on the label but more information is recorded in a field book and may be later transferred into a computerized database. Paper and pencil should be used for the labelling of samples in the field but a printed label on a good quality paper or hand written by permanent ink pen may be used for long-term storage. Laser printing is not permanent and should not be used but permanent pigment ink pens may be used. An example of a suitable label is provided below:

| |
|---|
| Museum accession no. (NMSA/OLIG.06364) |
| Country & province (RSA, KZN) |
| Locality (Queen Elizabeth Park) |
| Brief site info (grassland with few shrubs) |
| GPS coordinates (29.57148°S 30.32235°E) |
| Elevation (118 m a.s.l.) |
| Date (2 February 2012) |
| Collectors (JD Plisko, A. Mamlala leg.) |
| Person who classified (JDP det.) |

SUPPLEMENTARY MATERIAL 2

Species richness estimates for the sites from the data collected by the method producing the highest number of species: Cedara, Queen Elizabeth Park and Karkloof via digging [A to C] and Botanic Gardens via the mustard solution method [D]).

The analyses were carried out in EstimateS (Colwell 2013) using the default settings. Only specimens that could be identified to species level were included in the analyses. S(est) = estimated number of earthworm species (95% confidence intervals (CI) for S(est) indicated) using rarefaction; ACE = Abundance Coverage-based Estimator (Chazdon *et al.* 1998). The CIs are wider where species richness is higher (Karkloof and Botanic Garden) and do not converge on identified species richness. This, together with the relatively higher ACE mean estimates, indicates undersampling of species.

