

## **Change in the Distribution of National Bird (Himalayan Monal) Habitat in Gandaki River Basin, Central Himalayas**

Authors: Rai, Raju, Paudel, Basanta, Changjun, Gu, and Khanal, Narendra Raj

Source: Journal of Resources and Ecology, 11(2) : 223-231

Published By: Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences

URL: <https://doi.org/10.5814/j.issn.1674-764x.2020.02.010>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

J. Resour. Ecol. 2020 11(2): 223-231  
DOI: 10.5814/j.issn.1674-764x.2020.02.010  
www.jorae.cn

# Change in the Distribution of National Bird (Himalayan Monal) Habitat in Gandaki River Basin, Central Himalayas

Raju RAI<sup>1,2</sup>, Basanta PAUDEL<sup>1</sup>, GU Changjun<sup>1,2</sup>, Narendra Raj KHANAL<sup>1</sup>

1. Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China;

2. College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

**Abstract:** Gandaki River Basin (GRB) is part of the central Himalayan region, which provides habitat for numerous wild species. However, due to changes in climate and land cover, the habitats of many protected species are at risk. Based on the maximum entropy (MaxEnt) model, coupled with bioclimatic layers, land cover and DEM data, the impacts of environmental factors on habitat suitability of Himalayan Monal (*Lophophorus impejanus*), a national bird of Nepal, was quantified. This study further assessed the present and future habitat and distribution of the Himalayan Monal in the context of climate and land cover changes. The results of this study show that the highly suitable habitat of Himalayan Monal presently occupies around 749 km<sup>2</sup> within the northern, eastern and western parts, particularly protected areas such as Langtang National Park, Manaslu Conservation Area and Annapurna Conservation Area, while it is likely to decrease to 561 km<sup>2</sup> by 2050, primarily in the northern and northwestern parts (i.e., Chhyo, Tatopani, Humde and Chame). These expected changes indicate increasing risk for Himalayan Monal due to a decline in its suitable habitat area.

**Key words:** climate change; land cover; Himalayan Monal; habitat change; Gandaki River Basin

## 1 Introduction

The International Panel on Climate Change (IPCC) has reported that the global temperature is likely to increase by about 1.5°C between 2030 and 2052 (IPCC, 2018). This global increase in temperature will obviously have impacts on many wild species of the Nepal Himalayas. A study has revealed that species habitats are gradually migrating northward due to the rising temperatures in Nepal (Karki et al., 2009). In addition to climate change, land cover changes also affect many protected species globally (Thuiller, 2003; Jetz et al., 2007; Hofmeister et al., 2010; Grimmer et al., 2016). The increasing temperature has resulted in the shifting of vegetation to higher elevations which poses threats to high altitude endemic species that are likely to be reduced by around 77% by 2100 in the Austrian Alps (Dirnböck et al., 2011) and the Himalayan region

(Forrest et al., 2012; Chhetri et al., 2018). In addition to climate change, declining forest cover is also a fundamental cause of habitat losses and one study has projected that around 42% of forest is likely to be impacted in the Asian tropical region by 2100 (Sodhi et al., 2004). Thus, this region is likely to experience impacts on many native habitats in the future which will need conservation efforts in due time (Sodhi et al., 2004; Sodhi et al., 2010). A previous study has predicted that many Mountain species might be threatened with large ecological range reductions in the future, due to natural and anthropogenic factors (del Rosario Avalos and Hernández, 2015).

The changes in climate have exacerbated habitat loss and fragmentation in different ecological regions globally; and these changes have already impacted 54% of important terrestrial species during the 21st century (Segan et al., 2016).

**Received:** 2019-10-25 **Accepted:** 2020-01-20

**Foundation:** Chinese Academy of Sciences-The World Academy of Sciences (CAS-TWAS) President's Fellowship Program for PhD Study.

**First author:** Raju RAI, E-mail: rairaju@igsnr.ac.cn

**Citation:** Raju RAI, Basanta PAUDEL, GU Changjun, et al. 2020. Change in the Distribution of National Bird (Himalayan Monal) Habitat in Gandaki River Basin, Central Himalayas. *Journal of Resources and Ecology*, 11(2): 223–231.

One study has projected a reduction between 91% and 100% in the geographic distributions of endemic Andean bird species that currently extend from central Bolivia to southern Peru (del Rosario Avalos and Hernández, 2015). Likewise, a previous study has estimated that land cover dynamics are likely to affect around 400 species of land birds, out of the total of 8750 species, by the year 2050 in the tropical regions of the world (Jetz et al., 2007). Since historical periods, land cover dynamics have been known as a fundamental component of environmental change (Ganasri et al., 2013; Li et al., 2016), and they also play crucial roles in climate change (Pielke et al., 2002). Climate change will impact other important birds and their potential habitat distributions in future. For example, black woodpecker and boreal, tawny and Ural owls are all likely to change their geographic distributions due to changes in climate and forest cover (Brambilla et al., 2020). The changing patterns in climate and land cover, mainly decreases in vegetation, as well as illegal hunting, are already noted as the main threats to habitat protection in Mountain regions (Bhattacharya et al., 2009; Jnawali et al., 2011; Liu et al., 2017).

The Himalayan region is a shelter for numerous species (Liu et al., 2017; Nie et al., 2017). Gandaki River Basin (GRB) is a part of the central Himalayan region located in central Nepal that provides habitat for many kinds of birds, including the national bird: Himalayan Monal (HM). Among the various birds of Nepal, the HM is well-known to the Nepalese people. It is legally protected by the government and is also listed on CITES Appendix I, nationally categorized as near threatened (Inskipp et al., 2016) and listed in IUCN's least concern category (Inskipp and Baral,

2013). In addition, the HM was the first species recorded by Colonel Fitzpatrick in Nepal in 1793 (Inskipp and Baral, 2013). The higher Himalayan region, including the northern part of Gandaki Basin, has been experiencing a greater rate of temperature increase and a consequent change in the habitat of many threatened birds, including HM. However, the change in the distribution of suitable habitat for HM in the context of climate and land cover change is poorly understood. In order to address this research gap, this study assesses the current distribution and potential impacts of climate and land cover changes on the suitable habitat of HM within the GRB.

## 2 Materials and methods

### 2.1 Selection of the study area

GRB is located between 28.35°N and 29.33°N and between 82.87°E and 85.80°E. The basin covers around 32807 km<sup>2</sup> of the central Himalayan region (Fig. 1). The elevation ranges from 94 m above mean sea level (amsl) in the south to 8167 m amsl in the north. The sum of annual precipitation in GRB was recorded in 2014 as 285 mm (driest region) and 5160 mm (wettest region), and the average temperature was between 6.12 °C (lowest) and 32.35 °C (highest) (MPE, 2018). Due to heterogeneous landforms and altitudinal variations, the GRB contains various land covers and ecosystems (Rai et al., 2018). The predominant land covers are agricultural land and forest cover, which are mainly distributed in the central and southern parts of the GRB. The Annapurna Conservation Area, Manaslu Conservation Area, Chitwan National Park, Shivapuri Nagarjuna and Parsa Wildlife are located within the GRB, where large numbers




Fig. 1 Location of the GRB and species occurrences

of mammals, birds, and reptiles are found (DNPWC, 2018). The Himalayan Monal, also called Danfe, is the national bird of Nepal which has given it a very high profile (DNPWC and BCN, 2018), and it is distributed in higher elevations, particularly in Langtang National Park, Manaslu Conservation Area and Annapurna Conservation Area in the GRB (BirdLife International, 2016). According to the National Parks and Wildlife Conservation Act 1973, HM is included with eight other species of birds as protected species (Baral, 2009).

## 2.2 Datasets

### 2.2.1 Species occurrence

The points of known HM occurrences were extracted from the Global Biodiversity Information Facility (GBIF) portal (<http://www.gbif.org/>), a platform which provides more than 280 million records of different species observations worldwide (Telenius, 2011). GBIF is the largest global databank of biodiversity for scientific research (Beck et al., 2013; Robertson et al., 2014; Liu et al., 2017).

### 2.2.2 Bioclimatic variables

The bioclimatic variables (at 30 arc-second spatial resolution) were extracted from WorldClim: <http://www.worldclim.org/> (Table 1). The current WorldClim data layer was generated by the interpolation of average monthly climate data annually

for the period between 1970 and 2000 on a 30 arc-second resolution grid. The future climatic model used was the IPCC5-Global Climate Models (GCM) for four representative concentration pathways (RCP). Downscaled Coupled Model Intercomparison Project phase 5 (CMIP5) data at 30 arc-second resolution, based on Representative Concentration Pathways (RCP) 4.5 and the Community Climate System Model version 4 (CCSM4) for 2050, were used in this study. Values for each climatic variable were extracted for the corresponding species occurrence locations and used to perform the correlation analysis. Highly correlated variables were removed according to their Pearson correlation coefficients. Variables with a coefficient greater than 0.60 were excluded from the modeling.

### 2.2.3 Land cover data

National land cover data for 1990 were download from the International Centre for Integrated Mountain Development (ICIMOD) (<http://rds.icimod.org>). Similarly, land cover data for 2010 were extracted from Uddin et al. (2015). Based on the 1990 and 2010 data, land cover data for 2050 were simulated using a Cellular Automata (CA) Markov model in TerraSet version 18.21. The Markov model is often used for projecting future land cover changes (Mandal, 2014; Mondal et al., 2016). The Markov model was introduced by Brown (1963) for the projection of future land use change (Mandal, 2014). Since the model was introduced, numerous scholars have used it at different spatio-temporal scales for the prediction of land use changes. The CA-Markov model can predict both spatial and temporal changes in land use and land cover over an area (Ye and Bai, 2008; Zhao et al., 2017). It is a convenient tool for simulating land use and land cover changes (LUCC), which describes the LUCC from one period to another (in this case 1990 to 2010) and uses this as the basis for projecting future changes (in this case 2050) (Kumar et al., 2014). A digital elevation model (DEM) at 30 m spatial resolution was download from the United States Geological Survey Earth Explorer. The slope map was prepared from the DEM using ArcGIS tools.

## 2.3 Maximum Entropy (MaxEnt) model

The freely available MaxEnt version 3.4.1 software was used in this study. This model identifies wild species' environmental requirements and geographical distributions (Phillips et al., 2006; Baldwin, 2009) which are then used to estimate current and future distributions (Phillips et al., 2004; Phillips et al., 2006). The MaxEnt model is a simple and precise mathematical formulation that incorporates a number of features for estimating the geographic distribution of suitable habitat for a given species (Phillips et al., 2006). This approach has been applied across many disciplines including studies of cropland changes (Heumann et al., 2011; Gu et al., 2018) and infectious disease habitat suitability (Mweya et al., 2016; Acharya et al., 2018). In-

Table 1 Environmental variables and their descriptions

Variable	Description	Retained or excluded
Bio 1	Annual mean temperature	✓
Bio 2	Mean diurnal range	✓
Bio 3	Isothermality	✓
Bio 4	Temperature seasonality	✓
Bio 5	Max temperature of warmest month	✓
Bio 6	Min temperature of coldest month	✓
Bio 7	Temperature annual range (Bio 5, 6)	×
Bio 8	Mean temperature of wettest quarter	✓
Bio 9	Mean temperature of driest quarter	×
Climate Bio 10	Mean temperature of warmest quarter	✓
Bio 11	Mean temperature of coldest quarter	×
Bio 12	Annual precipitation	✓
Bio 13	Precipitation of wettest month	×
Bio 14	Precipitation of driest month	✓
Bio 15	Precipitation seasonality	✓
Bio 16	Precipitation of wettest quarter	✓
Bio 17	Precipitation of driest quarter	✓
Bio 18	Precipitation of warmest quarter	✓
Bio 19	Precipitation of coldest quarter	✓
Topography	Elevation, slope, aspect	✓
Land cover	Land cover data at 30 m spatial resolution	✓

Note: Abbreviations: ✓ means retained; × means excluded.

deed, more than 1000 studies have applied this model since 2006 because of its high predictive performance (Merow et al., 2013). Several existing studies also have recommended this model for predicting the future habitat of protected species under different environmental scenarios (Kumar and Stohlgren, 2009; Larson et al., 2010; Zhang et al., 2011; Remya et al., 2015; Liu et al., 2017). Species occurrences, climatic variables, land cover, DEM and slope were used as the source input data for the model. The results from the MaxEnt model were imported to ArcGIS to analyze the current and future habitat distribution of HM.

This study categorized habitat suitability by defining five classes: Very high suitability (greater than 70% probability); High suitability (from 50% to 70% probability); Medium suitability (from 30% to 50% probability); Low suitability (from 10% and 30% probability); and Very low suitability (less than 10% probability) (Liu et al., 2017).

3 Results

3.1 Land cover changes

Based on land cover data for 1990 and 2010, the CA Markov model was used to predict land cover data for 2050. This model predicts that agricultural land, shrubland and built-up areas are likely to increase in the future, while forest cover, grassland, barren land and water body area would decrease by 2050 (Table 2). It further predicts that largely occupied forest cover is likely to decrease from 40% to 35% between 2010 and 2050, while shrubland and agricultural land are likely to increase by about 1% and 2%, respectively (Table 2 and Fig. 2).

The results show that the predominant forest cover is likely to decrease in each of the elevation ranges, while the shrubland is likely to increase at elevations up to 3750 m (Fig. 3). Likewise, the grassland is likely to decrease between 3500 m and 6500 m elevation ranges, while the agricultural land is more likely to increase in each of the elevation ranges from 250 m to 3250 m within the GRB (Fig. 3).

Table 2 Current (2010) and projected (2050) land use and land cover (LULC) types

LULC types	2010		2050	
	Area (km <sup>2</sup> )	Percentage (%)	Area (km <sup>2</sup> )	Percentage (%)
Forest	13174	40.15	11344	34.58
Shrubland	821	2.50	1072	3.27
Grassland	3798	11.58	3366	10.26
Agricultural land	7574	23.09	8379	25.54
Barren land	3284	10.01	3147	9.59
Water body	168	0.51	115	0.35
Snow/Glacier	3887	11.85	5218	15.91
Built-up area	101	0.31	166	0.51
Total	32807	100	32807	100

3.2 Model evaluation and analysis of variables

In setting-up the MaxEnt model, 25% of the occurrence locations were used for testing and the remaining randomly selected 75% were used to train the model. This process found the area under the Receiver Operating Characteristics (ROC) curve, or mean Area Under ROC (AUC) test set data, to be 0.920. The overall accuracy of the model performance was high, which indicates that the distributions generated provide close estimations for the real-world distribution probabilities.

The environmental variables are fundamental to MaxEnt modeling and each one contributes to the spatial distribution of habitat. Different environmental variables play important roles in explaining whether or not the species exists in a particular area. Thus, if any environmental layer provides a higher contribution, then it represents a higher impact of the variables on the habitat prediction. The result shows that climatic layer Bio 8 (mean temperature of wettest quarter) provided the highest contribution to the HM, corresponding to 29%. Two others, Bio 14 (precipitation of driest month)




Fig. 2 Current (a, 2010) and projected (b, 2050) LULC






Fig. 3 Land cover changes along the altitudinal gradient

(12%) and elevation (9%) were also found to be important variables contributing to the habitat of HM (Fig. 4).

3.3 Habitat status and future distributions

Under the appropriate conditions of the climatic variables, topography and land cover, the modeling results reveal the suitable habitat across the GRB. The total area of HM habitat is projected to increase by 695 km<sup>2</sup> in the future, with corresponding decreases and stable areas projected to be 3177 km<sup>2</sup> and 995 km<sup>2</sup>, respectively, in the future (Fig. 5).




Fig. 4 Environmental variables and contributions  
Note: See Table 1 for descriptions of the variables.




Fig. 5 Projected habitat changes of HM within the GRB in the future (by 2050)

Most of the suitable habitat increase would be in the eastern parts such as Nyak, Langtang National Park and Manaslu Conservation Area, while in many places in the central and western parts such as Chame, Chhyo, Tatopani, and Humde, habitat areas are likely to decrease by 2050.

This study assessed the habitat suitability according to five classes: very high suitability (>70% probability); high suitability (50%–70% probability); medium suitability (30%–50% probability); low suitability (10%–30% probability); and very low suitability (<10% of probability) (Liu et al., 2017). The very high suitability habitat of the species is projected to decrease in the future at different spatial scales, while the high suitability habitat is likely to decrease around 188 km<sup>2</sup> in the future. Similarly, medium suitability habitats are also predicted to decrease in total area (Table 3 and Fig. 6).

3.4 Habitat changes along the altitudinal gradient

This study assessed the very high suitability area of greater than 70% prediction probability as it was mapped at different altitudes in the future. The habitat of HM is likely to decrease at elevations between 1750 m and 3750 m, and between 4500 m and 5500 m, while at elevation ranges from 3750 to 4500 m it will likely increase in the future. Overall, the projections found around 186 km<sup>2</sup> of total area loss within these elevation ranges (Fig. 7).

4 Discussion: The potential impact of climate and land cover changes on habitat of HM

Currently, the HM is estimated to occur within less than 20000 km<sup>2</sup> area globally, including Nepal, India, Bhutan, Afghanistan, Pakistan, Tibet and Myanmar (Baral, 2009;

Table 3 The suitability status of current and predicted future habitat areas (km<sup>2</sup>)

Habitat suitability	Current	Future
Very high	748.50	560.54
High	1532.99	846.76
Medium	2625.70	1849.90
Low	4520.88	9201.65
Very low	23379.74	20349.06




Fig. 6 Current (a) and predicted (b) habitat distributions in the GRB




Fig. 7 Changes in very high suitability habitat at different elevation ranges in the GRB

Miller, 2010; BirdLife International, 2016). It is found at high altitudes, in places with steep slopes, rocky slopes, cliffs, grasslands and wood patches (Inskipp et al., 2016; Inskipp and Inskipp, 1991). The total global population has not been calculated, though in Nepal it is estimated at around 3500 to 5000 individuals (Inskipp et al., 2016). Still, the total population within the GRB is unknown, though one previous study reported 26 individuals in winter and 51 in spring within the Annapurna Conservation Area (BCN, 2013). However, hunting and trapping have been heavily practiced by local hunters, herders and medicinal plant collectors in the Mountain region of the country for many years (Baral, 2009; Miller, 2010; Inskipp and Baral, 2013; Inskipp et al., 2016). This bird tends to move into surrounding farmlands where people kill it for its crest feathers as well as its meat, which reaches a high price (Kaul et al., 2004; Ma et al., 2011; Inskipp et al., 2016). Changes in forest cover, grassland, and human activities, as well as illegal activities, threaten the continuing presence of this bird in Nepal (BCN and DNPWC, 2011; Inskipp et al., 2016). The grassland area has a declining trend due to overgrazing and burning, such that no significant area remains outside of the protected areas in the country (Inskipp et al., 2016). The human disturbance, particularly livestock grazing, and natural disturbances were also noted as the key causes of

declining habitat in the Great Himalayan National Park, India (Miller, 2010). One recent study has observed that due to livestock grazing and large-scale collection of Mushrooms from the forest, many species are experiencing degraded situations in the western Himalayan region of Pakistan (Ahmad et al., 2019). The results of this study also project a decrease in the forest area by approximately 5% between 2010 and 2050. Nationally, the forest cover has shown a decreasing trend between 1930 to 2014, and a net loss of 37318 km<sup>2</sup> across the country has been observed (Reddy et al., 2018). Decreases in forest cover and grasslands are likely to reduce the habitat for HM within the GRB.

The habitat of HM is also projected to decrease at elevations between 1750 m and 3750 m, and to increase at elevations between 3750 m and 4500 m. Past studies also found the suitable habitat ranges between 2500 m and 4750 m in barren land and open forest of the Nepal Himalayan regions (BCN, 2013). The model used in this study found that a suitable habitat range might be at elevations between 3750 m and 4500 m. These elevation ranges encompass estimated decreases in forest cover, grassland and shrubland, which could influence these habitat ranges in the future. Land cover and changing climatic patterns determine future habitat changes for this species. Highly significant climatic variables, as well as elevation, are expected to be favorable for HM in the northeastern region. Habitat is projected to likely increase in the eastern part of LNP and ACA buffer zones, as well as in the MCA and surrounding protected areas that are especially close to agricultural land.

This species undergoes fluctuations with altitudinal and seasonal migrations (Inskipp et al., 2016). This study has projected a small increase in the elevation range between 2000 m and 2250 m, and the increase in this range could be due to the seasonal migration of HM. This species migrates with seasons mainly during winter, and moves to lower elevations, around 2500 m (Baral, 2009), bringing it close to human habitations and people hunting it for local consumption in the western Himalaya, India (WII, 2016). Similarly, during the spring and autumn seasons, the Himalayan

Monal shares habitat with herded livestock, which is also associated with negative factors for species occurrence, mainly shepherd dogs and human disturbances which have negatively impacted the MH in the western Himalayan region, India (Bhattacharya et al., 2009). During the summer season, the HM shifts up to 4300 m elevation and digs roots by using its strong bill (Ma et al., 2011). Therefore, this study has estimated that the mean temperature of wettest seasons will have a high impact on habitat suitability of the HM in the future within the GRB.

## 5 Conclusions

This study assessed the current and future distributions of the habitat of the nationally protected bird, Himalayan Monal, within the GRB using the MaxEnt model. Under the conditions of changing climatic variables, land cover changes were found likely to impact the habitat of the Himalayan Monal in 2050. The highly suitable habitat of the Himalayan Monal is likely to decrease by around 188 km<sup>2</sup> in the future, particularly in the northern and northwestern parts (i.e., Chhyo, Tatopani, Humde and Chame). For the most part, the suitable habitat would increase in the eastern parts such as Nyak, Langtang, Langtang National Park and Manaslu Conservation Area in 2050. This study highlights the fact that more attention should be given to HM, in terms of future habitat protection. This species is expected to experience considerable decreases in suitable habitat areas in the future. The northeast parts are particularly important regions and more attention should be paid to them, especially with regard to forest cover, grassland, and shrubland to protect the remaining habitat of HM.

## References

- Acharya B, Cao C, Xu M, et al. 2018. Present and future of dengue fever in Nepal: Mapping climatic suitability by ecological niche model. *International Journal of Environmental Research and Public Health*, 15(2): 1-16.
- Ahmad B, Noor F, Awan M, et al. 2019. Distribution and population status of Himalayan Monal pheasant (*Lophophorus impejanus*) in Salkhala game reserve, Neelum valley Azad Jammu and Kashmir (Pakistan). *The Journal of Animal Plant Science*, 29(4): 1150-1159.
- Baldwin R. 2009. Use of maximum entropy modeling in wildlife research. *Entropy*, 11(4): 854-866.
- Baral H. 2009. Protected birds of Nepal: A review of their status, distribution and habitat. *The Initiation*, 9: 66-80.
- BCN. 2013. Important bird areas in Nepal deliver vital ecosystem services to people. Kathmandu: Bird Conservation Nepal.
- BCN, DNPWC. 2011. The state of Nepal's Birds 2010. Kathmandu: Bird Conservation Nepal and Department of National Parks and Wildlife Conservation.
- Beck J, Ballesteros-Mejia L, Nagel P, et al. 2013. Online solutions and the 'W allacean shortfall': What does GBIF contribute to our knowledge of species' ranges? *Diversity and Distributions*, 19: 1043-1050.
- Bhattacharya T, Sathyakumar S, Rawat G. 2009. Distribution and abundance of Galliformes in response to anthropogenic pressures in the buffer zone of Nanda Devi Biosphere Reserve. *International Journal of Galliformes Conservation*, 1: 78-84.
- BirdLife International. 2016. *Lophophorus impejanus*. The IUCN Red List of Threatened Species 2016: e.T22679182A92806166. <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22679182A92806166.en>.
- Brambilla M, Scridel D, Bazzi G, et al. 2020. Species interactions and climate change: How the disruption of species co-occurrence will impact on an avian forest guild. *Global Change Biology*, 26(3): 1212-1224.
- Chhetri P K, Gaddis K D, Cairns D M. 2018. Predicting the suitable habitat of treeline species in the Nepalese Himalayas under climate change. *Mountain Research and Development*, 38(2): 153-163.
- del Rosario Avalos V, Hernández J. 2015. Projected distribution n shifts and protected area coverage of range-restricted Andean birds under climate change. *Global Ecology and Conservation*, 4: 459-469.
- Dirnböck T, Essl F, Rabitsch W. 2011. Disproportional risk for habitat loss of high-altitude endemic species under climate change. *Global Change Biology*, 17(2): 990-996.
- DNPWC. 2018. Protected areas of Nepal. Department of National Parks and Wildlife Conservation, Ministry of Forests and Soil Conservation, Kathmandu, Nepal.
- DNPWC, BCN. 2018. Birds of Nepal: An official checklist. Department of National Parks and Wildlife Conservation and Bird Conservation Kathmandu, Nepal.
- Forrest J L, Wikramanayake E, Shrestha R, et al. 2012. Conservation and climate change: Assessing the vulnerability of snow leopard habitat to treeline shift in the Himalaya. *Biological Conservation*, 150(1): 129-135.
- Ganasri B, Raju A, Dwarakish G. 2013. Different approaches for land use land cover change detection: A review. *Journal of Engineering and Technology*, 2: 44-48.
- Grimmett R, Inskipp C, Inskipp T, et al. 2016. Birds of Nepal (revised edition). Christopher Helm, London, UK: Bloomsbury Publishing.
- Gu C, Zhang Y, Liu L, et al. 2018. Comprehensive evaluation of the suitability of agricultural land in Myanmar. *Journal of Resources and Ecology*, 9(6): 609-622.
- Heumann B W, Walsh S J, McDaniel P M. 2011. Assessing the application of a geographic presence-only model for land suitability mapping. *Ecological informatics*, 6(5): 257-269.
- Hofmeister E, Rogall G M, Wesenberg K, et al. 2010. Climate change and wildlife health: Direct and indirect effects. Rep. No. 2327-6932. US Geological Survey.
- Inskipp C, Baral H. 2013. Danphe—A nationally threatened species? In: Danphe, Bird Conservation Nepal, 22: 1-12.
- Inskipp C, Baral H, Phuyal S, et al. 2016. The status of Nepal's birds: The national red list series. Vol. 1, Zoological Society of London, UK.
- Inskipp C, Inskipp T. 1991. A guide to the birds of Nepal (Second edition), London, UK: Christopher Helm.
- IPCC. 2018. Summary for policymakers. In: Masson-Delmotte V, Zhai P, Pörtner H O, et al. (eds.). Global warming of 1.5°C. An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty. Geneva, Switzerland: World Meteorological Organization.
- Jetz W, Wilcove D S, Dobson A P. 2007. Projected impacts of climate and land-use change on the global diversity of birds. *Plos Biology*, 5(6): 1211-1219.
- Jnawali S, Baral H, Lee S, et al. 2011. The status of Nepal mammals: The national red list series 4. Department of National Parks and Wildlife Conservation Kathmandu, Nepal.
- Karki M, Mool P, Shrestha A. 2009. Climate change and its increasing impacts in Nepal. *The Initiation*, 3: 30-37.



- Kaul R, Jandrotia J, McGowan P J. 2004. Hunting of large mammals and pheasants in the Indian western Himalaya. *Oryx*, 38(4): 426–431.
- Kumar S, Radhakrishnan N, Mathew S. 2014. Land use change modelling using a Markov model and remote sensing. *Geomatics, Natural Hazards and Risk*, 5(2): 145–156.
- Kumar S, Stohlgren T J. 2009. Maxent modeling for predicting suitable habitat for threatened and endangered tree *Canacomyrica monticola* in New Caledonia. *Journal of Ecology and the Natural Environment*, 1: 94–98.
- Larson S R, Degroot J P, Bartholomay L C, et al. 2010. Ecological niche modeling of potential west Nile virus vector mosquito species in Iowa. *Journal of Insect Science*, 10(1): 1–17.
- Li X, Wang Y, Li J, et al. 2016. Physical and socioeconomic driving forces of land-use and land-cover changes: A case study of Wuhan City, China. *Discrete Dynamics in Nature and Society*, 2016: 1–11. DOI: 10.1155/2016/8061069.
- Liu L, Zhao Z, Zhang Y, et al. 2017. Using MaxEnt model to predict suitable habitat changes for key protected species in Koshi basin, central Himalayas. *Journal of Resources and Ecology*, 8(1): 77–87.
- Ma X, Guo J, Yu X. 2011. Himalayan Monal (*Lophophorus impejanus*): Distribution, habitat and population status in Tibet, China. *Chinese Birds*, 2(3): 157–162.
- Mandal U K. 2014. Geo-information based spatio-temporal modeling of urban land use and land cover change in Butwal municipality, Nepal. *International Archives of Photogrammetry, Remote Sensing and Spatial Information Sciences*, XL-8(1): 809–815.
- Merow C, Smith M J, Silander J A. 2013. A practical guide to MaxEnt for modeling species' distributions: What it does, and why inputs and settings matter. *Ecography*, 36: 1058–1069.
- Miller J R. 2010. Survey of Western Tragopan, Koklass Pheasant, and Himalayan Monal populations in the Great Himalayan National Park, Himachal Pradesh, India. *Indian Birds*, 6: 60–65.
- Mondal M S, Sharma N, Garg P, et al. 2016. Statistical independence test and validation of CA Markov land use land cover (LULC) prediction results. *The Egyptian Journal of Remote Sensing and Space Science*, 19: 259–272.
- MPE. 2018. Weather summary of Nepal year–2014. Government of Nepal, Ministry of Population and Environment, Department of Hydrology and Meteorology, Kathmandu.
- Mweya C N, Kimera S I, Misinzo G, et al. 2016. Climate change influences potential distribution of infected *Aedes aegypti* co-occurrence with dengue epidemics risk areas in Tanzania. *Plos One*, 11: 1–13.
- Nie Y, Sheng Y, Liu Q, et al. 2017. A regional-scale assessment of Himalayan glacial lake changes using satellite observations from 1990 to 2015. *Remote Sensing of Environment*, 189: 1–13.
- Phillips S J, Anderson R P, Schapire R E. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190: 231–259.
- Phillips S J, Dudík M, Schapire R E. 2004. A maximum entropy approach to species distribution modeling. In: Proceedings of the twenty-first international conference on Machine learning, ACM, Banff, Canada.
- Pielke R A, Marland G, Betts R A, et al. 2002. The influence of land-use change and landscape dynamics on the climate system: Relevance to climate-change policy beyond the radiative effect of greenhouse gases. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 360: 1705–1719.
- Rai R, Zhang Y, Paudel B, et al. 2018. Land use and land cover dynamics and assessing the ecosystem service values in the Trans-Boundary Gandaki River Basin, Central Himalayas. *Sustainability*, 10: 1–22.
- Reddy C S, Pasha S V, Satish K, et al. 2018. Quantifying nationwide land cover and historical changes in forests of Nepal (1930–2014): Implications on forest fragmentation. *Biodiversity and Conservation*, 27: 91–107.
- Remya K, Ramachandran A, Jayakumar S. 2015. Predicting the current and future suitable habitat distribution of *Myristica dactyloides* Gaertn using MaxEnt model in the Eastern Ghats, India. *Ecological Engineering*, 82: 184–188.
- Robertson T, Döring M, Guralnick R, et al. 2014. The GBIF integrated publishing toolkit: Facilitating the efficient publishing of biodiversity data on the internet. *Plos One*, 9(8): 1–7.
- Segan D B, Murray K A, Watson J E. 2016. A global assessment of current and future biodiversity vulnerability to habitat loss–climate change interactions. *Global Ecology and Conservation*, 5: 12–21.
- Sodhi N S, Koh L P, Brook B W, et al. 2004. Southeast Asian biodiversity: An impending disaster. *Trends in Ecology & Evolution*, 19: 654–660.
- Sodhi N S, Posa M R C, Lee T M, et al. 2010. The state and conservation of Southeast Asian biodiversity. *Biodiversity and Conservation*, 19: 317–328.
- Telenius A. 2011. Biodiversity information goes public: GBIF at your service. *Nordic Journal of Botany*, 29: 378–381.
- Thuiller W. 2003. BIOMOD—Optimizing predictions of species distributions and projecting potential future shifts under global change. *Global Change Biology*, 9: 1353–1362.
- Uddin K, Shrestha H L, Murthy M, et al. 2015. Development of 2010 national land cover database for the Nepal. *Journal of Environmental Management*, 148: 82 – 90.
- WII. 2016. National studybook of Himalayan Monal (*Lophophorus impejanus*). Wildlife Institute of India, Deharadun and Central Zoo Authority, New Delhi.
- Ye B, Bai Z. 2008. Simulating land use/cover changes of Nenjiang County based on CA-Markov model. In: International Conference on Computer and Computing Technologies in Agriculture. Springer: 321–329.
- Zhang J, Zhang Y, Liu L, et al. 2011. Predicting potential distribution of Tibetan spruce (*Picea smithiana*) in Qomolangma (Mount Everest) National Nature Preserve using maximum entropy niche-based model. *Chinese Geographical Science*, 21(4): 417–426.
- Zhao Z, Wu X, Zhang Y, et al. 2017. Assessment of changes in the value of ecosystem services in the Koshi River Basin, central high Himalayas based on land cover changes and the CA-Markov model. *Journal of Resources and Ecology*, 8(1): 67–76.

## 喜马拉雅中部甘达基河流域尼泊尔国鸟（棕尾虹雉）生境分布变化

Raju RAI<sup>1,2</sup>, Basanta PAUDEL<sup>1</sup>, 谷昌军<sup>1,2</sup>, Narendra Raj KHANAL<sup>1</sup>

1. 中国科学院地理科学与资源研究所, 陆地表层格局与模拟重点实验室, 北京 100101;

2. 中国科学院大学资源与环境学院, 北京 100049

**摘 要:** 甘达基河流域 (Gandaki River Basin, GRB) 是喜马拉雅中部地区的一部分, 该地区栖息着许多珍稀的野生动物。由于气候和人类活动的影响, 许多珍稀保护物种的生境处于危险之中。本研究基于最大熵 (MaxEnt) 模型, 运用生物气候、土地覆被和 DEM 数据, 分析各环境要素对棕尾虹雉 (*Lophophorus impejanus*) 的生境适宜性的影响, 评估棕尾虹雉现在状况和未来栖息地分布的变化。研究表明, 目前棕尾虹雉的高度适宜栖息地面积约为 749 km<sup>2</sup>, 主要分布在流域北部、东部和西部, 尤其是郎塘国家公园、马纳斯卢峰自然保护区和安娜布尔纳峰自然保护区等保护区内。到 2050 年, 棕尾虹雉的高度适宜栖息地面积将减少至 561 km<sup>2</sup>, 主要在流域北部和西北部 (即 Chhyo, Tatopani, Humde 和 Chame 地区)。未来环境变化的模拟表明, 由于适宜栖息地面积的减少, 棕尾虹雉面临的生存风险将增加。

**关键词:** 气候变化; 土地覆被; 棕尾虹雉; 生境变化; 甘达基河流域