

BLOOD CHARACTERISTICS OF FREE-RANGING WHITE-TAILED DEER IN SOUTHERN TEXAS

Author: WHITE, MARSHALL

Source: Journal of Wildlife Diseases, 10(1) : 18-24

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-10.1.18>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BLOOD CHARACTERISTICS OF FREE-RANGING WHITE-TAILED DEER IN SOUTHERN TEXAS

MARSHALL WHITE*, Department of Forestry and Conservation, Purdue University, Lafayette, Indiana, U.S.A.

ROBERT S. COOK**, Department of Veterinary Science, University of Wisconsin, Madison, Wisconsin, U.S.A.

Abstract: We present an analysis of some blood characters in 31 newborn fawns and 79 older white-tailed deer (*Odocoileus virginianus*) from free-ranging populations in southern Texas. In the newborn fawns, averages were: Erythrocytes $6.95 \times 10^6/\text{mm}^3$, leucocytes $3.6 \times 10^3/\text{mm}^3$, hemoglobin 10.1 g/100 ml, sedimentation rate 2.2 mm/hr and hematocrit 29.1%. In the older fawns and adults averages were: Hematocrit 49.5%, serum calcium 5.2 mEq/L, serum sodium 157.6 mEq/L, serum potassium 6.8 mEq/L, BUN 14.4 mg%, total serum protein 6.4 g/100 ml, albumin 3.8 g/100 ml, globulins 2.6 g/100 ml, and albumin/globulin ratio 1.5. These values for fawns and older deer generally fell within the ranges reported in other studies. More data are needed, particularly from free-ranging populations, in order to determine which blood composition characters can be used efficiently to indicate subtle differences in health and condition of deer.

INTRODUCTION

The blood of mammals has a remarkable tendency to maintain constancy of composition during good health.¹⁸ Data on blood characteristics and composition from wildlife populations reportedly are helpful in contrasting different species and comparing health and disease status among populations of the same species.^{18,20}

We describe the blood of newborn fawns and older white-tailed deer from two populations in southern Texas. Samples of newborn fawn blood came from the Welder Wildlife Refuge, near Sinton, in San Patricio County. Samples of blood of older deer came from the Welder refuge and from the Aransas National Wildlife Refuge near Austwell, about 40 km to the northeast. Both populations are free-ranging, dense, and healthy.^{8,27,28}

There are few reports in the literature of blood analyses from free-ranging whitetail populations, particularly of newborn fawns. Seal and Erickson¹⁷ described the blood of free-ranging white-tails in Minnesota. Johnson et al.¹³ presented blood characteristics for a large sample of penned whitetails in Michigan that included newborn fawns. Youatt et al.³⁰ gave blood values for captive newborn fawns and their postpartum dams. Dommert et al.¹¹ and Tumbleson et al.^{22,28} analyzed the blood of penned whitetails in Missouri. Cowan and his colleagues^{4,10,14} in British Columbia described blood characteristics of various races of mule deer (*Odocoileus hemionus*), particularly the blacktail (*O. h. columbianus*). Theirs¹⁰ is the only large sample of free-ranging newborn fawns we find described in the literature. Rosen

* Presently, School of Forestry and Conservation, and Museum of Vertebrate Zoology, University of California, Berkeley, 94720, U.S.A.

**Presently College of Environmental Sciences, University of Wisconsin—Green Bay 54302, U.S.A.

and Bischoff¹⁶ in California, Browman and Sears⁷ and Taber et al.²⁰ in Montana, and Anderson et al.^{1,2,3} in Colorado also have presented the results of larger samples of blood of free-ranging mule deer.

MATERIAL AND METHODS

In June of 1965, 10 ml of blood were drawn from the jugular vein of newborn fawns, 1-14 days of age.⁵ Between September, 1964 and May, 1965, blood samples were taken from a total of 79 older deer that were shot in the neck with a rifle. Within 5 minutes of death samples of blood were taken from the heart.

Ten ml aliquots of each sample were placed in vials with EDTA-K anticoagulant. Samples were cooled in the laboratory and analyzed within 6 hours. From these, cell counts, and measures of hemoglobin, hematocrit (packed cell volume), and sedimentation rate were made according to standard procedures.⁵ A Spencer BriteLine hemacytometer was used for erythrocyte and leucocyte counts, a Haden-Hausser hemoglobino-meter for hemoglobin estimation, and Wintrobé hematocrit tubes for sedimentation rates and packed cell volume.

Portions of samples from the older deer were cooled in a refrigerator and allowed to coagulate. When a clot had formed serum was decanted, centrifuged, and frozen for chemical analysis. Samples were discarded if hemolysis obviously had occurred. The analyses were conducted at the Department of Veterinary Medicine of Purdue University during the summer of 1965. Serum calcium levels were estimated by titration procedures,¹³ sodium and potassium levels by flame photometry, BUN by Unitest methods (Biodynamics Inc., Indianapolis, Ind.), a total serum protein by Unitest methods using the Weichselbaum biuret reagent, and the albumin/globulin (A-G) protein fraction separation by paper electrophoresis using the Spinco apparatus (Spinco Tech. Bull. No. 6027A).

RESULTS

Newborn Fawns

The average rectal temperature of 50 newborn fawns on the Welder refuge was 39.9°C. There was no apparent difference related to sex, or age in days. Blood cell counts, hemoglobin, sedimentation rate, and hematocrit were measured in 31 newborn fawns. A separation was made between fawns 1-3, 4-7, and 8-14 days old, but no significant differences ($P < .05$) in these blood values were observed. Composite mean values (table 1) were: Erythrocytes $6.95 \times 10^6/\text{mm}^3$, leucocytes $3.6 \times 10^3/\text{mm}^3$, hemoglobin concentration 10.1 g/100 ml, sedimentation rate 2.2 mm/hr, and hematocrit 29.1%.

Older Fawns and Adults

Blood samples were analyzed from a total of 79 older fawns and adults, 28 from the Welder refuge, and 51 from the Aransas refuge. There were 24 male deer and 55 females, 16 fawns (4-11 months), and 63 adults. All were collected from September through May. No significant differences (of $P > .05$) were observed between males and females, between these older fawns and adults, with season of collection, or between the two deer populations. Hence, all of the samples were combined (table 2).

For these older deer there was considerable variation between individuals in the blood values measured. Accordingly, ranges are provided in addition to the mean values (table 2). The mean hematocrit percentage was 49.5; of this, erythrocytes and platelets represented 49.1% and leucocytes 0.4%. Serum electrolyte values were 5.2 mEq/L (= 10.3 mg/100 ml) for calcium, 157.6 mEq/L for sodium, and 6.8 mEq/L for potassium. Serum blood urea nitrogen (BUN) levels averaged 14.4 mg%. Total serum protein averaged 6.4 g/100 ml. Of this, Albumin made up 3.8 g/100 ml (59%) and Globulins 2.6 g/100 ml (41%). The A/G ratio was 1.5 (table 2).

TABLE 1. Blood characteristics for 31 newborn fawns on the Welder refuge.

Age in Days	Sample Size	Erythrocytes (10 ⁶ /mm ³)		Leucocytes (10 ³ /mm ³)		Hemoglobin (g/100 ml)		Sedimentation Rate (mm/hr)		Hematocrit (%)	
		Mean	S.E.	Mean	S.E.	Mean	S.E.	Mean	S.E.	Mean	S.E.
1-3	14	6.79	0.32	3.7	0.5	10.2	0.2	2	2	28	0.9
4-7	11	6.89	0.45	3.9	0.6	10.1	0.1	2	2	30	1.0
8-14	6	7.45	0.13	3.1	0.5	10.0	0.1	3	3	30	2.2
Composite mean	31	6.95		3.6		10.1		2.2		29.1	

TABLE 2. Blood characteristics for 79 deer, four months and older, on the Welder and Aransas Refuges.

Parameter	Mean	Range	S.E.
Whole Blood			
Hematocrit (%)	49.5	39.4-66.1	8.2
Erythrocytes and platelets	49.1	39.2-65.8	7.8
Leucocytes	.4	.1-1.5	.1
Serum			
Ca (mEq/L)	5.2	3.8-7.0	.2
Na (mEq/L)	157.6	152-170	1.9
K (mEq/L)	6.8	4.2-9.9	.5
BUN (mg %)	14.4	4.0-23.5	2.8
Total protein (g/100 ml)	6.4	3.8-8.3	.5
Albumin	3.8	2.4-5.1	.2
Globulin	2.6	1.4-3.7	.2
A/G ratio	1.5	1.0-2.2	.1

DISCUSSION

The rectal temperature average of 39.9°C in newborn fawns was higher than the only temperatures we find reported for deer, adult whitetails,²⁰ and older mule deer fawns.⁹ Cowan and Wood⁹ found a higher rectal temperature in 4-month mule deer fawns than in adult females.

Erythrocyte, leucocyte, hemoglobin, sedimentation rate and hematocrit values for these newborn fawns were similar to those reported for penned fawns in Michigan.^{12,30} Compared to mule deer fawns,^{10,14} erythrocyte averages were higher, and leucocyte, hematocrit and sedimentation rate values were lower in these newborn whitetails.

Mean blood values for the older whitetails generally fell within the parameters reported in other studies, according to a summary comparison presented in table 3. Variation with individuals is to be expected. In a natural population a few diseased and abnormal individuals often are present. Our mean values should characterize the typical blood values; some of the range extremes doubtless represent diseased or abnormal individuals. Values of 170 mEq/L for sodium, and 9.9 mEq/L for potassium (table 2), for example, probably represent abnormal individuals. Hemolysis also can affect the values of these electrolytes, however, all obviously hemolyzed samples were discarded.

There are few divergencies in blood characteristics between males and females, or with age after the first 12 months, reported in the literature for *Odocoileus*. However, there appear to be differences in several blood values between fawns, particularly newborn fawns, and adults. Higher levels in newborn fawns for albumin in proportion to globulins,¹⁰ leucocytes,^{10,12} and in rectal temperature⁹ have been reported. Lower values in young fawns appear to occur in total serum protein, hemoglobin, erythrocytes, and hematocrit.^{10,12,28,30}

Additionally, there are differences between whitetails and mule deer. Higher values for erythrocytes,^{1,12,18,16} hemoglo-

bin^{1,12,14,16} and higher rectal temperature;^{9,28} and a lower A/G ratio⁴ have been reported in whitetails.

The effects of handling and treatment on blood values require more investigation. Restraint, handling, and immobilization with drugs can result in a hemodilution or expansion of plasma volume.¹⁸ Blood samples obtained from deer that have been shot may show similar deviations from normal values, particularly in individuals that do not die instantly.

Although blood composition is suggested as an indicator for evaluating health and condition in mammals, the reported data at hand for deer, in most cases, do not appear to be sufficient to differentiate clearly between small changes in health and condition and other possible variation. Studies of the effects of restricted diet and winter weight and condition loss on the blood of penned whitetails suggest a tendency toward hemoconcentration, but the results have been variable.^{18,21,24,25,26} Rosen and Bischoff¹⁸ found a marked drop in erythrocyte and hemoglobin concentrations, but apparently only in those mule deer herds that were near starvation. Workers in British Columbia^{4,14} reported no reductions of hematocrit, sedimentation rate, hemoglobin, or total plasma proteins in mule deer receiving a low plane of nutrition. Youatt et al.³⁰ reported no correlation between blood composition and diet in feeding experiments in Michigan. Anderson et al.^{1,2} found some differences in blood chemistry associated with changes in habitat quality. Bolte et al.⁴ observed marked reductions in hemoglobin and hematocrit levels in newborn whitetails in Oklahoma, but in individuals in very poor condition with tissue damage from infestation of ticks. More data are needed, particularly from free-ranging populations, in order to determine which blood composition characters can be used efficiently to indicate subtle differences in health and condition of deer, and to describe the range of normal variations in individuals, populations, races, and species of *Odocoileus*.

TABLE 3. Representative blood values (means) for adult deer. Adopted from various sources.

	Hematocrit (%)	Serum Ca mEq/L ^a or mg/100 ml ^b	Serum Na (mEq/L)	Serum K (mEq/L)	BUN (mg %)	Total Serum protein	% Albumin	A/G Ratio
White-tailed deer								
**Michigan ^{11,19,24,25,26}	56.3	8.8 - 9.4 ^b	143 - 148	4.1 - 4.9	17 - 24	8.1	63.6	1:1.7
	39.5 - 55.2					5.1 - 5.9	55.6 - 57.1	1:1.2 - 1:1.3
	51 - 59				16.9 - 25.6	6.4 - 10.5	55.4 - 63.6	1:1.2 - 1:1.7
*Minnesota¹⁷								
	39 - 44				12 - 16	4.9 - 5.2	64.0 - 64.2	1:1.7 - 1:1.8
**Missouri^{11,28}								
	47.5 - 54.8	10.9 - 11.6 ^b	154 - 156	6.0 - 6.4	22 - 25	7.2 - 7.6		
*Maryland²⁰								
			174	10.4				
Mule deer								
*Montana ^{7,20}	39.6					5.9 - 6.4		
**British Columbia^{4,14}								
	58.2					6.1 - 6.7	64 - 72	1:2.5
*Colorado^{1,2,8}								
	46.7	6.1 - 6.4 ^a	140 - 164	7.4 - 9.9		5.5 - 7.1		
• free-ranging deer								
•• penned deer								

Acknowledgements

This was a cooperative study supported by the Welder Wildlife Foundation (Contribution No. 153); the Department of Veterinary Science, University of Wisconsin; and the Department of Forestry and Conservation, Purdue University (AES Journal Paper No. 5148). We thank all who assisted in the field, laboratory, and office, particularly D. L. Allen, E. L. Benham, C. Cottam, B. A. Fall, C. M. Fleming, S. M. Gaafar, W. C. Glazener, K. C. Matocha, R. L. Morter, R. H. Shields, and D. O. Trainer.

LITERATURE CITED

1. ANDERSON, A. E., D. E. MEDIN and D. C. BOWDEN. 1970. Erythrocytes and leucocytes in a Colorado mule deer population. *J. Wildl. Mgmt.* 34: 389-406.
2. ANDERSON, A. E., D. E. MEDIN and D. C. BOWDEN. 1972. Blood serum electrolytes in a Colorado mule deer population. *J. Wildl. Dis.* 8: 183-190.
3. ANDERSON, A. E., D. E. MEDIN and D. C. BOWDEN. 1972. Total serum protein in a population of mule deer. *J. Mammal.* 53: 384-387.
4. BANDY, P. J., W. D. KITTS, A. J. WOOD and I. McT. COWAN. 1957. The effect of age and the plane of nutrition on the blood chemistry of the Columbian black-tailed deer, (*Odocoileus hemionus columbianus*) B. blood glucose, non-protein nitrogen, total plasma protein, plasma albumin, globulin, and fibrinogen. *Can. J. Zool.* 35: 283-289.
5. BENJAMIN, M. M. 1961. *Outline of Veterinary Clinical Pathology*. 2nd ed. Iowa State Univ. Press, Ames. 186 pp.
6. BOLTE, J. R., J. A. HAIR and J. FLETCHER. 1970. White-tailed deer mortality following tissue destruction induced by Lone Star ticks. *J. Wildl. Mgmt.* 34: 546-552.
7. BROWMAN, L. G. and H. S. SEARS. 1955. Erythrocyte values, and alimentary canal pH values in the mule deer. *J. Mammal.* 36: 474-476.
8. COOK, R. S., M. WHITE, W. C. GLAZENER and D. O. TRAINER. 1971. Mortality of young white-tailed deer fawns in South Texas. *J. Wildl. Mgmt.* 35: 47-56.
9. COWAN, I. McT. and A. J. WOOD. 1955. The normal temperature of the Columbian black-tailed deer. *J. Wildl. Mgmt.* 19: 154-155.
10. COWAN, I. McT. and P. J. BANDY. 1969. Observations on the haematology of several races of black-tailed deer (*Odocoileus hemionus*). *Can. J. Zool.* 47: 1021-1024.
11. DOMMERT, A. R., M. E. TUMBLESON, R. B. WESTCOTT, D. A. MURPHY and L. J. KORSCHGEN. 1968. Hematologic values for dieltrin-treated white-tailed deer (*Odocoileus virginianus*) in Missouri. *Amer. J. Vet. Clin. Path.* 2: 181-184.
12. JOHNSON, H. E., W. G. YOUATT, L. D. FAY, H. D. HARTE and D. E. ULLREY. 1968. Hematological values of Michigan white-tailed deer. *J. Mammal.* 49: 749-754.
13. KITCHEN, H. and W. R. PRITCHARD. 1962. Physiology of blood. Pp. 109-114. *In Proc. First Nat'l. White-tailed Deer Disease Symposium*. Univ. Georgia, Athens. 202 pp.
14. KITTS, W. D., P. J. BANDY, A. J. WOOD and I. McT. COWAN. 1956. Effect of age and plane of nutrition on the blood chemistry of the Columbian black-tailed deer (*Odocoileus hemionus columbianus*) A. packed-cell volume, sedimentation rate, and hemoglobin. *Can. J. Zool.* 34: 477-484.

15. LEWIS, L. L. and L. M. MELNICK. 1960. Determination of calcium and magnesium with (ethyleal dinitrolo) tetracetic acid. *Analytical Chem.* 32: 38-42.
16. ROSEN, M. N. and A. I. BISCHOFF. 1952. The relation of hematology to condition in California deer. *N. Amer. Wildl. Conf. Trans.* 17: 482-496.
17. SEAL, U. S. and A. W. ERICKSON. 1969. Hematology, blood chemistry and protein polymorphisms in the white-tailed deer (*Odocoileus virginianus*). *Comp. Biochem. and Physiol.* 30: 695-713.
18. SEAL, U. S., J. J. OZOGA, A. W. ERICKSON and L. J. VERME. 1972. Effects of immobilization on blood analysis of white-tailed deer. *J. Wildl. Mgmt.* 36: 1034-1040.
19. SEAL, U. S., L. J. VERME, J. J. OZOGA and A. W. ERICKSON. 1972. Nutritional effects on thyroid activity and blood of white-tailed deer. *J. Wildl. Mgmt.* 36: 1041-1052.
20. TABER, R. D., K. L. WHITE and N. S. SMITH. 1959. The annual cycle of condition in the Rattlesnake, Montana, mule deer. *Montana Acad. Science* 19: 72-79.
21. TEERI, A. E., W. VIRCHOW, N. F. COLOVOS and F. GREELEY. 1958. Blood composition of white-tailed deer. *J. Mammal.* 39: 269-274.
22. TUMBLESON, M. E., M. G. WOOD, A. R. DOMMERT, D. A. MURPHY and L. J. KORSCHGEN. 1968. Biochemic studies on serum from white-tailed deer in Missouri. *Amer. J. Vet. Clin. Path.* 2: 121-125.
23. TUMBLESON, M. E., J. D. CUNEIO and D. A. MURPHY. 1970. Serum biochemical and hematological parameters of captive white-tailed fawns. *Can. J. Comp. Med.* 34: 66-71.
24. ULLREY, D. E., W. G. YOUATT, H. E. JOHNSON, P. K. KU and L. D. FAY. 1964. Digestibility of cedar and aspen browse for the white-tailed deer. *J. Wildl. Mgmt.* 28: 791-797.
25. ULLREY, D. E., W. G. YOUATT, H. E. JOHNSON, L. D. FAY and B. E. BRENT. 1967. Digestibility of cedar and jack pine browse for the white-tailed deer. *J. Wildl. Mgmt.* 31: 448-454.
26. ULLREY, D. E., W. G. YOUATT, H. E. JOHNSON, L. D. FAY, B. E. BRENT and K. E. KEMP. 1968. Digestibility of cedar and balsam fir browse for the white-tailed deer. *J. Wildl. Mgmt.* 32: 162-171.
27. WHITE, M. 1973. The whitetail deer of the Aransas National Wildlife Refuge. *Texas J. Science* 24: 457-489.
28. WHITE, M., F. F. KNOWLTON and W. C. GLAZENER. 1972. Effects of dam-newborn fawn behavior on capture and mortality. *J. Wildl. Mgmt.* 36: 897-906.
29. WILBUR, C. G. and P. F. ROBINSON. 1958. Aspects of blood chemistry in the white-tailed deer. *J. Mammal.* 39: 309-310.
30. YOUATT, W. G., L. J. VERME and D. E. ULLREY. 1965. Composition of milk and blood in nursing white-tailed does and blood composition of their fawns. *J. Wildl. Mgmt.* 29: 79-84.

Received for publication 24 May 1973
