

SOME HELMINTH PARASITES OF THE AMERICAN BALD EAGLE

Author: KOCAN, A. ALAN

Source: Journal of Wildlife Diseases, 10(1): 8-10

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-10.1.8

The BioOne Digital Library (https://bioone.org/) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (https://bioone.org/subscribe), the BioOne Complete Archive (https://bioone.org/archive), and the BioOne eBooks program offerings ESA eBook Collection (https://bioone.org/esa-ebooks) and CSIRO Publishing BioSelect Collection (https://bioone.org/esa-ebooks) and CSIRO Publishing BioSelect Collection (https://bioone.org/csiro-ebooks).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commmercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SOME HELMINTH PARASITES OF THE AMERICAN BALD EAGLE

A. ALAN KOCAN, Department of Parasitology and Laboratory Practice, School of Public Health, University of North Carolina, Chapel Hill, North Carolina 27514, U.S.A.

LOUIS N. LOCKE, Bureau of Sports Fisheries and Wildlife, Patuxent Wildlife Research Center, Laurel, Maryland 20810, U.S.A.

Abstract: Bald eagles (Haliaeetus leucocephalus) found dead or moribund in the United States and Canada and submitted to Patuxent Wildlife Research Center were examined for helminth parasites. Nine genera of helminths were reported which include new host records for Clinostomum complanatum, Neogogatea pandionis, Centrorhynchus sp., Serratospiculum amaculata, Capillaria contorta, and Habronema americanum.

INTRODUCTION

The American bald eagle (Haliaeetus leucocephalus) has shown a dramatic decline in numbers over the last two decades throughout the continental United States. A great deal of attention has been focussed on the importance of pesticide poisoning and other factors which have contributed to mortality and reduced fecundity. 5,5,2,11 Few reports are available on the helminth parasites of this North American bird, perhaps because of its general decline in numbers and intensified protection in recent years.

MATERIALS AND METHODS

Fifty-nine eagles, found dead or moribund in the United States and Canada and submitted to Patuxent Wildlife Research Center were examined for helminth parasites. These eagles were collected between 1963 and 1971. Because of the condition of the birds upon receipt and the variable care afforded the birds prior to arrival, no attempts could be made to determine prevalence or degree of infection.

RESULTS

Nine genera of helminths (four trematodes, one cestode, one acanthocephalan, and four nematodes) were recorded from the eagles examined (Table 1). Clinostomum complanatum, Neogogatea pandionis, Centrorhynchus sp., Capillaria contorta, Habronema americanum, and Serratospiculum amaculata are reported here for the first time from this host. Although this is the first report of these genera from bald eagles, many of them have been previously reported from European and Asian eagles and a number of other raptors. 10,6,5,7

DISCUSSION

The exact role of helminths in raptor populations is not clear. It appears that birds subjected to greater than normal stresses, hand reared birds, and captive birds are more susceptible to parasitic infections and other environmental and biological insults than are birds from wild populations. 18,4,8 Although no evidence of pathogenicity or mortality could be directly attributed to the helminths recovered in this study, pathogenicity has been previously reported for Serratospiculum amaculata by Bigland et al., and esophageal capillarids by Coopers and Trainer et al., is in other raptors.

The Present address: Dept. of Veterinary Parasitology and Public Health, Oklahoma State University, Stillwater, Oklahoma 74074.

TABLE 1. Helminths found in North American bald eagles.

Parasite	Collection location	Age	Sex	Necropsy diagnosi	s Date
Trematoda					
Clinostomum complanatum	Minnesota	fledgling	F	enteritis emaciation	1972
Phagicola longus	North Carolina	2 years	F	gunshot	1965
Neodiplostomum banghami	Arkansas Iowa	adult immatur e	M M	none gunshot	1963 1966
Neogogatea pandionis	Iowa Massachusetts Wisconsin	immature immature immature	M F M	gunshot	1966 1966 1966
Acanthocephala					
Centrorhynchus sp.	Maine Florida Iowa New Jersey	adult immature immature adult	F F F	none gunshot gunshot none	1967 1971 1966 1963
Cestodes					
Cladotaenia banghami	Minnesota Idaho Florida Minnesota Wisconsin	immature immature immature immature immature	M M F F M	gunshot impact injuries	1968 1968 1971 1968 1968
Nematodes					
Contracaecum sp.	New Jersey Minnesota South Dakota Illinois Minnesota Wisconsin South Dakota	adult immature immature immature adult adult	F M F M F		1963 1969 1968 1971 1969 1968 Irin)
Habronema americanum	Minnesota Iowa Illinois Wisconsin Illinois Minnesota	immature adult immature immature immature immature	F F M F F	impact injuries none gunshot	1967 1971 1970 1969 1967
Serratospiculum amaculata	Wisconsin Illinois	immature immature	M F	gunshot gunshot	1969 1971
Capillaria contorta	Iowa Missouri Minnesota Wisconsin	immature immature immature adult	F F M	strangled gunshot gunshot none	1967 1969 1966 1965

Acknowledgements

The authors wish to thank Mr. Stanley N. Wiemeyer, Wildlife Biologist, Patuxent Wildlife Research Center, for his critical review of the manuscript and for information on the histories of the eagles.

LITERATURE CITED

- BIGLAND, C. H., SI-KWANG, LIU and M. L. PERRY. 1964. Five cases of Serratospiculum amaculata (Nematoda: Filarioidea) infection in prairie falcons (Falco mexicanus). Avian Dis. 8: 412-419.
- COON, N. C., L. N. LOCKE, E. CROMARTIE and W. L. REICHEL. 1970. Causes of bald eagle mortality, 1960-1965. J. Wildl. Dis. 6: 72-76.
- COOPER, J. E. 1969. Oesophageal capillariasis in captive falcons. Vet. Rec. 84: 634-636.
- 4. COOPER, J. E. 1969. Some diseases of birds of prey. Vet. Rec. 84: 454-457.
- 5. COOPER, J. E. 1972. Hawks and parasites. Hawk Chalk 11: 31-35.
- 6. KEYMER, I. F. 1972. Diseases of birds of prey. Vet. Rec. 90: 579-594.
- MAROTEL, G. 1899. Sur un type particulier d'Acanthocephale (Echinorhynchus tenicandatus n. sp.) C.R. Soc. Biol. 6: 226-228.
- 8. MULHERN, B. M., W. L. REICHEL, L. N. LOCKE, T G. LAMONT, A. BESILE, E. CROMARTIE, G. E. BAGLEY and R. M. PROUTY. 1970. Organochlorine residues and autopsy data from bald eagles 1966-1968. Pesticides Monitoring J. 4: 141-144.
- REICHEL, W. L.,T. G. LAMONT, E. CROMARTIE and L. N. LOCKE. 1969. Residues in two eagles suspected of pesticide poisoning. Bull. Environ. Contamination and Toxicol. 4: 24-30.
- SHEN, S. and S. WU. 1964. A preliminary survey of trematode and nematode parasites in aquatic birds from Inner Mongolia, China (Eng. Sum.) Tung Heueh Pao. 16: 398-415.
- STICKEL, L. F., N. J. CHURA, P. A. STEWART, C. M. MENZIE, R. M. PROUTY and W. L. REICHEL. 1966. Bald eagle pesticide relations. Trans. 31st N. Amer. Wildl. Nat. Res. Conf. 190-200.
- 12. TRAINER, D. O., S. D. FOLZ and W. M. SAMUEL, 1968. Capilariasis in the gyrfalcon. Condor 70: 276-277.
- 13. WOODFORD, M. H. 1966. A Manual of Falconry. Adam and Charles Black, London.

Received for publication May 3, 1973