

SOME HEMATOLOGICAL VALUES OF FREE-RANGING AFRICAN ELEPHANTS 1

Authors: DEBBIE, J. G., and CLAUSEN, B.

Source: Journal of Wildlife Diseases, 11(1) : 79-82

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-11.1.79>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SOME HEMATOLOGICAL VALUES OF FREE-RANGING AFRICAN ELEPHANTS¹

J. G. DEBBIE² and B. CLAUSEN, Veterinary Research Laboratory, P.O. Kabete, Kenya

Abstract: Blood samples were collected from 23 free-ranging African elephants (*Loxodonta africana*) in Tanzania. Red and white blood cell counts, white blood cell differential, hemoglobin, blood copper, and blood inorganic phosphate were determined. No correlation of any of these measurements with age and sex was discernible, except that the total white count was significantly higher in females than in males in all age groups. The findings are compared with those previously reported for captive and free-ranging African elephants.

INTRODUCTION

An opportunity arose in March 1968 to obtain blood samples from African elephants cropped in a herd reduction scheme in Mkomasi Game Reserve, Tanzania. The limited published information on blood values for African elephants,^{1,7,8} especially free-ranging elephants, prompted this report.

MATERIALS AND METHODS

Free-flowing blood from a jugular vein puncture was collected in heparin within 10 min after the elephant was shot. Erythrocytes (RBC) and leukocytes (WBC) were counted in an "improved Neubauer" chamber within 6 h of collection. The RBC diluent was 0.9% physiologic saline solution; the WBC diluent, 2% glacial acetic acid with 1% gentian-violet. The differential blood count was determined by counting 100 leukocytes on thin Giemsa-stained blood smears. Hemoglobin levels were determined by the alkaline hematin method,⁹ copper by the modified method² of Eden and Green,³ and inorganic phosphate by the

method of Fiske and Subbarow.⁴ The last three tests were performed in the laboratory within 48 h of collection of the blood in vials containing an anticoagulant consisting of 6 mg of sodium fluoride and 1 mg of "Merthiolate" per ml of blood.⁵ The copper and inorganic phosphate determinations were selected because of the sandy, heavily weathered soil in the area inhabited by the elephants.

The age of the elephants was determined by dental criteria established by Laws.⁶

RESULTS

Table 1 gives the results obtained. It is important to remember that the normal sedimentation rate (PCV) of elephant blood is very high (29 mm/h)¹⁰ in comparison with that of other living animals. For this reason samples were obtained only from animals that could be bled within 10 min of death.

The average RBC counts for males ($3.78 \pm 0.48 \times 10^6$) and females ($3.75 \pm 0.17 \times 10^6$) did not differ appreciably

¹ A project of the Government of Kenya, supported by the United Nations Development Program with the Food and Agriculture Organization of the United Nations as the executing agency.

² Present address: Division of Laboratories and Research, New York State Department of Health, Albany, New York 12201, U.S.A.

TABLE 1. Some Hematological Values of Blood from Free-Ranging African Elephants (*Loxodonta africana*).

Elephant No. (Sex, age in yrs.)	Red blood cells (10 ⁶ /mm ³)	White blood cells (mm ³)	Lymphocytes (%)	Nonsegmented neutrophils (%)	Segmented neutrophils (%)	Monocytes (%)	Eosinophils (%)	Basophils (%)	Hemoglobin (g/100ml)	Copper (ppm)	Inorganic phosphate (mg/100ml)
120 (M, 2-1/4)	3.76	12,000	79	2	11	8	0	0			
124 (M, 3-1/2)	2.20	8,600	56	5	22	16	1	0			
212 (M, 24)	1.92	8,200	85	2	7	5	1	0			
214 (M, 7)	6.74	8,200	72	0	21	6	1	0	5.2	12.8	1.13
218 (M, 1/2)	3.21	7,150	69	2	11	17	1	0	9.1	11.8	1.05
223 (M, 5)	3.11	5,950	69	2	18	7	4	0			
228 (M, 14)	4.32	13,450	63	1	27	12	1	1			
229 (M, 8)	4.10	9,700	76	0	18	6	0	0			
231 (M, 7-1/2)	4.66	5,700	81	1	10	6	2	0			
108 (F, 48)	4.39	7,850	58	0	31	10	1	0			
110 (F, 9)	3.58	21,200	66	1	24	6	3	0			
112 (F, 4)	3.76	12,000									
118 (F, 23)	4.28	7,950									
121 (F, 24)	3.12	13,000									
123 (F, 2-1/2)	4.01	15,300									
126 (F, 7-1/2)	3.32	15,500	65	0	33	2	0	0			
216 (F, 30)	3.11	16,000	70	1	17	11	1	0	6.3	10.4	0.89
217 (F, 1)	4.22	10,300	60	1	31	5	2	1			
206 (F, 8)			68	0	21	10	1	0	8.0	9.4	1.11
211 (F, 26)									7.5	7.1	0.86
215 (F, 23)			62	1	32	3	2	0	7.3	11.1	
221 (F, 3-1/2)			76	1	12	9	2	0	7.3	11.4	1.19
226 ((F, 17)			68	1	22	8	1	0	5.9	10.1	1.08
Sample											
size	23	18	18	18	18	18	18	18	8	8	7
Mean	12.6	3.77	11,000	69.1	1.17	20.17	8.17	1.33	0.11	7.08	10.51
		±	±	±	±	±	±	±	±	±	±
		0.25*	975	1.89	0.03	0.61	0.03	0.25	0.08	0.44	0.06
Range		1.92-	5,700-	56-	0-	7-	2-	0-	5.2-	7.1-	0.86-
		6.74	21,200	85	5	33	17	4	9.1	12.8	1.19

* Standard deviation.

($P < 0.05$). However, there was a significant difference ($P > 0.05$) in WBC count by sex, with the average for males being $8,700 \pm 860/\text{mm}^3$ and for females, $13,230 \pm 1142/\text{mm}^3$. The remaining values in the table are self-explanatory, with no discernible ($P < 0.05$) correlation between age and sex.

DISCUSSION

This study was conducted under less than ideal conditions. The red and white blood counts were performed up to 6 h after collection, which may bias the determined values. Moreover, all the elephants in this study were found infected with the bile duct hookworm *Grammocephalus* sp., and the presence of these and other parasites could have a bearing on the blood values obtained. However, the data may be useful until true norms can be determined.

Young and Lombard¹⁰ found the RBC count for 11 male elephants averaged 5.02×10^6 , which is much higher than our results. Since their samples were from immobilized elephants, the speed of sedimentation of the RBC might explain the discrepancy, although one might expect the opposite effect, namely a decrease in erythrocyte count and PCV, as described in reports which compare the effects of drugs and physical restraint in white-tailed deer.^{7,8} Bartels *et al.*¹ quote 3.20×10^6 as the RBC count for one

African elephant in captivity while Schmitt⁹ cites 4.64×10^6 for four elephants, of which only one was African.

WBC determinations for elephant blood have been reported to range from 10,000 to 14,400/ mm^3 and hemoglobin from 12 to 14 g/100 ml. Our results agree for WBC but are lower for hemoglobin. However, we made only eight hemoglobin determinations.

Previously published results on the differential blood count of elephants are minimal. Our finding of 69% lymphocytes is high compared to Young and Lombard's 58%, but more striking is our finding of 8% monocytes and 1% nonsegmented neutrophils compared with 1.5% and 13%, respectively, reported by Young and Lombard.¹⁰ Each of two elephants had 1% basophils (three cells actually seen) indicating that elephants do have basophils, a fact not previously reported. The low percentage of eosinophils is surprising considering the large number of tissue eosinophils observed to be associated with the bile duct lesions of *Grammocephalus* sp.

Determinations of copper and inorganic phosphate in elephant blood appear not to have been previously published. Although our sample size is small, there is close correlation between samples. The average for blood inorganic phosphate (7.08 ± 0.44 mg/100 ml) is higher than the 4.5 mg/100 ml seen in mature cattle blood. The average for copper corresponds with results in cattle.⁵

Acknowledgements

The authors express their gratitude to Wildlife Services Ltd., Nairobi, for permission to collect specimens and providing accommodation; to the Tanzania Veterinary and Game Departments for permission to collect specimens in Tanzania; and to the Director of Veterinary Services, Kenya, for permission to publish this paper.

LITERATURE CITED

1. BARTELS, H., P. HILPERT, K. BARBEY, K. BETKE, K. RIEGEL, E. M. LANG and J. METCALFE. 1963. Respiratory functions of blood of the yak, llama, camel, Dybowski deer and African elephant. *Am. J. Physiol.* 205: 331-336.
2. CLARE, N. T., I. J. CUNNINGHAM and D. D. PERRIN. 1945. The determination of copper in pastures and livers. *New Zealand J. Sci. Tech.* 26: 340-350.

3. EDEN, A. and H. H. GREEN. 1940. Micro-determination of copper in biological material. *Biochem. J.* 34: 1202-1208.
4. FISKE, C. H. and Y. SUBBAROW. 1925. The colorimetric determination of phosphorus. *J. Biol. Chem.* 66: 375-400.
5. LAMPKIN, K. and D. A. HOWARD. 1962. Studies on the production of beef from Zebu cattle in East Africa. IV. Variation in blood hemoglobin and its response to the feeding of dicalcium phosphate. *J. Agric. Sci.* 59: 245-249.
6. LAWS, R. M. 1966. Age criteria for the African elephant, *Loxodonta a. africana*. *E. Afr. Wildl. J.* 4: 1-37.
7. PRESIDENTE, P. J. A., J. H. LUMSDEN, K. R. PRESNELL, W. A. RAPLEY and B. M. McCRAW. 1973. Combination of etorphine and xylazine in captive white-tailed deer. II. Effects on hematologic, serum biochemical and blood gas values. *J. Wildl. Dis.* 9: 342-348.
8. SEAL, U. S., J. J. OZAGA, A. W. ERICKSON and L. J. VERME. 1972. Effects of immobilization on blood analyses of white-tailed deer. *J. Wildl. Mgmt* 36: 1034-1040.
9. SCHMITT, J. 1964. Hematological studies in elephants. *Vet. Med. Res.* 2: 87-95.
10. YOUNG, E. and C. J. LOMBARD. 1967. Physiological values of the African elephant (*Loxodonta africana*). *The Veterinarian* 4: 169-172.

Received for publication 15 August 1974