

THE OCCURRENCE OF GAPEWORM (Syngamus trachea) IN WILLOW GROUSE

Authors: WISSLER, KATHRIN, and HALVORSEN, ODD

Source: Journal of Wildlife Diseases, 11(2): 245-247

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-11.2.245

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

THE OCCURRENCE OF GAPEWORM (Syngamus trachea) IN WILLOW GROUSE

KATHRIN WISSLER and ODD HALVORSEN, Institute of Biology and Geology University of Tromsø, 9001 Tromsø, Norway

Abstract: 21.4% of the willow grouse (*Lagopus lagopus*) examined from a coastal locality in northern Norway were infected with *Syngamus trachea*. None of the willow grouse examined from an inland locality with colder and longer winters were infected with the parasite. It is suggested that the parasite may influence survival of the grouse population.

INTRODUCTION

The nematode S. trachea parasitizes the trachea of a large number of bird species.⁶ The infection causes the socalled gapeworm disease well known in poultry.

Gapeworms have been recorded from wild birds in Norway including capercaillie (*Tetrao urogallus*),⁴ black grouse (*Lyrurus tetrix*)⁵, magpie (*Pica pica*), house sparrow (*Passer domesticus*), arctic tern (*Sterna paradisaea*), and fieldfare (*Turdus pilaris*).¹ In the cases reported by Bakke¹ only one to three infected birds of each species were recorded.

Bakke noted that all known cases of gapeworm infection in birds in Norway were from the southern and western part of the country, and he suggested that the distribution of S. trachea is restricted to these areas for climatic reasons.¹

We have found *S. trachea* in a population of willow grouse in northern Norway.

MATERIAL AND METHODS

Willow grouse were collected from one island locality (Karls ϕ y, approx. 9° E, 70°N) and one inland locality (Sennaland, approx. 10°E, 70° 10'N).

The main climatic difference between the coast and the interior is a colder and longer winter in the interior. In 1972 Solovomi near Sennaland had monthly average temperatures below 0C from January to April and from October to December. Monthly averages below 0C in Troms ϕ near Karls ϕ y were recorded from January to March and in November. The lowest monthly average temperature at Solovomi was —15,9C and in Troms ϕ —2,4C, both in February.

On Karls ϕ y samples of grouse were collected monthly from July to December 1973 and from Sennaland the sampling period included September, October and November of the same year. Most of the adult grouse were obtained by shooting, while chicks were collected by a variety of methods. The birds were deep frozen as soon as possible after being killed and were examined for helminths later.

RESULTS

From the inland locality (Sennaland) 48 willow grouse were collected in September, 12 in October and 5 in November. None of these birds were infected with gapeworms.

Of the 70 willow grouse examined from the island locality (Karls ϕ y) 15 (21.4%) were infected with from 1 to 23 pairs of *S. trachea.* Only chicks and young birds were infected, and only birds collected in July, August and September (Table 1).

MonthexaminedInfecteJuly910July910August10August1820September1210October1330		Older than	ı 28 days	28 days or	. younger	Total	Intensity of infection
July 9 1 0 August 10 September 18 2 0 October 12 1 0 November 13 3 0	Examined Infected	Examined	Infected	Examined	Infected	- prevalence of infection	rairs of S. trachea
August 10 <th< td=""><td>1 0</td><td>4</td><td>ব</td><td>4</td><td>7</td><td>66.7</td><td>6.1 (1-23)</td></th<>	1 0	4	ব	4	7	66.7	6.1 (1-23)
September 18 2 0 October 12 1 0 November 13 3 0		10	٢	I	1	70.0	3.4 (1-7)
October 12 1 0 November 13 3 0	2 0	16	7	I	I	1.11	1.0 (1)
November 13 3 0	1 0	11	0	1	1	0	
•	3 0	10	0	I	I	0	
December 8 3 0	3 0	S	0	I	I	0	

Journal of Wildlife Diseases Vol. 11, April, 1975

TABLE 1. The prevalence and intensity of infection with Syngamus trachea in willow grouse from the island Karlsøy.

DISCUSSION

Our observations show that *S. trachea* is endemic in a coastal locality in northern Norway and thus not restricted to southern parts of the country for climatic factors. The absence of the parasite in grouse samples from an inland locality in the north may however be caused by the more severe winters in that locality.

The life cycle of *S. trachea* may include an invertebrate transport host in which the parasite may live for a considerable length of time.⁷ According to Barth² insects are an important part of the diet of willow grouse during the first 2 weeks after hatching. Most of the

willow grouse eggs on Karlsøy hatched in 1973 during the last week of June and the first week of July (Parker pers. comm.) Barus³ showed that S. trachea may live from 48 to 108 days with a mean life of 72 days in the final host. Based on these data one would expect a decrease in the gapeworm infection in grouse on Karlsøy around mid-September. The decrease and disappearance of the infection in the material examined appear more abrupt than one would expect from the life data of the parasite alone, and indicate that S. trachea may influence survival in this population of willow grouse.

LITERATURE CITED

- BAKKE, T. A. 1973. Studies of the helminth fauna of Norway XXVII: Syngamiasis in Norway. Norw. J. Zool. 21: 299-303.
- 2. BARTH, E. K. 1949. Hva spiser lirypekyllingene? Jeger & Fisker 6: 134-140.
- BARUS, V. 1964. The morphological and biometrical variability of the nematode Syngamus (Syngamus) trachea (Montagu 1811) Chapin 1925 and a revision of the species composition of the subgenus Syngamus. Vest. Csl. Spol. Zool. 28: 290-304.
- 4. LUND, H. M.-K. 1946. Endoparasites in the capercaillie (Tetrao urogallus). Skand. Vet. Tidsskr. 36: 641-662.
- 5. LUND, H. M.-K. 1954. Nematodes, cestodes and coccidia found in black grouse (Lyurus tetrix) ia Norway. Statens Viltundersøkelser, 1-42.
- 6. MADSEN, H. 1950. On the systematics of Syngamus trachea (Montagu, 1811) Chapin 1925. J. Helminth. 24: 33-46.
- 7. SOULSBY, E. J. L. 1968. Helminths, Arthropods and Protozoa of Domestic Animals. Bailliere, Tindall and Cassell. London.

Received for publication 17 September 1974