

EXPERIMENTALLY TRANSMITTED MARBLE SPLEEN DISEASE IN PEN-RAISED WILD TURKEYS *

Authors: ILTIS, J. P., JAKOWSKI, R. M., and WYAND, D. S.

Source: Journal of Wildlife Diseases, 11(4) : 484-485

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-11.4.484>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

EXPERIMENTALLY TRANSMITTED MARBLE SPLEEN DISEASE IN PEN-RAISED WILD TURKEYS*

J. P. ILTIS, R. M. JAKOWSKI and D. S. WYAND, Department of Pathobiology,
University of Connecticut, Storrs, Connecticut 06268, USA

Abstract: Pen-raised North American wild turkeys (*Meleagris gallopavo* L.) were experimentally infected with marble spleen disease (MSD) to determine their susceptibility to this disease. Gross and microscopic lesions were consistent with experimental MSD in pheasants and domestic turkeys: an enlarged mottled spleen, intranuclear inclusion bodies, and absence of pulmonary edema and hemorrhage. Detectable levels of viral antigen were not demonstrable in sera of turkeys using the agar gel precipitin test.

INTRODUCTION

Marble spleen disease (MSD) of ring-necked pheasants (*Phasianus colchicus* L.) was experimentally transmitted to pen-raised ring-necked pheasants and domestic turkeys (*Meleagris gallopavo* L.) by Iltis *et al.*¹ This study¹ proved that a filterable agent, probably a virus, was the causative agent of MSD.^{1,2,4,5} by demonstrating infectivity with 0.20 μ m filtered inoculum. However, cases of MSD transmitted in the laboratory⁴ lacked the terminal pulmonary involvement occurring in field cases.^{1,2,5}

Pen-raised wild turkeys bred for state repopulation programs represent an analogous situation to pen-raised ring-necked pheasants. Whereas field outbreaks of MSD occur in pen-raised pheasants none have been reported in wild turkeys raised in captivity. Domestic turkeys are susceptible to experimental MSD infection but domestic and wild turkeys must have some genetic differences; therefore, the present study was undertaken to determine the wild turkey's susceptibility.

MATERIALS AND METHODS

Experimental Birds

Twelve turkeys, approximately 8 weeks old, from the Pennsylvania Game Commission's Wild Turkey Farm in Williamsport, Pennsylvania were supplied by Harvey A. Roberts¹ with the assistance of Eugene P. Nelson.² Of the nine birds used for this study, five were housed in an outdoor pen off the ground and four were housed indoors in two separate units off the floor.

Inoculations

Spleens (kept at -60°C for 3 months) from two domestic turkeys experimentally infected with MSD and killed on postinoculation day 5 (representing the fourth passage in turkeys of the pheasant derived infectious agent in the laboratory) were prepared by adding 10 ml of sterile phosphate buffered saline (PBS), pH 7.2, then homogenizing in a Ten Broeck grinder followed by two freeze-thaw cycles in a dry ice-alcohol bath. One ml of this inoculum was administered orally via canula to the five turkeys

* Scientific Contribution No. 606, Storrs Agricultural Experiment Station, University of Connecticut, Storrs, Connecticut 06268. This study was supported by a grant from the Northeastern Research Center for Wildlife Diseases, College of Agriculture and Natural Resources, University of Connecticut.

¹ Deputy Executive Director, Pennsylvania State Game Commission.

² Superintendent of Wild Turkey Farm, Williamsport, Pennsylvania.

housed outdoors and also to two turkeys housed indoors. The other two turkeys kept in the other indoor unit served as uninoculated controls. The turkeys were 12 weeks old at inoculation.

Gross and Microscopic Examinations

On postinoculation day 5 all nine turkeys were killed and viscera examined for gross lesions. Sections of spleen, liver and lung were fixed in 10% formalin, embedded in paraffin, sectioned at 5 μ m and stained with hematoxylin and eosin.

Agar Gel Precipitin Tests

Prior to inoculation, sera from the nine turkeys were tested for agar gel precipitin (AGP) antibodies to marble spleen antigen as previously described.⁸ In addition, serum samples from each bird were taken on post-inoculation days 1 to 5 and tested for AGP antigen reactive to antibody in convalescent serum from a pheasant exposed to MSD.

RESULTS AND DISCUSSION

The only gross lesions observed were enlargement of the spleen and slight splenic mottling in four of the seven experimentally infected turkeys. The two uninoculated controls had no gross or microscopic lesions. No clinical signs were observed in any of the birds.

Microscopic lesions observed in the

seven inoculated turkeys consisted primarily of intranuclear inclusion bodies (INIB) typical of MSD.^{1,2,5} The INIB occurred in cells of spleen, liver, and lung. In addition, splenic lymphoid hyperplasia also was noted. The gross and microscopic findings observed in experimentally transmitted MSD in these seven pen-raised wild turkeys were consistent with those previously reported for the pen-raised pheasant and domestic turkey.⁸

No AGP antibody was detected in sera prior to experimental infection. Also, no antigen, indicative of viremia, was detected by the AGP test in sera of nine turkeys on postinoculation days 1 to 5.

Presumably, outdoor housing of turkeys and pheasants experimentally infected with MSD might better simulate field conditions, which could possibly induce typical field cases of MSD,^{1,6} rather than the experimental MSD (typified by absence of terminal lung involvement and presence of INIB on postinoculation days 3, 4, and 5).⁸ Results of this study demonstrate that while pen-raised wild turkeys are susceptible to experimentally transmitted MSD, no change in the course of the disease between birds housed indoors and outdoors had occurred by postinoculation day 5. The possibility of field outbreaks of MSD in pen-raised wild turkeys remains unknown; however, serum samples taken from these birds in the fall and tested for AGP antibody to MSD antigen should indicate if inapparent infections occur.

LITERATURE CITED

1. CARLSON, H. C., J. R. PETTIT, R. V. HEMSLEY and W. R. MITCHELL. 1973. Marble spleen disease of pheasants in Ontario. *Can. J. Comp. Med.* 37: 281-286.
2. ILTIS, J. P. and D. S. WYAND. 1974. Indications of a viral etiology for marble spleen disease in pheasants. *J. Wildl. Dis.* 10: 272-278.
3. ———, R. M. JAKOWSKI and D. S. WYAND. 1975. Transmission of marble spleen disease in turkeys and pheasants. *Am. J. vet. Res.* 36: 97-101.
4. JAKOWSKI, R. M. and D. S. WYAND. 1972. Marble spleen disease in ring-necked pheasants: Demonstration of agar gel precipitin antibody in pheasants from an infected flock. *J. Wildl. Dis.* 8: 261-263.
5. WYAND, D. S., R. M. JAKOWSKI and C. N. BURKE. 1972. Marble spleen disease in ring-necked pheasants: Histology and ultrastructure. *Avian Dis.* 16: 319-329.

Received for publication 3 March 1975