

Dermatophilus DERMATITIS ENZOOTIC IN DEER IN NEW YORK STATE AND VICINITY

Authors: GORDON, M. A., SALKIN, I. F., and Stone, W. B.

Source: Journal of Wildlife Diseases, 13(2) : 184-190

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.2.184>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Dermatophilus DERMATITIS ENZOOTIC IN DEER IN NEW YORK STATE AND VICINITY

M. A. GORDON and I. F. SALKIN, Division of Laboratories and Research,
New York State Department of Health, Albany, New York 12201, USA

W. B. STONE, Wildlife Research Center, New York State Department of Environmental
Conservation, Delmar, New York 12054, USA

Abstract: In a recent 2-year period *Dermatophilus congolensis* infection, ranging from mild to very severe, was found in 15 white-tailed deer (*Odocoileus virginianus*) taken in southeastern New York State (12) adjacent New Jersey (2) and central Vermont (1). The host range for *Dermatophilus congolensis* is given.

INTRODUCTION

Dermatophilus congolensis infection was first noted in 1915 in African domestic cattle³⁶ and is usually manifested as an exudative epidermitis (dermatophilosis, streptotrichosis, mycotic dermatitis, lumpy wool). The disease commonly affects cattle, sheep, goats and domesticated equines in many parts of the world, often creating severe economic problems.^{1,35} Infection has been reported in a number of additional mammalian hosts, including the polar bear (*Thalarctos maritimus*);^{24,33} domestic cat (*Felis domesticus*);^{3,36} chamois (*Rupicapra rupicapra*);²⁵ domestic dog (*Canis familiaris*);⁵ eland (*Taurotragus oryx*);¹⁴ fox (*Vulpes vulpes*);² giraffe (*Giraffa camelopardalis*) and Thomson's gazelle (*Gazella thomsonii*);²¹ gerbil (*Gerbillus* sp.); hare (*Lepus* sp.) and hedgehog (*Erinaceus europaeus*);¹⁹ owl monkey (*Aotus trivirgatus*);^{7,17,22} woolly monkey (*Lagothrix lagotrichia*);⁸ pig (*Sus scrofa*);³⁴ cottontail rabbit (*Sylvilagus floridanus*);⁴ raccoon (*Procyon lotor*);³⁰ South American seal (*Otaria bryonia*);⁹ Columbian ground squirrel (*Citellus columbianus*);³⁹ and zebra (*Equus* sp.).¹³ The only non-mammalian host thus far encountered is the Australian bearded lizard (*Amphibolurus barbatus*);^{23,32} Ten human cases also have been reported.^{6,11,15,18,20}

The first report of *Dermatophilus* infection in a white-tailed deer (*Odocoileus virginianus*) was by Dean *et al.*⁶

in New York State. In 1970, the death of a white-tailed deer fawn in South Carolina was attributed to dermatophilosis,¹⁸ and in 1972 two more cases in this host were reported from New York.^{36,37}

The present report was occasioned by the finding, in a recent 2-year period, of 15 additional cases of *Dermatophilus* dermatitis in wild deer, 12 within the State of New York, 2 in adjacent New Jersey, and 1 in nearby Vermont. Nine occurred between June and December, 1974 and 5 between September, 1975 and January, 1976.

MATERIALS AND METHODS

Specimens brought to the microbiology laboratory for examination included ears, or portions thereof, excised portions of skin bearing crusts or scabs, eroded or denuded areas of skin, pustules and detached scabs. Crusts, scabs and pustules were rinsed several times in sterile distilled water, and the underside of each was rubbed on an area approximately 2 cm² of a beef heart infusion—5.5% horse blood agar plate. This was streaked for bacterial isolation, and the plates were incubated at 35 C. The same specimens also were rubbed on microscope slides and stained by the Giemsa method or with methylene blue. Denuded

skin was decontaminated with 70% ethanol and scraped deeply with a sterile scalpel, and smears and cultures were prepared from this material as described above. Portions of all specimens also were placed in 10% formalin and were subsequently sectioned and stained with hematoxylin and eosin, Giemsa stain and methenamine-silver.

In some instances, cultures of *D. congolensis* were obtained by rabbit passage. An area approximately 4 cm² on the outer surface of the ear of an albino laboratory rabbit (*Oryctolagus cuniculus*) was disinfected with 70% ethanol, lightly scarified with a scalpel, and then rubbed with affected tissue from the deer.

RESULTS

Data on the 15 affected deer are presented in Table 1. Reports of the first three of these cases have been publish-

ed.^{24,25} Geographic distribution is indicated in Figure 1. The gross appearance of a particularly severely diseased animal (case 10) is shown in Figure 2.




FIGURE 1. Distribution of cases of dermatophilosis in deer in New York State and adjacent Vermont and New Jersey.



FIGURE 2. Severe *Dermatophilus* dermatitis in adult male deer (case no. 10), with desquamation of skin of the snout, antlers and ears. The entire body was covered with eruptions.

TABLE 1. *Dermatophilus* infection in white-tailed deer, 1974-76.

Case no.	Date	County ^a	Age, Sex	Description	Confirmed by ^b		
					Direct examination	Direct culture	Rabbit passage
1	7/74	Morris, NJ	2 wk, M	Encrusted papules on muzzle and ears.	+	+	ND
2	7/74	Morris, NJ	2 wk, F	Encrusted papules on muzzle and ears.	+	ND	ND
3	10/74	Greene	5.5 yr, F	Lactating; emaciated; scattered encrusted lesions on legs, flanks and back. Cultures from leg. Also had alternariosis.	+	+	+
4	7/74	Columbia	5 wk, M	Complete body involvement; died shortly after arrival at the laboratory.	+	+	ND
5	10/74	Roxbury, VT	Fawn, M	Extensive hair loss everywhere except legs and tail, especially head and neck. A few encrusted lesions. Cultures from base of ears.	+	+	ND
6	10/74	Columbia	1.5 yr, M	Paintbrush effect; acute lesions along back with spreading.	+	—	ND
7	10/74	Columbia	? F	"Brushfire" effect.	+	—	+
8	12/74	Columbia	2.5 yr, M	"Brushfire" effect. Alopecia over sacral, lumbar and thoracic vertebrae, and on both sides.	+	—	+

TABLE 1 (Continued)

Case no.	Date	County ^a	Age, Sex	Description	Confirmed by ^b		
					Direct examination	Direct culture	Rabbit passage
9	12/74	Columbia	6.5 yr, F	"Brushfire" effect. Alopecia over sacral, lumbar and thoracic vertebrae, and on both sides.	+	—	+
10	9/75	Sullivan	1.5 yr, M	Entire body covered with eruptions; skin rotted on ear, snout and antlers. Sections full of <i>D. congolensis</i> .	+	+	ND
11	11/75	Columbia	1.5 yr, M	"Brushfire" effect. Alopecia over sacral, lumbar and thoracic vertebrae and on both sides.	+	+	ND
12	12/75	Greene	7.5 yr, F	Denuded along back and much of both sides of thorax and abdomen.	+	—	ND
13	12/75	Dutchess	2.5 yr, F	Area of alopecia over lumbar region and several more on thorax, with whitish pustules.	+	—	+
14	1/76	Dutchess	15 yr, F	Extensive infection of skin over sacral and lumbar vertebrae, extending down the flanks.	+	—	ND
15	8/76	Washington	12 wk, F	Found moribund. Pustules and scabs about the nose, lips, eyes, ears, legs, haired regions of the toes, abdomen, and back.	+	+	+

^aIn New York State except where noted.^b+ = positive, — = negative, ND = not done.

Bacteriology

In the cases in which *D. congolensis* was cultured directly from lesions, typical colonies¹⁰ appeared within 48 h. They were 0.5-1 mm in diameter, opaque white to yellowish, glabrous, rough, raised, round to square or irregular in shape, tough and adherent. They pitted the agar and left mycelia in the medium when the colonies were removed.

Methylene blue- or Giemsa-stained smears from lesions (Fig. 3), as well as tissue sections, showed the diagnostic branching filaments of *D. congolensis*, which divide both transversely and longitudinally and become converted into packets of cocci.

Inoculation of rabbit ears with infected tissues resulted in the appearance of an erythematous rash within 72 h, followed by multiple small pustules which developed scabs. The infection was self-limiting. *D. congolensis* was observed in




FIGURE 3. Smear of exudate from lesion of case no. 10, illustrating septate branching filaments of *D. congolensis*. Older hyphae are dividing in three planes to form cocci. Note long, slender, nonseptate, young branches. Giemsa stain, X 1200.

Giemsa-stained smears of pustules or of the underside of scabs and was cultured from these materials.

DISCUSSION

Dermatophilosis (streptotrichosis) has been defined³⁵ as "an acute or chronic infection of the epidermis resulting from an exudative dermatitis with scab formation, caused by *Dermatophilus congolensis*." In domestic cats, granulomata of the tongue and of the urinary bladder also have occurred.

The clinical spectrum of epidermitis ranges from a paintbrush effect (tufts of hair fused by exudate) through nodular and pustular lesions to severe exudative epidermitis with scabbing and crusting, often accompanied by secondary bacterial infection. A "brushfire" effect (generalized hair loss over the saddle and rump, with solitary tufts of hair at the periphery) also can be seen during the rutting season.

Obtaining pure cultures is often difficult because of bacterial contamination but usually can be accomplished by informed selection of specimens and careful bacterial streaking on blood agar plates. Alternatively, tissues from a diseased deer are examined microscopically, and samples in which *D. congolensis* is observed are inoculated into a rabbit's ear. The inocula raise typical lesions on the ear, and scrapings from these lesions are cultured. In our hands *D. congolensis* always has been recovered by this procedure.¹²

Transmission of infection is by direct contact and possibly by means of vectors.²⁷ Deer must, therefore, be considered a potential source of infection for domestic cattle. However, although dermatophilosis in cattle has been reported from many parts of the United States, thus far it has not been recognized in New York. Infection of horses has been noted in New York State and Vermont.⁴ The first four known cases of spontaneous dermatophilosis of humans, contracted from a single infected deer, were reported from New York State⁸ and the fifth from Toronto, Canada.¹¹

Dermatophilus infection of deer has not, except for the single case in South Carolina,¹⁸ been reported anywhere except in New York State and its immediate vicinity, where there are now 18 known instances. The sighting of many

additional deer with the "brushfire" effect has been reported to us by wildlife technicians. The disease undoubtedly occurs elsewhere and probably is misdiagnosed clinically—but as what?

LITERATURE CITED

1. AINSWORTH, G. C. and P. K. C. AUSTWICK. 1973. *Fungal Diseases of Animals* p. 216, 2nd Ed., Commonwealth Agricultural Bureaux, England.
2. AUSTWICK, P. K. C. 1968. Mycotic infections. Symp. Zool. Soc. Lond., 24: 249-271.
3. BAKER, G. J., R. G. BREEZE and C. O. DAWSON. 1972. Oral dermatophilosis in a cat: A case report. *J. Small Anim. Pract.* 13: 649-653.
4. BENTINCK-SMITH, J., F. H. FOX and D. W. BAKER. 1961. Equine dermatitis (cutaneous streptothricosis) infection with dermatophilus in the United States. *Cornell Vet.* 51: 343-349.
5. BLANCOU, J. 1973. Infection du chien par *Dermatophilus congolensis* (Van Saceghem, 1915). *Rev. Elev. Méd. Vét. Pays Trop.* 26: 289-291.
6. DEAN, D. J., M. A. GORDON, C. W. SEVERINGHAUS, E. T. KROLL and J. R. REILLY. 1961. Streptothricosis: A new zoonotic disease. *N.Y. State J. Med.* 61: 1283-1287.
7. FOX, J. G., L. H. CAMPBELL, C. REED, S. B. SNYDER and O. A. SOAVE. 1973. Dermatophilosis (cutaneous streptothricosis) in owl monkeys. *J. Am. vet. med. Ass.* 163: 642-644.
8. FRASER, C. E. O. and F. G. GARCIA. 1971. Cutaneous streptothricosis (dermatophilosis) in a woolly monkey. *Primate Zoonosis Surveillance*, Report No. 5, January-March 1971. U.S. Department of Health, Education and Welfare, Center for Disease Control, Atlanta, Georgia.
9. FRESE, K. and A. WEBER. 1971. Eine dermatitis bei mähenrobber (*Otaria bryonia* Blainville) hervorgerufen durch *Dermatophilus congolensis*. *Berlin und München. Tierarzt. Wochenschrift* 84: 50-54.
10. GORDON, M. A. 1964. The genus *Dermatophilus*. *J. Bacteriol.* 88: 509-522.
11. ———. 1976. Characterization of *Dermatophilus congolensis*: Its affinities with the Actinomycetales and differentiation from *Geodermatophilus*. In: *Dermatophilus Infection in Animals and Man* p. 187-201, Ed. by D. H. Lloyd and K. C. Sellers, Academic Press, London.
12. ——— and U. PERRIN. 1971. Pathogenicity of *Dermatophilus* and *Geodermatophilus*. *Infect. Immun.* 4: 29-33.
13. GREEN, H. F. 1960. Streptothricosis in zebra and donkeys and demodectic mange in eland in Kenya. *Vet. Rec.* 72: 1098.
14. HORNBY, H. E. 1920. A contagious impetigo of cattle. *Vet. J.* 76: 210-216.
15. Infections in cattle pose new occupational perils for DVMs. 1964. *Vet. Dispatch* 6 (7): 1-2.
16. KAMINSKY, G. W. 1971. *Dermatophilus congolensis*—human infection in Australia p. 76-77, C. Rend. Commun. V. Cong. SIMHA, Paris.
17. KING, N. W., C. E. O. FRASER, F. G. GARCIA, L. A. WOLF and M. E. WILLIAMSON. 1971. Cutaneous streptothricosis (dermatophiliasis) in owl monkeys. *Lab. An. Sci.* 21: 67-74.
18. KISTNER, T. P., E. B. SHOTTS and E. W. GREENE. 1970. Naturally occurring streptothricosis in a white-tailed deer. *J. Am. vet. med. Ass.* 157: 633-635.

19. KUSELTAN, I. V. 1968. Nocardiosis of lambs in Tadzhik S.S.R. Vet. Bull. 38: 590 (Abstract).
20. LONDERO, A. T., C. D. RAMOS and L. P. SOUZA. 1974. Human dermatophilosis, its occurrence in Brazil. Mykosen 17: 111-113.
21. MacKENZIE, P. A. and R. M. SIMPSON. 1956. *The African Veterinary Handbook* 2nd Ed., Pittman, Nairobi.
22. McCLURE, H. M., W. KAPLAN, W. B. BONNER and M. E. KEELING. 1971. Dermatophilosis in owl monkeys. Sabouraudia 9: 185-190.
23. MONTALI, R. J., E. E. SMITH, M. DAVENPORT and M. BUSH. 1975. Dermatophilosis in Australian bearded lizards. J. Am. vet. med. Ass. 167: 553-555.
24. NEWMAN, M. S., R. W. COOK, W. K. APPELHOF and H. KITCHEN. 1975. Dermatophilosis in two polar bears. J. Am. vet. med. Ass. 167: 561-564.
25. NICOLET, J., J. KLINGER and H. FEY. 1967. *Dermatophilus congolensis* agent de la streptotrichose du chamois. Path. Microbiol. 30: 831-837.
26. O'HARA, P. J. and D. O. CORDES. 1963. Granulomata caused by *Dermatophilus* in two cats. New Zealand Vet. J. 11: 151-154.
27. RICHARD, J. L. and A. C. PIER. 1966. Transmission of *Dermatophilus congolensis* by *Stomoxys calcitrans* and *Musca domestica*. Am. J. vet. Res. 27: 419-423.
28. ROSCOE, D. E., R. C. LUND, M. A. GORDON and I. F. SALKIN. 1975. Spontaneous dermatophilosis in twin white-tailed deer fawns. J. Wildl. Dis. 11: 398-401.
29. SALKIN, I. F., M. A. GORDON and W. B. STONE. 1975. Dual infection of a white-tailed deer by *Dermatophilus congolensis* and *Alternaria alternata*. J. Am. vet. med. Ass. 167: 571-573.
30. ———. 1975. Dermatophilosis: Increased prevalence in New York State with involvement of wild raccoons. p. 93, 75th Ann. Mtg., American Society for Microbiology (Abstract).
31. SHOTTS, E. B. and T. P. KISTNER. 1970. Naturally occurring cutaneous streptothricosis in a cottontail rabbit. J. Am. vet. med. Ass. 157: 667-670.
32. SIMMONS, G. C., N. D. SULLIVAN and P. E. GREEN. 1972. Dermatophilosis in a lizard (*Amphibolurus barbatus*). Austral. Vet. J. 48: 465-466.
33. SMITH, C. F. and D. O. CORDES. 1972. Dermatitis caused by *Dermatophilus congolensis* in polar bears (*Thalarctos maritimus*). Br. Vet. J. 128: 366-371.
34. STANKUSHEV, Kh., I. SIMOV, I. DENEV, N. KYURTOV and M. ALEKSANDROV. 1968. Mycotic dermatitis in pigs. Vet. Sbir. Sofia 65: 3-5.
35. STEWART, G. H. 1972. Dermatophilosis: A skin disease of animals and man. Part I. Vet. Rec. 91: 537-544.
36. STONE, W. B. 1972. Dermatophilosis (cutaneous streptothricosis) in a white-tailed deer fawn. N.Y. Fish Game J. 19: 92-95.
37. ———. 1972. Another case of dermatophilosis in a white-tailed deer. N.Y. Fish Game J. 19: 184.
38. VAN SACEGHEM, R. 1915. Dermatoze contagieuse (impétigo contagieux). Bull. Soc. Path. Exot. 8: 354-359.
39. WOBESER, G. and M. A. GORDON. 1969. *Dermatophilus* infection in Columbian ground squirrels. Bull. Wildl. Dis. Ass. 5: 31-32.

Received for publication 2 September 1976