

Mycobacterium intracellulare INFECTION IN A WATER MONITOR

Authors: FRIEND, S.C.E., and RUSSELL, E.G.

Source: Journal of Wildlife Diseases, 15(2) : 229-233

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-15.2.229>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Mycobacterium intracellulare INFECTION IN A WATER MONITOR

S.C.E. FRIEND[□] and E.G. RUSSELL, Department of Agriculture, Veterinary Research Institute, Parkville, Victoria 3052, Australia

Abstract: Mycobacteriosis caused by *Mycobacterium intracellulare* serotypes Davis (8) and Altman (18) is described in a water monitor (*Varanus semiremex*). Infection with this organism has not been reported previously in reptiles in Australia.

INTRODUCTION

Tuberculosis has been reported in many poikilothermic animal species, viz. snakes, turtles, lizards and crocodiles.^{14,16,24} Validly described species of mycobacteria associated with naturally-occurring infections in lower vertebrates include *M. chelonae*,⁵ *M. fortuitum*,¹¹ *M. thamnophaeos*,¹ *M. marinum*,⁷ *M. xenopi*²¹ and *M. avium*.^{21,22} Mycobacteriosis in turtles, snakes, lizards and crocodiles has been characterized by a granulomatous inflammation with involvement of skin, kidney, liver, spleen and lung.^{2,14,16,25} This report describes a miliary granulomatous mycobacterial infection caused by *M. intracellulare* serotypes Davis (8) and Altman (18) in a water monitor (*Varanus semiremex*).

CASE HISTORY

A young male water monitor, a large Australian lizard, was acquired from Queensland in June, 1976, and housed with several other species of lizard in a large enclosure. It became ill shortly thereafter, refused food and began to show abnormal postural positions. It was presented live to the Veterinary Research Institute for examination and was observed to circle to the left and rest in a tightly wound position. The lizard was killed and examined at necropsy.

NECROPSY FINDINGS

The lizard was thin and the bones were prominent. Hard yellowish-white nodules, 2 to 10 mm in diameter, with dry caseous centres were present throughout the mesentery, kidneys, liver, lung, under the fascia of the thigh muscles, retroperitoneally along the spine and beneath the periosteum of the sacrum (Fig. 1).

MICROSCOPIC RESULTS

Tissues for histologic examination were fixed in 10% buffered formalin, sectioned at 6 µm and stained with haematoxylin and eosin and by the Ziehl-Neelsen (ZN) and Auramine Fluorescent¹³ method for acid-fast bacteria.

Microscopically, the nodules consisted of intensely eosinophilic, amorphous, caseous necrotic debris enclosed by a mantle of epithelioid cells and heterophils (Fig. 2). Fibrin and granulation tissue often were associated with these areas, but there was no evidence of giant cell formation. Acid-fast organisms were not found in tissues or on impression smears with routine or modified ZN staining methods, but were present in sections stained by the fluorescent method.

[□] Present address: Department of Veterinary Pathology, Western College of Veterinary Medicine, University of Saskatchewan, Saskatoon, Saskatchewan, Canada S7N 0W0.




FIGURE 1. Multiple white nodules visible in lung, liver, kidney and mesentery of a water monitor with *M. intracellulare* infection.




FIGURE 2. Caseous necrotic nodule within the kidney enclosed by a mantle of epithelioid cells and heterophils (ZN $\times 160$).

MICROBIOLOGIC RESULTS

Portions of liver and samples of faeces were homogenized in phosphate-buffered saline (PBS) and then shaken in an equal volume of 4% sodium hydroxide for 15 min. The resultant suspension was neutralized with 10% hydrochloric acid and centrifuged at $250 \times g$ for 5 min to remove debris. The supernatant was then centrifuged at $1600 \times g$ for 30 min. The pellet was resuspended in a small quantity of PBS and streaked onto slopes of Lowenstein-Jensen, Stonebrink and modified Herold with mycobactin media.²³ Slopes were incubated at 22 C and 37 C for at least 12 weeks.

After 6 weeks, a pure growth of pale yellow pigmented colonies was observed on Lowenstein-Jensen slopes inoculated from the liver specimen and incubated at 22 C and 37 C. Later, growth occurred on other slopes. The isolate was acid-fast, negative in the Tween 80 hydrolysis, nitrate reduction and semiquantitative catalase tests²⁶ and positive in the tellurite reduction test.¹⁰ Inoculation of 1.4×10^5 organisms into the wing vein of a 6-week-old chicken produced lesions characteristic of avian tuberculosis.

A suspension was tested by Schaefer's seroagglutination test,¹⁹ as modified by Reznikov and Leggo¹⁵ and found to consist of a mixture of *Mycobacterium intracellulare* serotypes Davis (8) and Altman (18).

M. intracellulare serotype Davis was recovered from tissues of experimentally infected chickens.

DISCUSSION

Reptiles debilitated by injury, malnutrition or other disease are susceptible to mycobacterial disease.¹⁴ Isolation of mycobacteria alone does not necessarily indicate disease, as they are commonly found in aquaria, water in contact with animals, and water from lakes and streams.⁶ Furthermore, frogs experimentally infected with *M. intracellulare* serotype Davis shed bacteria into the environment without showing signs of disease.⁸ However, the granulomata found in the viscera, mesentery and bone of this monitor are consistent with those lesions described in reptiles with tuberculosis.^{2,3,7,12,14,16} Involvement of fascia and musculature of the thigh, and the adjacent tissues and periosteum of the spine and sacrum may have resulted in the circling behavior shown by the lizard.

The distribution of *M. intracellulare* is widespread and it has been isolated from soil,²⁷ house-dusts⁴ and water.^{6,18} In Australia, infection by *M. intracellulare* (Battey disease) is a serious and important medical problem.¹⁷ Both serotypes isolated have been implicated in human pulmonary and lymph node infections.²⁰ *M. intracellulare* serotype Davis is frequently found in water⁹ and it seems very likely that the environment was the source of the causative organisms of this infection. Lizards may, therefore, act as a possible reservoir and a health hazard to humans, wildlife and domestic animals.

Acknowledgements

We thank David J. Dawson of the Tuberculosis Section of the Laboratory of Microbiology and Pathology, Queensland Department of Health, Brisbane, for serotyping the organisms, Ms. J.F. Bell for excellent technical assistance and Ian Shirley and Gunther Appl for the photography.

LITERATURE CITED

1. ARONSON, J.D. 1929. Spontaneous tuberculosis in snakes. *J. infect. Dis.* 44: 215-223.

2. BROCK, J.A., R.M. NAKAMURA, A.Y. MIYAHARA and E.M.L. CHANG. 1976. Tuberculosis in Pacific green sea turtles, *Chelonia mydas*. Trans. Am. Fish. Soc. 105: 564-566.
3. COWAN, D.F. 1968. Diseases of captive reptiles. J. Am. vet. med. Ass. 153: 848-859.
4. DAWSON, D.J. 1971. Potential pathogens among strains of mycobacteria isolated from house-dust. Med. J. Aust. 1: 679-681.
5. FRIEDMAN, F.F. 1903. Der Schildrotentuberkelbazillus, seine Zuchtung, Biologie und Pathogenität. Zentbl. Bakt. Parasitkde., Abt. I. 34: 647.
6. GOSLEE, S. and E. WOLINSKY. 1976. Water as a source of potentially pathogenic mycobacteria. Am. Rev. resp. Dis. 113: 287-292.
7. GRIFFITH, A.S. 1941. The susceptibility of the Water or Grass Snake (*Trepidonotus natrix*) to the avian tubercle bacillus and to reptilian strains of acid-fast bacilli. J. Hyg., Camb. 41: 284-288.
8. KAZDA, J. and R. HOYTE. 1972. Zur Ökologie von *Mycobacterium intracellulare* serotype Davis. Zentbl. Bakt. Hyg., I. Abt. Orig. A. 222: 506-509.
9. ———. 1973. Die Bedeutung von Wasser für die Verbreitung von potentiell pathogenen Mycobakterien. Zentbl. Bakt. Hyg., I. Abt. Orig. B. 158: 161-169.
10. KUBICA, G.P. 1973. Differential identification of mycobacteria. VII. Key features for identification of clinically significant mycobacteria. Am. Rev. resp. Dis. 107: 9-21.
11. KUSTER, E. 1905. Über Kaltblutertuberkulose. Münch. med. Wschr. 52: 57-59.
12. MARCUS, L.C. 1971. Infectious disease of reptiles. J. Am. vet. med. Ass. 159: 1626-1631.
13. MOTE, R.F., R.L. MUHM and D.C. GIGSTAD. 1975. A staining method using acridine orange and auramine O for fungi and mycobacteria in bovine tissue. Stain Technol. 50: 5-9.
14. REICHENBACH-KLINKE, H. and E. ELKA. 1965. *The Principal Diseases of Lower Vertebrates*, p. 385-543. Academic Press, London and New York.
15. REZNIKOV, M. and J.H. LEGGO. 1972. Modification of Schaefer's procedure for serotyping of organisms of the *Mycobacterium avium* - *M. intracellulare* - *M. scrofulaceum* complex. Appl. Microbiol. 23: 819-823.
16. RHODIN, A.G.J. and M.R. ANVER. 1977. Mycobacteriosis in turtles: Cutaneous and hepatosplenic involvement in a *Phrynosoma hilari*. J. Wildl. Dis. 13: 180-183.
17. RUNYON, E.H. 1971. Whence Mycobacteria and Mycobacterioses. Ann. intern. Med. 75: 467-468.
18. SAITO, H. and M. TSUKAMURA. 1976. *Mycobacterium intracellulare* from public bath water. Jap. J. Microbiol. 20: 561.
19. SCHAEFER, W.B. 1965. Serologic identification and classification of atypical mycobacteria by their agglutination. Am. Rev. resp. Dis. 92: 85-93.
20. ———. 1968. Incidence of the serotypes of *Mycobacterium avium* and atypical mycobacteria in human and animal diseases. Am. Rev. resp. Dis. 97: 18-23.
21. SCHWABACHER, H. 1959. A strain of mycobacterium isolated from skin lesions of the cold-blooded animal, *Xenopus laevis*, and its relation to atypical acid-fast bacilli occurring in man. J. Hyg., Camb. 57: 57-67.
22. THOEN, C.O., W.D. RICHARDS and J.L. JARNAGIN. 1977. Mycobacteria isolated from exotic animals. J. Am. vet. med. Ass. 170: 987-990.

23. USDA, APHIS, Veterinary Services Laboratories. 1974. *Laboratory Methods in Veterinary Mycobacteriology*, Ames, Iowa.
24. VOGEL, H. 1958. Mycobacteria from cold-blooded animals. *Am. Rev. Tuberc. pulm. Dis.* 77: 823-838.
25. WALLACH, J.D. 1969. Medical care of reptiles. *J. Am. vet. med. Ass.* 155: 1017-1034.
26. WAYNE, L.G. and J.R. DOUBEK. 1968. Diagnostic key to mycobacteria encountered in clinical laboratories. *Appl. Microbiol.* 16: 925-931.
27. WOLINSKY, E. and T.K. RYNEARSON. 1968. Mycobacteria in soil and their relation to disease associated strains. *Am. Rev. resp. Dis.* 97: 1032-1037.
28. ——— and W.A. SCHAEFER. 1973. Proposed numbering scheme for mycobacterial serotypes by agglutination. *Int. J. Syst. Bacteriol.* 23: 182-183.

Received for publication 7 June 1978
