



TICK PARALYSIS IN A GREY FOX

Author: JESSUP, DAVID A.

Source: Journal of Wildlife Diseases, 15(2) : 271-272

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-15.2.271>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TICK PARALYSIS IN A GREY FOX

DAVID A. JESSUP, Wildlife Investigations Laboratory, California Department of Fish and Game, 987 Jedemith Drive, Sacramento, California 95819, USA.

Abstract: One hundred seventeen *Dermacentor variabilis* were removed from the head and back of a grey fox (*Urocyon cinereoargenteus*) showing paresis and diminished motor reflexes of the hind limbs. Rapid and total recovery led to a diagnosis of tick paralysis.

INTRODUCTION

A number of species of ticks are capable of producing progressive ascending paralysis in man, cattle, dog and bison (*Bison bison*) as well as other animals.^{5,8} *Dermacentor andersoni* produces a neuroparalytic substance in the saliva which interferes with the synthesis or release of acetylcholine at motor end plates.^{3,4,5,7,8} *Dermacentor occidentalis* previously has been reported as a cause of tick paralysis in cattle and blacktailed deer (*Odocoileus hemionus columbianus*) in California.^{1,6} The purpose of this report is to present the signs of tick paralysis in the grey fox and verify the tick involved as *Dermacentor variabilis*.

CASE HISTORY

An adult male grey fox (*Urocyon cinereoargenteus*) found along a country road was unable to use its hind limbs and was brought into the Wildlife Disease Section on 17 April 1978. The animal weighed 4 kg, was alert, responsive and only mildly aggressive. In attempting to escape it would drag its hindquarters but offered no resistance when grasped by the nape of the neck. Ocular or nasal discharges were not present. Examination of the oral cavity and pharynx revealed pale, pink, dry mucous membranes, a rapid capillary refill time but no lesions. Pupils were of equal size, responsive to light and the anterior

chamber and the retinal fundus appeared normal. The rectal temperature was 39 C, the heart rate 80/min., and the respiratory rate 20/min., all considered normal. Stethoscope auscultation of both lung fields was unremarkable. A loss of tone to the anal sphincter and sluggish response to stimulus was noted. The hind limbs were semi-flaccid and flexor or toe pinch and patellar reflexes were absent. No pain or crepitation could be elicited on palpation of the hind limbs, pelvis, sacral, lumbar, or thoracic spine; moreover, no fractures, dislocations or evidence of soft tissue or skin trauma could be found. A strong femoral pulse was present in both hind limbs. The urinary bladder was extremely distended.

A number of ticks were visible about the head, neck and shoulders. Further exploration revealed many additional ticks at various locations, resulting in a tentative diagnosis of tick paralysis. Ticks were carefully removed manually and with forceps. The animal was given subcutaneous fluids, Benzathine and Procaine Penicillin intramuscularly, canine distemper vaccinated, and sprayed with a carbaryl and pyrethrin containing insecticide spray. The bladder was manually expressed and the animal was placed in a cage and given food and water.

Within 24 h, partial use of the hind limbs had returned and at 48 h, use of the

hind limbs and anal sphincter were normal. Patellar and flexor reflexes were present. Appetite was good and aggression manifested as fear toward humans, was marked. The animal was returned to the locale where it was found and released.

Examination of the ticks removed revealed 15 fully engorged females, 26 partially or unengorged females and 76 males. They were identified as *Dermacentor variabilis* on the basis of scutal pattern, spiracular plates and host and geographic location.²

Acknowledgements

Oscar A. Brunetti, Wildlife Pathologist, California Department of Fish and Game identified the ticks and assisted in literature review.

LITERATURE CITED

1. BRUNETTI, O. 1965. Tick paralysis in California deer. *Calif. Fish and Game*. 51: 208-210.
2. COOLEY, R.A. 1938. The genera *Dermacentor* and *Otocentor* (Ixodidae) in the United States, with studies in variation. *Nat. Inst. of Hlth. Bull. No. 171*: 21-29.
3. ESPLIN, D.W., C.B. PHILLIP and L.E. HUGHES. 1960. Impairment of muscle stretch reflexes in tick paralysis. *Science*. 132: 958-959.
4. ETTINGER, S.J. 1975. *Textbook of Veterinary Internal Medicine*. Vol. 1., pp. 413-414. W.B. Saunders, Philadelphia. 862 pp.
5. JUBB, K.V.F. and P.C. KENNEDY. 1970. *Pathology of Domestic Animals*. Vol. 2., p. 632 Academic Press, New York. 697 pp.
6. LOOMIS, E.C. and R.B. BUSHNELL. 1968. Tick paralysis in California livestock. *Am. J. Vet. Res.* 29: 1089-1093.
7. MURNAGHEN, M.F. 1960. Conduction block of terminal somatic motor fibres in tick paralysis. *Canad. J. Biochem. & Physiol.* 38: 287-295.
8. SMITH, H.A., T.C. JONES and R.D. HUNT. 1972. *Veterinary Pathology*, p. 813. Lea and Febiger, Philadelphia, Pennsylvania. 1521 pp.

Received for publication 14 September 1978