

HELMINTH PARASITES OF FISHER *Martes pennanti* (ERXLEBEN) FROM MANITOBA, CANADA

Authors: DICK, T. A., and LEONARD, R. D.

Source: Journal of Wildlife Diseases, 15(3) : 409-412

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-15.3.409>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

HELMINTH PARASITES OF FISHER *Martes pennanti* (ERXLEBEN) FROM MANITOBA, CANADA

T. A. DICK and R. D. LEONARD, Department of Zoology, University of Manitoba, Winnipeg, Manitoba, R3T 2N2, Canada.

Abstract: Seven species of helminths were recovered during a survey of 162 fisher (*Martes pennanti*) from four areas of Manitoba: *Baylisascaris devosi* in 52 fisher; *Taenia sibirica* in 25; *Physaloptera* sp. in nine; *Alaria mustelae* in two; *Metorchis conjunctus* in one; *Trichinella spiralis* in one of 81; *Molineus* sp. in one. *B. devosi* was the most prevalent parasite and differences in its geographical distribution were possibly related to population density of fisher. The prevalence of other parasites did not appear to be related to density of fisher.

INTRODUCTION

Few studies have been done on parasites of fisher and none on a fisher (*Martes pennanti*) population in the central to northern parts of its range. A few studies have incorporated parasite data in broader ecological and management studies.^{3,5,6} A list of parasites from fisher collected in Maine,⁷ and a new species of nematode, *Baylisascaris devosi*, from fisher and marten (*Martes americana*) have been described.¹¹ This study was undertaken to determine the parasites of fisher in Manitoba, their distribution in the province and parasite prevalence and intensity based on host age.

MATERIALS AND METHODS

One hundred and fifty-seven fisher were collected from four study areas: 105 from area A (50° 40'N; 95° 30'W); nine from area B (50° 0'N; 95° 10'W); 17 from area C (49° 20'N; 95° 30'W); 26 from area D (52° 30'N; 101° 30'W); one from the Berens River (52° 20'N; 97° 0'W), one from the Pigeon River (52° 10'N; 97° 0'W) and three from the Thompson area (55° 45'N; 97° 50'W). Carcasses, frozen from one to ten months, were obtained from trappers and the Manitoba Wildlife Branch. Five specimens were fresh. Carcasses were thawed at room temperature and the entire gastrointestinal tract,

lungs and heart were removed for examination.

The gastrointestinal tract was divided into three major sections: esophagus-stomach, small intestine (separated into four sections), and large intestine. The mucosa was scraped off each section, stirred in a tray and the suspended material examined and then decanted. Parasites were removed, fixed in FAA and stored in 70% ethanol. Nematodes were cleared in lactophenol and *en face* preparations made. Cestodes and trematodes were stained with acetocarmine.

Adult fisher were distinguished from juveniles by the presence of open sutures in the skulls of fisher under one year old.¹⁵

RESULTS

Results are summarized in Table 1. Helminths were recovered from 52 of 162 animals examined. Only *T. sibirica* recovered from 19 females and 6 males showed a significant difference ($p < 0.01$) in prevalence between sexes. Juveniles had higher prevalences of *B. devosi* and *Physaloptera* sp.: 36.0% had *B. devosi* compared to 18.9% of adults; 6.4% had *Physaloptera* sp. compared to 2.7% of adults. There was not a significant

TABLE 1. Prevalence and intensity of gastrointestinal parasites of 162 fisher from Manitoba, Canada (1975-77).

Parasite	NMCIC(P) ¹	Prevalence of Infection			Intensity ⁴
		Total Sample	Juveniles	Adults	
<i>Baylisascaris devosi</i> Sprent (1952)	1979-234	32.0	36.0	18.9	2.17±1.68
<i>Taenia sibirica</i> Dubnitzky (1952)	1979-235	15.4	16.0	13.5	1.84±1.31
<i>Physaloptera</i> sp.	1979-236	5.6	6.4	2.7	1.44±1.01
<i>Alaria mustelae</i> Bosma (1934)	1979-238	1.2	1.6	2.7	55.00±18.03
<i>Metorchis conjunctus</i> ² (Cobbold, 1860) Looss, 1899	1979-237	0.6	—	—	25
<i>Trichinella spiralis</i> ³ (Owen, 1835) Raillet, 1895		1.2	—	—	1
<i>Molineus</i> sp. ²		0.6	—	—	1

¹Accession number to National Museum of Canada Invertebrate Collection (Parasites).

²one specimen infected

³one infection in 81 examined

⁴mean ± SD

difference between prevalences of *T. sibirica* in juvenile and adult fisher.

Prevalence of parasites in fisher varied between sample areas: 44.8% of the fisher population from area A was infected with *B. devosi* and 18.1% infected with *T. sibirica*. Prevalence values for *B. devosi* in areas B, C, and D were 11.1, 5.9, and 11.7, respectively. Prevalence values for *T. sibirica* in areas B, C, and D were 0, 11.7, and 15.4, respectively. Significant differences ($p < 0.01$) in levels of parasitism were found between areas for *B. devosi* but not for *T. sibirica*.

DISCUSSION

Low numbers of species recorded and low prevalence values are similar to other studies.^{3,5,6} This study, like that of De Vos,⁵ found *B. devosi* in fisher. Recently, *B. devosi* had been found in fisher in New Brunswick (C. Bursey,

pers. comm.) thus indicating a much wider geographical distribution for this nematode than previously known.

Although *P. maxillaris* has been reported from fisher,⁶ recent experimental evidence indicated a high degree of host specificity for this species in skunks.² We have identified specimens from fisher as *Physaloptera* sp.; however, arrangement of anal papillae on the only male specimen recovered does not fit published descriptions for *Physaloptera* spp.⁹

The prevalence and intensity of *Physaloptera* sp. are low suggesting that intermediate or paratenic hosts are not eaten frequently.

T. sibirica in fisher is a new host and North American record. Our identification of *T. sibirica*, based on shape, size and number of hooks has been verified by C. Bursey (pers. comm.) and compared to

indistinguishable specimens obtained from fisher in New Brunswick. The life cycle of *T. sibirica* is not known but since squirrels are a major food item of fisher they are a potential intermediate host.¹⁰ Tapeworms found in other studies^{3,5,6,7} were identified as *Mesocestoides variabilis*, although a *Taenia* sp. was mentioned in one study.³

Alaria sp. has been reported from fisher.^{3,8} Our findings extend the geographical range of *Alaria* in fisher. This is the first report of *M. conjunctus* from fisher and the first report in Manitoba east of Lake Winnipeg. Although fisher are reported to eat fish,^{1,10} no fish were recovered from stomach samples examined in this study.

T. spiralis has been reported from fisher from the Rocky Mountain Region of the United States.¹⁴ Our observation extends the range of *T. spiralis* in wild carnivores into Southern Manitoba. Since the one infected fisher was trapped in cottage country the infection may be due to scavenging around dumps.

The absence of lungworms, kidney worms and guinea worm from fisher in Manitoba is of interest. Lungworms were recorded in fisher collected from Ontario,⁴ and *Dracunculus insignis* from fisher in New York State.¹⁵ *Diectophyma renale* has been reported in fisher from Ontario¹³ and from Maine.³ Furthermore, in area A, where most fisher were trapped, *D. renale* was recovered from a wolf and mink. Examination of fisher stomachs from Manitoba by R. Leonard revealed a high proportion of their diet was mammals and birds. No fish,

amphibians or reptiles were recovered suggesting that these animals were not plentiful or that fisher did not frequent aquatic habitats.

The geographical differences noted in the prevalence of *B. devosi* may be related to current population densities (based on trapping records) and the absence of fisher from areas for an extended time (R. Stardom, pers. comm.). Area A has always had fisher in moderate numbers. In contrast, fisher numbers in area B (Whiteshell Park) have been reduced by trapping and destruction of habitat, and extirpated from area C in the past 30-40 years due to agriculture, logging and trapping. In the last 10 years fisher have returned and are increasing in numbers in areas B and C. Fisher were reduced to very low numbers in area D through intensive trapping and logging operations, but have started to increase in the last 5-10 years. *B. devosi* requires a small rodent to complete its life cycle.¹² Consequently, the absence of fisher from an area, or very low populations of fisher, could reduce the prevalence to a negligible value, especially since wolverines (*Gulo gulo*) have been absent from these areas for 100 years and marten are rarely trapped (R. Stardom, pers. comm.). With increasing numbers of fisher in areas B, C and D we predict an increase in the prevalence of *B. devosi* in these fisher populations. *T. sibirica* did not show obvious geographical differences in levels of infection, thus it appears that its prevalence in fisher populations may be independent of fisher density.

Acknowledgements

We are grateful to the staff of the Manitoba Department of Renewable Resources and Transportation Services for their help in collecting specimens. We particularly thank Mr. R. Stardom, Fur Bearer Specialist, Manitoba Research Section, for his support and Mr. C. Bursey, Department of Biology, University of New Brunswick, Canada, for his invaluable help by loaning specimens and identifying *T. sibirica*. Financial support from the National Research Council of Canada is gratefully acknowledged.

LITERATURE CITED

1. BANFIELD, A.W.F. 1974. *The Mammals of Canada*. University of Toronto Press, Toronto. 438 pp.
2. CAWTHORN, R.J. and R.C. ANDERSON. 1976. Development of *Physaloptera maxillaris* (Nematoda: Physalopteroidea) in skunk (*Mephitis mephitis*) and the role of paratenic and other hosts in its life cycle. *Can. J. Zool.* 54: 313-323.
3. COULTER, M.W. 1966. Ecology and management of fishers in Maine. Ph.D. thesis, State University College of Forestry at Syracuse University.
4. CRAIG, R.E. and R.A. BORECKY. 1976. Metastrongyles (Nematoda: Metastrongyloidea) of fisher (*Martes pennanti*) from Ontario. *Can. J. Zool.* 54: 806-807.
5. DE VOS, A. 1952. The ecology and management of fisher and marten in Ontario. *Tech. Bull. Ont. Dept. of Lands and Forests.* 90 pp.
6. HAMILTON, W.J. JR. and A.H. COOK. 1955. The biology and management of the fisher in New York. *N.Y. Fish and Game J.* 2: 13-35.
7. MEYER, M.C. and B.G. CHITWOOD. 1951. Helminths from fisher (*Martes p. pennanti*) in Maine. *J. Parasit.* 37: 320-321.
8. PEARSON, J.C. 1956. Studies on the life cycles and morphology of the larval stages of *Alaria arisaemoides* Augustine and Uribe, 1927 and *Alaria canis* LaRue and Fallis, 1936 (Trematoda: Diplostomidae). *Can. J. Zool.* 34: 295-387.
9. MORGAN, B.B. 1944. The *Physaloptera* (Nematoda) of carnivores. *Trans. Wis. Acad. Sci. Arts Lett.* 36: 375-388.
10. SETON, E.T.S. 1909. *Life Histories of Northern Animals. An Account of the Mammals of Manitoba*. Vol. II. *Flesh-eaters*. Charles Scribner's Sons, New York. 1267 pp.
11. SPRENT, J.F.A. 1952a. On an *Ascaris* parasite of the fisher and marten, *Ascaris devosi* sp. nov. *Proc. Helm. Soc. Wash.* 19: 27-37.
12. ———. 1952b. On the life history of *Ascaris devosi* and its development in the white mouse and the domestic ferret. *Parasitology*, 42: 244-258.
13. STRICKLAND, M. 1974. Fisher and marten study, Algonquin region (Ontario). Progress report, Ontario Ministry of Natural Resources. 57 pp.
14. WORLEY, D.E., J.C. FOX, J.B. WINTERS and K.R. GREER. 1974. Prevalence and distribution of *Trichinella spiralis* in carnivorous mammals in the United States Northern Rocky Mountain Region. pp. 597-602. In: *Trichinellosis*. Charles W. Kim, ed., Intext Educational Publishers, New York.
15. WRIGHT, P.L. and M.W. COULTER. 1967. Reproduction and growth in Maine fishers. *J. Wildl. Manage.* 31: 70-87.

Received for publication 17 November 1978
