

SEROLOGIC AND HEMATOLOGIC VALUES OF WILD COYOTES IN WISCONSIN

Authors: SMITH, GREGORY J., and Rongstad, Orrin J.

Source: Journal of Wildlife Diseases, 16(4) : 491-497

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-16.4.491>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SEROLOGIC AND HEMATOLOGIC VALUES OF WILD COYOTES IN WISCONSIN

GREGORY J. SMITH and ORRIN J. RONGSTAD, Department of Wildlife Ecology, University of Wisconsin, Madison, Wisconsin 53706, USA.

Abstract: Blood samples were obtained from 30 coyotes (*Canis latrans*) captured in northern Wisconsin in conjunction with radio-telemetry studies. Samples were assayed for seven hematologic values, seven serum chemistries, serum albumin, globulin and total protein. Results are given with respect to sex and age and are compared with available data for captive wild and pen-raised coyotes. Leukocyte counts were greater for males than females and packed cell volumes were greater for adults than young, possibly due to differential response to capture and handling stress. Hemoglobin concentrations and calcium levels suggest differences in nutrition between pen-raised and wild coyotes. Sex and age differences in serum calcium for wild coyotes probably reflect nutritional differences between groups examined. Juvenile coyote serum alkaline phosphatase levels declined curvilinearly with age for coyotes less than one year old, suggesting a possible technique for separating juveniles and yearlings captured in autumn that are released for research purposes. Elevated glucose levels and leukocyte counts in wild coyotes may reflect greater handling stress than for pen-raised and captive coyotes. No significant sex or age effects were found for levels of serum urea nitrogen, total protein, cholesterol, and total bilirubin.

INTRODUCTION

This paper describes serologic and hematologic values of wild, free-ranging coyotes (*Canis latrans*) in northern Wisconsin. These values are examined for normal physiologic variation in terms of age and sex, and are compared with data for wild captive and pen-raised coyotes under controlled conditions.

The coyote has been subject to an increasing amount of research, largely stimulated by the need for predator control. Coyotes are important not only in terms of livestock depredations but also as fur-bearers, game animals and for esthetics.⁵ An increase in coyote field research has created a demand for baseline physiological data.⁷ Seal *et al.*¹⁰ described the potential for integrating field and laboratory studies to provide a more complete understanding of animal populations.

Hematology and serum chemistry analyses have been described for captive

wild coyotes^{2,4} and for pen-raised coyotes.⁷ These studies have provided comparative data for evaluating blood values of wild, free-ranging coyotes. Blood values for animals in captivity reflect constant and adequate levels of nutrition.

Blood values may be influenced by nutrition, sex, age, habitat, season, reproductive status, trauma and other factors due to environmental stress.^{1,7,10} Several studies have evaluated blood parameters of wild ungulates with respect to age, sex, habitat, nutrition, season, immobilization, and capture stress.^{6,9,11,12} There are few data available on the effects of these factors on wild canid blood values. Comparative baseline studies are essential for determining the effects of these factors on blood analysis for wild populations.¹² Blood studies of wild, free-ranging mammals may not only serve as indicators to habitat condition, but may provide

useful techniques for assessment of environmental impacts on animal and habitat quality.

METHODS

Blood samples were obtained from undrugged coyotes, captured using No. 3 leg-hold traps, during all seasons of the year. Blood was removed from the lateral saphenous vein on a hind leg. Blood for hematological studies was placed in a 2 cc vacutainer tube containing a pre-measured amount of ethylenediaminetetraacetic acid disodium salt (EDTA). Air-dried blood smears were prepared in the field, and whole blood was refrigerated until mailed to the laboratory. Blood for serological studies was placed in a 10 cc vacutainer tube, centrifuged, the serum removed and immediately frozen until sent to the laboratory.

Whole blood and serum analysis were done by the Central Animal Health Laboratories, Madison, Wisconsin. Hemoglobin (Hb) concentrations were determined by spectrophotometric methods. Packed cell volumes (PCV) were determined by the microhematocrit method, and white blood cell (WBC) counts were made with a hemocytometer. Differential WBC counts were made from Wright-stained blood smears using standard counting procedures. Mean corpuscular hemoglobin concentration (MCHC) was determined by calculation.⁶ Serum chemistries were analyzed using a Technicon auto-analyzer (SMA 12/60).

Coyotes were aged on the basis of tooth eruption or tooth wear as described by Gier.⁴ Animals greater than one year old are treated here as adults. Juveniles were assigned a standardized age (in weeks) using a 1 May parturition date. This approximate whelping date for our study area, was estimated by monitoring denning activities of radio-tagged adult females.

Sex and age effects on blood parameters were statistically analyzed

using analysis of variance.¹⁴ Comparisons of pooled adult male and female blood values with similar samples from other studies were made using the Student's "t" test for unpaired variates.¹⁴

RESULTS

Blood samples were obtained from 30 coyotes captured in northern Wisconsin from February 1978 to November 1979. Of the 30 coyotes sampled, 16 were captured in autumn, eight in winter, five in summer, and one in spring. One sample obtained during the summer period was analyzed for serum chemistries but not for hematological parameters.

The results of hematological assays with respect to age and sex are compared with data for 18-month-old pen-raised coyotes⁷ (Table 1). Serum chemistry (Table 2) and protein analysis (Table 3) are also compared with data for pen-raised coyotes.⁷ Serum alkaline phosphatase levels for juvenile coyotes decreased curvilinearly with age until about January (Fig. 1).

A summary of significant differences between sex and age groups for hematologic values, serum chemistries, and serum protein values is given in Table 4. All comparisons within our wild, free-ranging coyote sample were made using analysis of variance.

DISCUSSION

Blood samples were obtained from manually restrained, undrugged coyotes. Interpretations of these blood data must take these methods into account. Several studies have related capture and handling effects to elevated blood values.^{2,11,12} Baseline data for captive coyotes reported by Gates and Goering,² Goering *et al.*,⁴ and Rich and Gates⁷ were also collected from restrained, undrugged coyotes. Coyotes we sampled were subject to capture and handling related bias, while captive coyote blood values likely reflect only handling bias.

TABLE 1. Hematologic values of coyotes captured in northern Wisconsin, 1978-79. Captive coyote data from Rich and Gates (1979). Mean values \pm 1 standard deviation.

Cohort	Sample Size	Hematologic value ¹				Leukocyte differential ²			
		Hb g/dl	PCV %	MCHC g/dl	Total WBC $\times 10^3/\mu\text{l}$	N %	L %	M %	E %
Adult ♂	6	14.2 \pm 2.5	47.7 \pm 8.7	29.8	20.3 \pm 8.5	80	12	4	4
Adult ♀	4	15.0 \pm 0.8	49.0 \pm 0.8	30.6	15.5 \pm 5.8	89	5	5	—
Juvenile ♂	12	12.8 \pm 1.5	41.2 \pm 4.0	31.1	24.0 \pm 8.0	84	4	4	7
Juvenile ♀	7	13.2 \pm 3.2	41.7 \pm 9.2	31.6	17.5 \pm 6.2	84	12	3	3
Captive ♂ + ♀ ³	48	17.1 \pm 1.2	50.5 \pm 3.4	33.9	9.0 \pm 1.9	62	26	5	6

¹Hb, hemoglobin; PCV, packed cell volume; WBC, white blood cells; MCHC, mean corpuscular hemoglobin concentration.²N, neutrophils; L, lymphocytes; M, monocytes; E, eosinophils. Mean values only.³Data from Rich and Gates (1979) represents blood obtained from 18-month-old pen-raised coyotes. Coyotes were undrugged and restrained by hand.TABLE 2. Serum chemistries of coyotes in northern Wisconsin, 1978-79. Captive coyote data from Rich and Gates (1979). Mean value \pm 1 standard deviation.

Cohort	Sample Size	Calcium mg/dl	Inorganic phosphorus		Cholesterol mg/dl	Glucose mg/dl	BUN mg/dl	Bilirubin mg/dl
			mg/dl	mg/dl				
Adult ♂	6	8.6 \pm 0.6	3.4 \pm 0.5	153 \pm 35	161 \pm 44	28.2 \pm 13.2	0.1 \pm 0.1	
Adult ♀	5	8.8 \pm 0.7	3.5 \pm 0.7	157 \pm 40	181 \pm 47	21.2 \pm 8.3	0.2 \pm 0.1	
Juvenile ♂	12	9.6 \pm 0.5	6.3 \pm 0.1	196 \pm 54	159 \pm 34	17.2 \pm 9.1	0.1 \pm 0.1	
Juvenile ♀	7	8.9 \pm 0.4	5.1 \pm 1.0	178 \pm 53	158 \pm 44	19.4 \pm 9.2	0.1 \pm 0.1	
Captive ♂ + ♀ ¹	48	9.8 \pm 1.4	3.8 \pm 1.0	—	125 \pm 28	26.6 \pm 7.1	0.2 \pm 0.1	

¹Data from Rich and Gates (1979) represents blood obtained from 18-month-old pen-raised coyotes. Coyotes were undrugged and restrained by hand.

TABLE 3. Serum protein levels of coyotes in northern Wisconsin, 1978-1979. Captive coyote data from Rich and Gates (1979). Mean values \pm 1 standard deviation.

Cohort	Sample Size	Total Protein gm/dl	Albumin ² gm/dl	Globulin gm/dl	Albumin/Globulin ratio
Adult ♂	6	6.4 \pm 0.5	2.9 \pm 0.2	3.5 \pm 0.4	0.9 \pm 0.1
Adult ♀	5	6.4 \pm 0.6	3.1 \pm 0.4	3.3 \pm 0.5	1.0 \pm 0.2
Juvenile ♂	12	6.4 \pm 0.4	3.1 \pm 0.2	3.3 \pm 0.4	0.9 \pm 0.1
Juvenile ♀	7	6.0 \pm 0.3	2.7 \pm 0.4	3.2 \pm 0.4	0.9 \pm 0.2
Captive ♂ + ♀ ¹	48	6.5 \pm 0.3	2.7 \pm 0.3	3.7 \pm 0.3	0.7 \pm 0.1

¹Data from Rich and Gates (1979) represents blood obtained from 18-month-old pen-raised coyotes. Coyotes were undrugged and restrained by hand.

²Technicon SMA 12/60 using BCG dye-binding method.




FIGURE 1. The relationship between serum alkaline phosphatase levels and juvenile coyote age. Coyote ages (in weeks) are standardized to a 1 May parturition date. The mean adult alkaline phosphatase value and 95% confidence interval is given in the shaded region.

Total WBC counts were higher in males than females in our sample ($P < 0.05$). Pen-raised coyotes⁷ and captive wild coyotes² had significantly lower total WBC counts compared with our

adult sample ($P < 0.001$). Male coyotes we sampled generally were more active and aggressive in the trap and during handling than females, resulting in higher stress and increased muscle activity

TABLE 4. Summary of significant age and sex difference in blood values of coyotes in northern Wisconsin, 1978-79.

Assay	Factor	Relative magnitude of blood values in group	Significance probability
Packed cell volume	Age	Adults > Juveniles	P<0.05
White blood cells	Sex	Males > Females	P<0.05
Albumin	Sex × Age	interaction	P<0.05
Calcium	Sex × Age	interaction	P<0.05
Calcium	Age	Juveniles > Adults	P<0.05
Inorganic phosphorus	Age	Juveniles > Adults	P<0.001
Alkaline phosphatase	Age	Juveniles > Adults	P<0.001

reflected in higher WBC counts. These elevated WBC counts probably represent 'stressed' neutrophilia due to capture stress and resulting splenic contraction.⁸ Wild coyotes we sampled were likely to be more active and stressed than conditioned captive coyotes, again resulting in the observed elevated WBC counts in our sample.

Higher packed cell volumes in adults compared with juveniles ($P < 0.05$) may reflect nutritional differences between these age cohorts. However, capture stress may also have contributed to this difference. Seal *et al.*⁹ suggested that elevated hematological values may result from handling stress in white-tailed deer (*Odocoileus virginianus*).

No significant sex or age effects were found in hemoglobin concentrations in coyotes we sampled. Mean hemoglobin values for pen-raised coyotes were significantly higher than for wild coyotes ($P < 0.001$). Lower hemoglobin concentrations in wild coyotes likely reflect a lower nutritional plane for these animals. Seal and Hoskinson¹¹ reported that hematology values may reflect long term nutritional status in ungulates. Gates and Goering reported hemoglobin values for wild captive coyotes that were conditioned a minimum of 3 months prior to sampling. Hemoglobin values and packed cell volumes reported for these captive wild coyotes did not differ significantly from values we observed. Again, long term nutrition influencing

these values may explain this lack of difference.

Juvenile coyotes had higher serum calcium levels than adults ($P < 0.05$). However, a significant sex × age interaction for calcium levels was determined by analysis of variance ($P < 0.05$). Calcium values were higher for pen-raised coyotes than for adult wild coyotes we sampled ($P < 0.05$). Seal *et al.*¹⁰ reported lower calcium values for wild wolf (*Canis lupus*) pups than dogs (*Canis familiaris*), and attributed this difference to nutrition. The major factor for lower calcium levels in wild coyotes compared with pen-raised animals may also be nutritional.

Juvenile coyotes had significantly higher inorganic phosphorus levels than adults ($P < 0.001$). Phosphorus absorption correlates positively with calcium concentrations and is important in bone metabolism.¹³

No significant sex or age differences were found in serum glucose levels among coyotes we sampled from northern Wisconsin. However, adult coyotes we sampled did have significantly higher serum glucose levels than pen-raised coyotes⁷ ($P < 0.001$). Elevated glucose levels likely reflected transitory hyperglycemia caused by increased adrenal activity in response to capture stress.¹ Seal *et al.*¹⁰ noted that high serum glucose levels in wolves, captured and sampled by methods similar to ours, may be attributable to capture stress. Capture effects have also been related to

high glucose levels in pronghorn antelope (*Antilocapra americana*).¹¹

No significant age or sex effects were found for serum cholesterol, serum urea nitrogen (BUN), and total bilirubin.

There were no significant differences between age or sex cohorts examined for total serum protein or globulin values. Georing *et al.*⁴ reported that captive wild coyotes had lower total protein and albumin values than dogs, and males had higher total protein levels than females. Globulin levels were higher in pen-raised coyotes⁷ than in wild coyotes we sampled ($P < 0.01$). A significant sex \times age interaction was determined for albumin values by analysis of variance ($P < 0.05$). The pattern of albumin values for each age class differed in males and females. Elevated albumin levels may occur with shock or dehydration, while low protein intake may result in depressed albumin values.¹ Adult wild coyotes we sampled had higher serum albumin levels than pen-raised coyotes⁷ ($P < 0.01$). Pen-raised coyotes were maintained on commercial dog food⁷ and, therefore, lower albumin values likely reflect a lower protein intake compared with wild coyotes.

Juvenile coyotes had significantly higher serum alkaline phosphatase levels than adults ($P < 0.001$). These levels decreased curvilinearly with time up to about 8 months of age (Figure 1). The equation for the curve fitted to juvenile alkaline phosphatase levels, given in Figure 1, is $Y = 634 - 28.8x + .37x^2$, where x is juvenile coyote age (weeks), and Y is serum alkaline phosphatase (mU/ml). Fitting this curve

to decreasing juvenile alkaline phosphatase levels reduced the residual variance about the line from $s^2=2486$ using a linear model to $s^2=1143$.

The mean adult alkaline phosphatase level for coyotes we sampled was 40.8 mU/ml with a standard deviation of 18.0 mU/ml. The juvenile alkaline phosphatase curve does not intersect the adult 95% confidence interval until about January, assuming a 1 May parturition date. Therefore, serum alkaline phosphatase levels for coyotes captured in autumn are useful for separating juveniles and yearlings when animals are to be released for research. Some juvenile coyotes are adult weight by October and tooth wear may not always be used as a practical aging tool. Alkaline phosphatase levels are not affected by capture stress,¹¹ and are directly related to the rate of bone formation and osteoblast differentiation.¹³ Coyotes attain adult size and weight within their first year,³ by which time major bone formation should be complete.

Although some blood parameters such as leukocyte counts and glucose levels may reflect capture stress and may differ between sex and age cohorts, other blood values such as hemoglobin may provide a useful measure of long term nutrition. The relationship between serum alkaline phosphatase levels and juvenile coyote age may be integrated with measurements of coyote weight, tooth wear, and body size to reduce ambiguity in field techniques for separating juvenile and yearling coyotes, prior to January.

Acknowledgements

The authors thank R. Adamcik, M. Backes, M. Foy, E. Haug, M. Korb (deceased), C. Morgan, R. Rolley, K. Smith, R. Kennedy, and A. Kennedy for assistance with field work. We thank J. Cary for statistical advice and the personnel of the Central Animal Health Lab, Madison. Research was supported by the College of Agriculture and Life Sciences, Univ. of Wisconsin, Madison, and by the Patuxent Wildlife Research Center (U.S. Fish and Wildlife Service). This paper is dedicated to the memory of Mark J. Korb.

LITERATURE CITED

1. COLES, E.H. 1974. *Veterinary Clinical Pathology*. 2nd ed. W.B. Saunders Co., Philadelphia, Pa. 615pp.
2. GATES, N.L. and E.K. GOERING. 1976. Hematologic values of conditioned, captive wild coyotes. *J. Wildl. Dis.* 12: 402-404.
3. GIER, H.T. 1968. Coyotes in Kansas. *Kans. State Coll. Agric. Exp. Stn. Bull.* 393. 118pp.
4. GOERING, E.K., C.S. CARD, D.F. BROBST and N.L. GATES. 1976. Electrophoretic protein analysis in the conditioned captive wild coyote. *J. Wildl. Dis.* 12: 498-503.
5. KNOWLTON, F.F. 1972. Preliminary interpretations of coyote population mechanics with some management implications. *J. Wildl. Manage.* 36: 369-382.
6. LERESCHE, R.E., U.S. SEAL, P.D. KARNS and A.W. FRANZMANN. 1974. A review of blood chemistry of moose and other Cervidae with emphasis on nutritional assessment. *Nat. Can.* 101: 263-290.
7. RICH, J.E. and N.L. GATES. 1979. Hematologic and serum chemistry values of pen-raised coyotes. *J. Wildl. Dis.* 15: 115-119.
8. SCHALM, O.W., N.C. JAIN and E.G. CARROLL. 1975. *Veterinary Hematology*. Lea and Febiger Co., Philadelphia, Pa. 807pp.
9. SEAL, U.S., J.J. OZOGA, A. W. ERICKSON and L.J. VERME. 1972. Effects of immobilization on blood analysis of white-tailed deer. *J. Wildl. Manage.* 36:1034-1040.
10. ———, L.D. MECH and V. VAN BALLEMBERGHE. 1975. Blood analyses of wolf pups and their ecological and metabolic interpretation. *J. Mammal.* 56: 64-75.
11. ——— and R.L. HOSKINSON. 1978. Metabolic indicators of habitat condition and capture stress in pronghorns. *J. Wildl. Manage.* 42: 755-763.
12. ———, M.E. NELSON, L.D. MECH and R.L. HOSKINSON. 1978. Metabolic indicators of habitat differences in four Minnesota deer populations. *J. Wildl. Manage.* 42:746-754.
13. SEARCY, R.L. 1969. *Diagnostic Biochemistry*. McGraw-Hill Co., New York, N.Y. 660pp.
14. SOKAL, R.R. and F.J. ROHLF. 1973. *Introduction to Biostatistics*. W.H. Freeman and Co., San Francisco, Ca. 368pp.

Received for publication 26 February 1980