

The Occurrence of *Trichomonas gallinae* in White-winged Doves in Arizona

Authors: TOEPFER, E. W., and LOCKE, L. N.

Source: Bulletin of the Wildlife Disease Association, 2(1) : 13

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-2.1.13>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The Occurrence of *Trichomonas gallinae* in White-winged Doves in Arizona

E. W. TOEPFER¹ AND L. N. LOCKE

Patuxent Wildlife Research Center, Laurel, Maryland

L. H. BLANKENSHIP

Migratory Bird Populations Station, Laurel, Maryland

Received for publication 23 August 1965

The occurrence of *Trichomonas gallinae* in white-winged doves (*Zenaida a. asiatica*) in Texas was reported by Locke and Kiel in 1960 (Proc. Helm. Soc. Wash., 27: 128) and by Stabler in 1961 (J. Parasitol., 47: 309-311). During 1963, a survey was conducted near Tucson, Arizona, to obtain data on the incidence of *Trichomonas* in mourning doves (*Zenaidura macroura*). In the process several isolations of *Trichomonas gallinae* were made from white-winged doves (*Zenaida asiatica mearnsi*). These are apparently the first records of this flagellate from white-winged doves in Arizona.

Thirty-two of 42 white-winged doves examined between June and

August 1963 by a throat swabbing and culture technique (Toepfer, 1964. J. Parasitol., 50:703) were found to be infected with *T. gallinae*. All but two of 23 swabbings obtained from nestling white-winged doves were positive. No typical trichomonad cankers were seen in any of the white-winged doves. Most of these doves were trapped in desert habitat in ranching country 20 miles from Tucson. Only one flock of 3 or 4 pigeons (*Columba livia*) frequented the area.

The levels of infection reported herein indicate the minimal level of infection present and should not be regarded as measures of the true incidence of infection.

¹Deceased