



## **Carcass Temperature in Relation to Botulism**

Author: Wobeser, G.

Source: Journal of Wildlife Diseases, 24(1) : 184

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-24.1.184>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## LETTER TO THE EDITOR . . .

### Carcass Temperature in Relation to Botulism

**G. Wobeser**, Department of Veterinary Pathology, Western College of Veterinary Medicine, University of Saskatchewan, Saskatoon, Saskatchewan, Canada S7N 0W0

In 1984, we described greatly elevated temperatures within decaying duck carcasses and suggested that this feature of the carcass microenvironment provided a "milieu" favourable for growth and toxin production by *Clostridium botulinum* during cool weather (Wobeser and Galmut, 1984). At the time I was unaware of any other report of this phenomenon. I have recently found a study in which carcass temperatures were measured. Payne (1965) used neonatal pig carcasses for the study of invertebrates associated with carrion. As occurred in ducks, the internal temperature of the pig carcasses rose much above that of either soil or air on day 3, and then as the flesh disappeared, declined to approximate that of the soil. Payne (1965) attributed the high internal temperature "to the high metabolic rates of the bacteria and dipterous larvae present."

The maximum temperature recorded in a pig carcass was 37.7 C, which is lower than that found in duck carcasses, but this may be a result of the difference in insulative value of pelage and plumage, respectively. These observations strengthen the potential importance of vertebrate carcasses as substrate for botulism toxin production, as the carcass may provide a microenvironment largely independent of ambient conditions.

#### LITERATURE CITED

- PAYNE, J. A. 1965. A summer carrion study of the baby pig *Sus scrofa* Linnaeus. *Ecology* 46: 592-602.
- WOBESER, G., AND E. GALMUT. 1984. Internal temperature of decomposing duck carcasses in relation to botulism. *Journal of Wildlife Diseases* 20: 267-271.

*Received for publication 31 August 1987.*