

Use of Xylazine Sedation with Yohimbine Antagonism in Captive Gray Wolves

Authors: Kreeger, Terry J., Seal, Ulysses S., Callahan, Margaret, and Beckel, Mark

Source: Journal of Wildlife Diseases, 24(4) : 689-690

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-24.4.689>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SHORT COMMUNICATIONS

Journal of Wildlife Diseases, 24(4), 1988, pp. 689-690
© Wildlife Disease Association 1988

Use of Xylazine Sedation with Yohimbine Antagonism in Captive Gray Wolves

Terry J. Kreeger,¹ Ulysses S. Seal,² Margaret Callahan,² and Mark Beckel,² ¹ Department of Fisheries and Wildlife, University of Minnesota, St. Paul, Minnesota 55108, USA; ² Veteran's Administration Medical Center, Minneapolis, Minnesota 55417, USA

ABSTRACT: Captive gray wolves (*Canis lupus*) were given 2.2 mg/kg xylazine hydrochloride intramuscularly resulting in profound sedation in 9.1 ± 0.6 min (mean \pm SE). Heart rate was 42.0 ± 1.0 beats per minute and respiratory rate was 20.1 ± 1.6 respirations per minute during sedation. A variety of manipulations could be performed on sedated animals in relative safety. Thirty min after xylazine administration, the animals were given either 0.15 mg/kg yohimbine hydrochloride or 5% dextrose solution intravenously causing recovery in 5.3 ± 1.0 and 97.1 ± 14.0 min, respectively ($P < 0.001$).

Key words: Gray wolves, *Canis lupus*, xylazine, yohimbine, chemical immobilization, experimental study.

Xylazine hydrochloride (XYL) is an α_2 -adrenergic agonist that has been used singly or as an adjunct to other drugs for the immobilization of many wild species (Bauditz, 1972; Roughton, 1975; Philo, 1978). Ungulates appear to be more sensitive to this drug than do carnivores; domestic cattle, for instance, require one-tenth the dose per kg for sedation than do dogs or cats (Knight, 1980). Although xylazine does not technically cause anesthesia (analgesia and unconsciousness), it can cause profound sedation characterized by hypotension, bradycardia, respiratory depression, analgesia and muscle relaxation (Knight, 1980). Xylazine has not been widely used alone in wild carnivores because sedation can be prolonged and sedated animals can be aroused unpredictably (Booth, 1982).

Yohimbine hydrochloride (YOH) is an α_2 -adrenergic antagonist which has been used to antagonize XYL or the XYL

component of drug combinations in many wild species (Jessup et al., 1983, 1985; Hsu and Shulaw, 1984; Mech et al., 1985; Kreeger and Seal, 1986a; Kreeger et al., 1987). Yohimbine markedly reduces recovery times for wolves immobilized with XYL and ketamine hydrochloride (KET) while having minimal physiological or behavioral alterations when used at recommended doses (Kreeger et al., 1987).

Philo (1978) used XYL to chemically restrain captive wolves (*Canis lupus*), but did not evaluate YOH. Our study reports on the efficacy and uses of XYL with YOH antagonism in the management of captive wolves.

This study took place in October 1987 in east central Minnesota. Thirteen adult wolves (seven female, six male) were used. The husbandry of these animals has been previously reported (Kreeger et al., 1987). All animals were given 2.2 mg/kg (1.0 mg/lb) XYL (Rompun, Haver-Lockhart Laboratories, Shawnee, Kansas 66201, USA) intramuscularly (i.m.) via pole syringe. Injections were in the dorsal musculature of the hindlimb. Thirty min after XYL administration, six animals (three female, three male) received 0.15 mg/kg YOH (Sigma Chemical Co., St. Louis, Missouri 63178, USA) and seven animals (four female, three male) received an equal volume of 5% dextrose (control) in the cephalic vein. Preparation of YOH has been previously reported (Kreeger et al., 1987). Test and control animals were selected at random.

Just prior to administration of YOH or 5% dextrose, heart rates (HR) and respiratory rates (RR) were determined by auscultation and recorded. Induction and recovery times also were recorded. Induction time was defined as the time from injection of XYL to recumbency with eyes closed; recovery was the time from injection of either YOH or dextrose to return to normal motor function, alertness and appropriate responses to stimuli. We were unable to record physiological parameters after YOH administration.

Statistical analyses were by one- and two-way ANOVA (Number Cruncher Statistical Systems, Kaysville, Utah 84307, USA). Statistical significance was determined at $P \leq 0.05$. Means are reported with standard errors (SE).

Induction time was 9.1 ± 0.6 min for all animals. Sedation was usually characterized by sternal or lateral recumbency, eyes closed, regular but shallow respirations and variable responses to sound stimuli. Heart rate was 42.0 ± 1.0 beats per minute (bpm) and respiratory rate was 20.1 ± 1.6 respirations per minute (rpm) 30 min after injection of XYL. Wolves receiving YOH recovered significantly faster than did control animals (5.3 ± 1.1 versus 97.1 ± 14.2 min; $P < 0.001$). There were no differences in induction times, physiological indices, or recovery times between sexes ($P > 0.05$).

Sedated wolves could be approached and handled safely if no loud noises or abrupt contact with the animal were made. Intravenous injections usually caused no response. If necessary, a muzzle could be gently placed on the animal without arousal. Heart rate of sedated wolves (42.0 ± 1.0 bpm) was lower than that of unanesthetized, sleeping wolves (54.0 ± 5.0 bpm) as measured by radiotelemetry (T. J. Kreeger, unpubl. data). This is consistent with the bradycardic properties of XYL (Klide et al., 1975). Respiratory rate was similar to those of undrugged, sleeping wolves (19 ± 4 rpm) (T. J. Kreeger, unpubl. data).

Philo (1978) used a higher dose of XYL (>3.1 mg/kg) than used in this study to sedate wolves, but induction times, level of sedation and recovery times were similar. Respiratory rates, however, were lower. He reported a mean respiratory rate of 15 rpm with periods of apnea of ≥ 30 sec. Decreased respiratory rate was not observed at the doses we used.

We have used XYL and YOH for a variety of animal management purposes: translocating animals, taking blood and urine samples, cleaning and treating wounds, administering vaccinations and other medications, and attaching radio collars. If gross manipulation of the animal is required, a muzzle should always be used. If an animal becomes aroused, ceasing any manipulations and remaining quiet usually results in the animal reverting to profound sedation. The use of XYL sedation is best reserved for short procedures (<15 min) requiring moderate handling of the wolf. For situations involving deep pain (i.e., surgery) or lengthy procedures, we recommend that complete anesthesia be induced with KET and XYL. The main advantage of using XYL alone is the rapid and complete recovery upon antagonism. Using KET and XYL offers total immobilization, but recoveries are lengthier and characterized by ataxia and disorientation (Kreeger et al., 1987). Recoveries are further complicated and lengthened if boosters of KET are required to maintain anesthesia. Yohimbine does not fully antagonize the KET component of this combination resulting in the residual effects of KET probably prolonging recovery (Kreeger and Seal, 1986b).

We feel that the use of XYL sedation with YOH antagonism can be useful for management of captive wolves. Xylazine produces a tractable animal with minimal physiological alterations which can be quickly returned to normalcy when desired by the use of YOH.

We would like to thank Dr. L. David Mech for his review of this manuscript; the U.S. Fish and Wildlife Service for the

use of the wolves; and the Minnesota Department of Natural Resources for the use of their facilities. This work was supported by the Veteran's Administration Medical Center, Minneapolis, Minnesota.

LITERATURE CITED

- BAUDITZ, R. 1972. Sedation, immobilization and anaesthesia with Rompun in captive and free-living wild animals. *Veterinary Medical Review* 3: 204-226.
- BOOTH, N. H. 1982. Nonnarcotic analgesics. In *Veterinary pharmacology and therapeutics*, N. H. Booth and L. E. McDonald (eds.). Iowa State University Press, Ames, Iowa, p. 314.
- HSU, W. H., AND W. P. SHULAW. 1984. Effect of yohimbine on xylazine-induced immobilization of white-tailed deer. *Journal of the American Veterinary Medical Association* 185: 1301-1303.
- JESSUP, D. A., W. E. CLARK, P. A. GULLET, AND K. R. JONES. 1983. Immobilization of mule deer with ketamine and xylazine, and reversal of immobilization with yohimbine. *Journal of the American Veterinary Medical Association* 183: 1339-1340.
- , K. JONES, R. MOHR, AND T. KUCERA. 1985. Yohimbine antagonism to xylazine in free-ranging mule deer and bighorn sheep. *Journal of the American Veterinary Medical Association* 187: 1251-1253.
- KLIDE, R. J., H. W. CALDERWOOD, AND L. R. SOMA. 1975. Cardiopulmonary effects of xylazine in dogs. *American Journal of Veterinary Research* 36: 931-935.
- KNIGHT, A. P. 1980. Xylazine. *Journal of the American Veterinary Medical Association* 176: 454-455.
- KREEGER, T. J., A. M. FAGGELLA, U. S. SEAL, L. D. MECH, M. CALLAHAN, AND B. HALL. 1987. Cardiovascular and behavioral responses of gray wolves to ketamine-xylazine immobilization and antagonism by yohimbine. *Journal of Wildlife Diseases* 23: 463-470.
- , AND U. S. SEAL. 1986a. Immobilization of coyotes with xylazine hydrochloride-ketamine hydrochloride and antagonism by yohimbine hydrochloride. *Journal of Wildlife Diseases* 22: 604-606.
- , AND ———. 1986b. Failure of yohimbine hydrochloride to antagonize ketamine hydrochloride immobilization of gray wolves. *Journal of Wildlife Diseases* 22: 600-603.
- MECH, L. D., G. D. DEL GIUDICE, P. D. KARNS, AND U. S. SEAL. 1985. Yohimbine hydrochloride as an antagonist to xylazine hydrochloride-ketamine hydrochloride immobilization of white-tailed deer. *Journal of Wildlife Diseases* 21: 405-410.
- PHILO, L. M. 1978. Evaluation of xylazine for chemical restraint of captive arctic wolves. *Journal of the American Veterinary Medical Association* 173: 1163-1166.
- ROUGHTON, R. D. 1975. Xylazine as an immobilizing agent for captive white-tailed deer. *Journal of the American Veterinary Medical Association* 167: 574-576.

Received for publication 26 January 1988.