



## **Natural Infections of *Clinostomum complanatum* (Trematoda: Clinostomatidae) in Wild Herons and Egrets, Tottori Prefecture, Japan**

Authors: Aohagi, Y., Shibahara, T., Machida, N., Yamaga, Y., Kagota, K., et al.

Source: Journal of Wildlife Diseases, 28(3) : 470-471

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.3.470>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Natural Infections of *Clinostomum complanatum* (Trematoda: Clinostomatidae) in Wild Herons and Egrets, Tottori Prefecture, Japan

Y. Aohagi,<sup>1</sup> T. Shibahara,<sup>1</sup> N. Machida,<sup>2,4</sup> Y. Yamaga,<sup>2</sup> K. Kagota,<sup>2</sup> and T. Hayashi,<sup>3</sup> <sup>1</sup> Laboratory Animal Research Center, Tottori University School of Medicine, Nishimachi, Yonago 683, Japan; <sup>2</sup> Department of Veterinary Internal Medicine, Faculty of Agriculture, Tottori University, Koyama, Tottori 680, Japan; <sup>3</sup> University Veterinary Hospital, Faculty of Agriculture, Tottori University, Koyama, Tottori 680, Japan. <sup>4</sup> Present address: Department of Veterinary Pathology, Faculty of Agriculture, Tokyo University of Agriculture and Technology, Fuchu 183, Japan

**ABSTRACT:** *Nycticorax nycticorax*, *Ardea cinerea*, *Egretta garzetta*, and *Egretta intermedia* were naturally infected with *Clinostomum complanatum* (Trematoda: Clinostomatidae) among fourteen wild herons, seven wild egrets and one wild bittern evaluated at the Veterinary Hospital of Tottori University, Tottori, Japan. The latter three species of heron and egrets are reported for the first time as definitive hosts of this parasite in Japan.

**Key words:** *Clinostomum complanatum*, natural infection, *Nycticorax*, *Egretta*, *Ardea*, wild herons, wild egrets, Japan.

Adult *Clinostomum complanatum* are found in the oral cavity and pharynx of many fish-eating birds. While a few species of the birds have been recorded as definitive hosts of this fluke in Asia (Lo et al., 1981), only night herons, *Nycticorax nycticorax* have been identified as definitive hosts in Japan (Yamaguti, 1933; Kagei et al., 1988). Four human cases of *C. complanatum* infection have been reported in Japan (Yamashita, 1938; Hirai et al., 1987; Umegai et al., 1990; Yoshimura et al., 1991). Adult flukes were recovered from the pharynx of the patients. Our objective was to determine the prevalence of *C. complanatum* among herons, egrets and bittern from a site in Japan.

From April 1987 to March 1991, 22 birds from the eastern part of Tottori Prefecture (134°00' to 134°50'E, 35°10' to 35°35'N) were brought to the Veterinary Hospital of Tottori University, Tottori, Japan: nine night herons, five grey herons (*Ardea cinerea*), five little egrets (*Egretta garzetta*), two intermediate egrets (*Egretta intermedia*), and one Chinese little bittern (*Ixobrychus sinensis*).

The herons, egrets and bittern survived between 1 and 35 days. Medical treatment was limited to treatment of injuries. No anthelmintic drugs were administered. The birds were fed fresh squid and small marine fish in the course of medical treatment; they never were fed any freshwater fish which could serve as second intermediate hosts of *C. complanatum*.

The complete digestive tract of each dead bird was examined for *C. complanatum* infection. The recovered worms were fixed in 70% alcohol under pressure of a cover glass, stained with Borax-carmin, dehydrated, cleared in xylene and mounted in balsam. Eggs were preserved in 10% formalin. These specimens were measured and observed by light microscopy. All the worms were identified as *C. complanatum*.

The worms were found in five of nine night herons, all five grey herons, two of five little egrets, and one of two intermediate egrets. No worms were recovered from the Chinese little bittern. Intensity ranged from 1 to 139, with a median of 6 worms per infected bird. Of 248 worms recovered from the pharynx of these birds, 210 were adult flukes with eggs in their uteri and uterine-sacs. The remaining 38 flukes were immature.

In the eastern part of Tottori Prefecture, six species of freshwater fishes were known as second intermediate hosts of *C. complanatum* (Aohagi et al., 1992). The birds in our study probably became infected with *C. complanatum* after feeding on infected fish in the area.

Many genera of birds have been re-

ported as definitive hosts of *C. complanatum* (Lo et al., 1982). All infected birds in this survey fell within those genera. However, three species of birds (*A. cinerea*, *E. garzetta*, *E. intermedia*) are recorded as definitive hosts of *C. complanatum* for the first time in Japan.

Representative specimens were deposited in the Laboratory Animal Research Center, Tottori University School of Medicine, Japan (CA8704001-9072003).

The authors are grateful to Professor K. Kawashima, Kyushu University School of Health Sciences, Japan, for a critical reading of the manuscript. We also are thankful to Professor M. Kamiya, Department of Parasitology, Faculty of Veterinary Medicine, Hokkaido University, for his valuable comments.

#### LITERATURE CITED

- AOHAGI, Y., T. SHIBAHARA, N. MACHIDA, Y. YAMAGA, AND K. KAGOTA. 1992. *Clinostomum complanatum* (Rudolphi, 1814) (Trematoda: Clinostomatidae) in five new fish hosts in Japan. *Journal of Wildlife Diseases* 28: In Press.
- HIRAI, H., H. OISO, T. KIFUNE, T. KIYOTA, AND Y. SAKAGUCHI. 1987. *Clinostomum complanatum* infection in posterior wall of the pharynx of a human. *Japanese Journal of Parasitology* 36: 142-144.
- KAGEI, N., Y. YANOHARA, R. UCHIKAWA, AND A. SATO. 1988. Natural infection with *Clinostomum complanatum* (Rud., 1819) in the birds of southern Japan. *Japanese Journal of Parasitology* 37: 254-257.
- LO, C. F., F. HUBER, G. H. KOU, AND C. J. LO. 1981. Study of *Clinostomum complanatum* (Rud., 1819). *Fish Pathology* 15: 219-227.
- , C. H. WANG, F. HUBER, AND G. H. KOU. 1982. The study of *Clinostomum complanatum* (Rudolphi, 1814) II. The life cycle of *Clinostomum complanatum*. CAPD Fisheries Series No. 8, Reports on Fish Disease Research (IV) 26-56.
- UMEGAI, T., T. SHIN, M. ODA, T. KIFUNE, AND M. MOGI. 1990. A case of acute laryngitis caused by *Clinostomum complanatum* with a complaint of throat irritation. *Jibi to Rinsho* 36: 665-668 (in Japanese).
- YAMAGUTI, S. 1933. Studies on the helminth fauna of Japan. Part 1. Trematodes of birds, reptiles and mammals. *Japanese Journal of Zoology* 5: 66-71.
- YAMASHITA, J. 1938. *Clinostomum complanatum*. A trematode parasite new to man. *Annotationes Zoologicae Japonenses* 17: 563-566.
- YOSHIMURA, K., S. ISHIGOOKA, I. SATOH, AND S. KAMEGAI. 1991. *Clinostomum complanatum* from the pharynx of a woman in Akita, Japan. *Japanese Journal of Parasitology* 40: 99-101.

Received for publication 30 August 1991.