

## Knemidocoptes sp. on Wild Passerines at the Mai Po Nature Reserve, Hong Kong

Authors: Mainka, Susan A, Melville, David S., Galsworthy, Antony, and Black, S. R.

Source: Journal of Wildlife Diseases, 30(2): 254-256

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-30.2.254

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## SHORT COMMUNICATIONS

Journal of Wildlife Diseases, 30(2), 1994, pp. 254-256 © Wildlife Disease Association 1994

## *Knemidocoptes* sp. on Wild Passerines at the Mai Po Nature Reserve, Hong Kong

Susan A Mainka,<sup>1</sup> David S. Melville,<sup>2</sup> Antony Galsworthy,<sup>2</sup> and S. R. Black,<sup>3</sup> <sup>1</sup> World Wide Fund for Nature International, 1196 Gland, Switzerland; <sup>2</sup> World Wide Fund for Nature Hong Kong, GPO 12721, Central, Hong Kong; <sup>3</sup> Calgary Zoological Society, P.O. Box 3036, Station B, Calgary, AB, Canada, T2M 4R8

ABSTRACT: Free-ranging passerine birds banded at the Mai Po Nature Reserve, Hong Kong, were affected with a skin disease characterized by crusty, proliferative lesions on legs, feet and beaks. Based on retrospective examination of 1990 to 1992 banding records, 83 of 16,353 birds and 5 of 161 species banded at Mai Po were identified as having these lesions. Eurasian tree sparrows (Passer montanus) were most commonly affected (58 of 411 birds) although 19 of 428 black-faced buntings (Emberiza spodocephala), 1 of 43 little buntings (Emberiza pusilla), 3 of 92 spotted doves (Streptopelia chinensis), and 2 of 616 eastern great reed warblers (Acrocephalus orientalis) also had clinical signs of infestation. Adult Eurasian tree sparrows were affected more often than juveniles and males were affected more often than females. Lesions on juveniles usually were on the feet while on adults lesions also were found on legs and beak. The effects of this parasitic infestation on body weight of wild Eurasian tree sparrows was not significant.

Key words: Ectoparasite, Knemidoctoptes sp., Eurasian tree sparrow, Passer montanus.

Infestation of free-ranging birds by Knemidocoptes sp. mites is uncommon. Reports of this disease in one great horned owl (Bubo virginianus) (Schulz et al., 1989), two sedge warblers (Acrocephalus schoenobaenus) (Fry et al., 1969), a chaffinch (Fringilla coelebs) (McDonald, 1962), and a snowy owl (Nyctea scandiaca), and accipters in Great Britain (Cooper, 1985) have been made. In these cases, other than scaly lesions, effects on the host were minimal. Kirmse (1966) reported the prevalence of mite infestation among affected species of birds ranging from 40% in red-winged blackbirds (Agelaius phoenicus) to 1% in cowbirds (*Molothrus ater*). We examined the prevalence of mites in the free-ranging population of birds at the World Wide Fund for Nature (WWF) Hong Kong Mai Po Nature Reserve, Hong Kong (22°30'N, 114°02'W). Effects of this parasitism as reflected in body weight also were analyzed.

Banding records from the Mai Po Nature Reserve for 1990 through 1992 were retrospectively examined for comments on prevalence of scaly, crusty skin lesions. During routine banding sessions, mist nets were set up at regular locations and birds caught in the nets were removed and banded. We recorded estimated age based on plumage characteristics, sex by cloacal protuberance, and body weight. Presence and location of any dry, scaly, crusty lesions were noted. Smooth, bulbous swellings of the toes of some birds occasionally were seen but were not included in this study. During 1992 skin scrapings were done on several birds with severe lesions. Two of the affected birds which died were necropsied and samples of lesions were fixed in 10% buffered formalin. Fixed tissues were embedded in paraffin and sectioned at 3  $\mu$ m and stained with hematoxylin and eosin for examination by light microscopy.

An analysis of species affected, sex, age class and body part affected was done. Statistical analysis of effect of mite infestation on body weight was done using a Student *t*-test (Kwikstat 3.01©, TexaSoft 1984– 1992, Cedar Hill, Texas, USA). Significance was assessed at a level of P = 0.05.

	Total birds examined	Mite-infested birds number (%)
Number of birds banded Number of species banded	16,353 161	106 (0.65) 5 (3.1)
Passer montanus banded	411	58 (14)
Males	87	23 (26)
Females <sup>.</sup>	84	13 (16)
Juveniles	196	16 (8.2)
Adults	215	40 (19)
Emberiza spodocephala	428	19 (4.4)
Streptopelia chinensis	92	3 (3.2)
Emberiza pusilla	43	1 (2.3)
Acrocephalus orientalis	616	2 (0.3)

TABLE 1.Prevalence (%) of Knemidocoptes sp. miteson free-ranging bird populations at the Mai Po NatureReserve, Hong Kong, 1990 through 1992.

TABLE 2.Distribution of scaly crusty lesions on wildEurasian tree sparrows (*Passer montanus*) banded atthe Mai Po Nature Reserve, Hong Kong.

Group	Foot	Leg	Beak
All sparrows $(n = 411)$	40	32	2
Males $(n = 87)^{\circ}$	9	15	1
Females $(n = 84)$ .	6	8	0
Juveniles $(n = 196)$	18	2	0
Adults $(n = 215)$	21	29	2

\* Sex was not determined for all birds.

asian tree sparrows identified as males and females, respectively, were banded; however, a preponderance of infection in male birds was noted (Table 1). Based on a chisquare test, these differences were not significant. Similarly, the percentage of adult and juvenile birds banded was approximately equal; however, using a chi-square test, there was a significantly (P = 0.007) higher prevalence of mite infestation in adults (Table 1). The increased prevalence in adults may have been partly related to older birds having more time to develop clinically recognizable lesions.

Distribution of mite lesions appeared to progress from foot lesions in juveniles to leg lesions and occasionally beak lesions in adults (Table 2). There were no significant differences in the body weights of infected and uninfected Eurasian tree sparrows, when corrected for age and sex differences.

The WWF Hong Kong banding program is sponsored by the Rotary Club of Hong Kong South. We would like to thank the pathology staff of the Alberta Agriculture Lab in Airdrie, Alberta, Canada for their assistance with section preparation.

## LITERATURE CITED

- COOPER, J. E. 1985. Veterinary aspects of captive birds of prey. The Standfast Press, Gloucestershire, England, 256 pp.
- FRY, C. H., I. J. FERGUSON-LEES, AND J. S. ASH. 1969. Mite lesions in sedge warblers and beeeaters in Africa. Ibis 111: 611–613.
- KIRMSE, P. 1966. Cnemidocoptic mite infestations in wild birds. Bulletin of the Wildlife Disease Association 2: 86–99.

Sex was not determined for all birds.

Skin scrapings taken from severe lesions confirmed the presence of a mite infestation in affected birds. One Eurasian tree sparrow and one reed warbler were necropsied; both had lesions on the feet and legs consisting of cornified tissue on the plantar surface of the feet and extending partway up the tarsus. These cornified masses were broken at the joint area.

On histopathology, crusty lesions had sections of mites within the epidermis in an area of massive compact orthokeratotic hyperkeratosis. The mass of stratum corneum had a honeycombed appearance and several mites were present in each section. Some mild degree of acanthosis was noted; otherwise deeper layers of the epidermis were not affected. Mites were tentatively identified as *Knemidocoptes* sp. based on the lesions seen grossly and the distribution of mites in burrows of the stratum corneum as seen on histopathological examination. No attempt was made to identify the species.

Eurasian tree sparrows represented only 2.6% of birds banded but were most commonly infested by mites at 14% (Table 1). As with Kirmse (1966), we found that some species of birds had a higher prevalence of infestation than others.

Approximately equal numbers of Eur-

MCDONALD, J. W. 1962. Chaffinch with cnemidocoptic mange. British Birds 55: 421.

SCHULZ, T. A., J. S. STEWART, AND M. E. FOWLER. 1989. Knemidocoptes mutans (Acari: Knemidocoptidae) in a great-horned owl (Bubo virginianus). Journal of Wildlife Diseases 25: 430-432.

Received for publication 2 August 1993.