

## ONE-SAMPLE PREGNANCY DIAGNOSIS IN ELK USING FECAL STEROID METABOLITES

Authors: Garrott, Robert A., Monfort, Steven L., White, P. J., Mashburn, Kendall L., and Cook, John G.

Source: Journal of Wildlife Diseases, 34(1) : 126-131

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-34.1.126>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ONE-SAMPLE PREGNANCY DIAGNOSIS IN ELK USING FECAL STEROID METABOLITES

Robert A. Garrett,<sup>1</sup> Steven L. Monfort,<sup>2</sup> P. J. White,<sup>3</sup> Kendall L. Mashburn,<sup>2</sup> and John G. Cook<sup>4</sup>

<sup>1</sup> Biology Department, Montana State University, 310 Lewis Hall, Bozeman, Montana 59717, USA

<sup>2</sup> Conservation and Research Center, National Zoological Park, Smithsonian Institution, Front Royal, Virginia 22630, USA

<sup>3</sup> Biological Sciences Department, California Polytechnic State University Foundation, San Luis Obispo, California 93451, USA

<sup>4</sup> National Council of the Paper Industry for Air and Stream Improvement, Forestry and Range Sciences Lab, 1401 Gekeler Lane, La Grande, Oregon 97850, USA

**ABSTRACT:** Recent research has demonstrated the potential of pregnancy diagnosis in elk (*Cervus elaphus nelsoni*) using immunoassays of fecal steroid concentration. However, multiple samples are required to insure accurate results, limiting its utility for free-ranging animals. We attempted to develop an accurate one-sample pregnancy diagnosis using 153 fecal samples that were collected from free-ranging, radio-collared, adult female elk in Yellowstone National Park (Wyoming, USA) and from captive elk maintained at the Starkey Research Facility (La Grande, Oregon, USA) February through April 1992 and 1997. The pregnancy status of each animal was diagnosed using serum pregnancy-specific protein B (PSPB) assays providing fecal samples from 38 nonpregnant and 115 pregnant animals. Fecal radioimmunoassay (RIA) indicated that mean ( $\pm$ SD) progestagens ( $P_4$ ) were elevated significantly in pregnant ( $2.96 \pm 1.49 \mu\text{g/gm}$ ) compared to nonpregnant ( $0.43 \pm 0.26 \mu\text{g/gm}$ ) individuals. Confidence intervals ( $1.96 \pm \text{SE}$ ) for the two groups were widely separated (nonpregnant 0.34–0.51, pregnant 2.69–3.24) with little overlap in the range of concentrations measured for each group (nonpregnant 0.09–0.98, pregnant 0.90–8.29). These results indicate that fecal progestagens RIA provides a reliable method of noninvasive pregnancy diagnosis using single fecal samples collected from elk during late gestation. However, independent validation of the suggested discrimination criteria should be performed before routine application.

**Key words:** *Cervus elaphus*, physiology, radioimmunoassay, reproduction, techniques.

### INTRODUCTION

The development of noninvasive techniques for assessing pregnancy in uncaptured ungulates is an active area of research that promises to provide biologists with a new and useful tool for the study of reproduction and neonatal mortality (Messier et al., 1990). Methodologies are based on the measurement of reproductive steroids or their metabolic byproducts excreted in feces using immunoassay procedures (Schwarzenberger et al., 1996). Fecal steroids concentrations have been explored in studies on muskoxen (*Ovibos moschatus*) by Desaulniers et al. (1989), caribou (*Rangifer tarandus*) by Messier et al. (1990), feral horses (*Equus caballus*) by Kirkpatrick et al. (1990) and Lucas et al. (1991), moose (*Alces alces*) by Monfort et al. (1993), bison (*Bison bison*) by Kirkpatrick et al. (1993) and bighorn sheep (*Ovis*

*canadensis*) by Borjesson et al. (1996). All of these studies have demonstrated the potential of diagnosing pregnancy using fecal steroid metabolites. However, the ability to discriminate between pregnant and nonpregnant animals from a single fecal sample has varied from 55 to 100% depending on the species, extraction and assay procedures, steroid metabolites measured, and timing of sample collection during gestation.

In an effort to validate this technique for monitoring the long-term reproductive success of cow elk (*Cervus elaphus nelsoni*) in Yellowstone National Park (Wyoming USA) White et al. (1995) used enzyme immunoassay (EIA) to explore the potential of three steroid metabolites for diagnosing pregnancy. They found that free progesterone ( $P_4$ ) and pregnanediol-3-glucuronide (PdG) concentrations in fecal samples collected during late gestation

(March and April) were the best indicators of pregnancy status. Discrimination from a single sample was only 74 to 84% correct, leading these investigators to recommend assaying multiple samples in order to ensure accurate results. The collection of numerous fecal samples from a single animal will limit the application of the technique to captive animals and relatively tractable, individually identifiable free-ranging animals. The purpose of this study was to determine if a more reliable single-sample pregnancy diagnosis for elk could be obtained by using alternative procedures.

### METHODS

One hundred fifty three fecal samples from adult female elk of known pregnancy status were used in this study. During 12 February to 28 April 1992 fecal samples were collected from 23 individual free-ranging instrumented cow elk occupying the Madison-Firehole winter range in Yellowstone National Park (44°37'N, 110°52'W). An additional 18 animals were sampled 14 March to 8 April 1996 (13 of these animals in common with the 1992 collection). Samples were collected by locating each animal using telemetry and observing the animal until a defecation event was observed (White et al. 1995). One hundred twelve additional fecal samples were collected 8 to 12 April 1996 ( $n = 57$ ) and 10 April, 1997 ( $n = 55$ ) from captive adult female elk maintained at the Starkey Research Facility near La Grande (Oregon, USA; 45°30'N, 118°20'W). All samples from each annual Starkey collection originated from different elk, but the same animals were sampled in both years. Descriptions of the Madison-Firehole area and the Starkey Research Facility are provided in Garrott et al. (1996) and Cook et al. (1996), respectively. The pregnancy status of each animal was diagnosed using serum pregnancy-specific protein B (PSPB; Sasser et al., 1986; Hein et al., 1991). Blood samples were collected from free-ranging elk while they were chemically immobilized for the purposes of attaching radio collars (White et al., 1995). Blood samples from captive elk were collected during December with pregnancy status confirmed at parturition in May and June.

Fecal sample processing and extraction procedures were modified from previously described methods (Wasser et al., 1991, 1994; Monfort et al., 1993; Brown et al., 1994; Schwartz et al., 1995). Fecal samples first were dried using a rotary evaporator and 0.1 g feces

was boiled (20 min) in 16 × 125-mm glass tubes containing 10 ml of absolute ethanol and approximately 2,000 c.p.m. radio-labeled <sup>3</sup>H-progesterone to monitor procedural losses. After boiling, tubes were centrifuged at room temperature (10 min, 1,500 × g) and the ethanol supernate was decanted into clean 16 × 125-mm tubes, evaporated to dryness, and reconstituted in 1 ml methanol. After vortexing, extractants were sonicated (10 min) and 25 µl was withdrawn for radioactive counting to determine extraction efficiency. Fecal P<sub>4</sub> was assayed using an radioimmunoassay (RIA) procedure described previously (Brown et al., 1994; Wasser et al., 1994). Briefly, serial dilutions (range = 1:32 to 1:1024) of a pooled fecal extract from randomly selected female elk of unknown reproductive status yielded a displacement curve parallel to the standard curve. Recovery of known amounts of progesterone (7.5 to 120.0 pg/tube) added to a pool of diluted fecal extract (1:400, 50 µl) was 95% ( $y = -1.78 + 1.04x$ ,  $r^2 = 0.99$ ). Inter-assay coefficients of variation for two separate internal controls were 14% ( $n = 6$ , 79 to 95% binding) and 6% ( $n = 6$ , 43 to 69% binding). Intra-assay coefficients of variation were <5%, and assay sensitivity was 3.75 pg/100 µl. All hormone concentrations are expressed as mass units of hormone excreted per gm of dry feces. A *t*-test was used to evaluate differences in mean fecal P<sub>4</sub> concentrations for pregnant versus nonpregnant animals using STATISTICA (StatSoft, Inc., Tulsa, Oklahoma, USA).

### RESULTS

PSPB assays indicated that 20 of the 23 elk from Yellowstone sampled in 1992 were pregnant, and 16 of the 18 animals sampled in 1996 were pregnant. Forty-three of the 57 animals from Starkey sampled in 1996 and 36 of the 55 animals sampled in 1997 were pregnant. Thus, of the 153 samples collected, 38 were from nonpregnant animals and 115 were from pregnant animals. Thirty seven of 38 samples from nonpregnant animals had fecal P<sub>4</sub> concentrations <1.0 µg/gm, while 112 of 115 samples from pregnant animals had concentrations ≥1.0 µg/gm. One 1996 sample from a nonpregnant animal from Starkey had a P<sub>4</sub> concentration of 1.26 and a single 1997 sample from a pregnant animal from Starkey had a P<sub>4</sub> concentration of 0.64. These two samples appear to be




FIGURE 1. Distribution of progesterone ( $P_4$ ) concentrations ( $\mu\text{g/gm}$  dry fecal weight) in 151 fecal samples collected from nonpregnant ( $n = 37$ ) and pregnant elk ( $n = 114$ ), collected from free-ranging radio-collared elk in Yellowstone National Park (Wyoming USA) and captive animals maintained at the Starkey Research Facility (La Grande, Oregon USA).

outliers and we suspect errors during sample collection and/or labeling. Mean ( $\pm\text{SD}$ ) fecal  $P_4$  concentrations, excluding the two apparent outliers, were  $0.43 \pm 0.26 \mu\text{g/gm}$  and  $2.96 \pm 1.49 \mu\text{g/gm}$  for nonpregnant and pregnant animals, respectively (Fig. 1). Confidence intervals ( $1.96 \pm \text{SE}$ ) for the two groups were widely separated (nonpregnant 0.34 to 0.51, pregnant 2.69 to 3.24), indicating higher concentrations for samples from pregnant animals ( $P < 0.01$ ). Overlap in fecal  $P_4$  concentrations for samples from two pregnant animals and two nonpregnant animals was noted in the 0.9 to 0.98  $\mu\text{g/gm}$  range (preg.—0.90, 0.95; nonpreg.—0.96, 0.98).

#### DISCUSSION

White et al. (1995) demonstrated that fecal concentrations of both PdG and  $P_4$  were significantly higher in samples collected from pregnant elk during late gestation compared to those collected from nonpregnant animals. EIA procedures were employed for measuring fecal steroid concentrations as they felt such assay methodology would be more economical and require less sophisticated laboratory equipment than RIA procedures. However, results of their study indicated that some er-

rors in discrimination between pregnant and nonpregnant animals occurred when pregnancy diagnosis was attempted based on a single fecal sample; false negatives (classifying a nonpregnant animal as pregnant) were the most common error (White et al., 1995). These results led the investigators to conclude that multiple late-gestation fecal samples were needed to accurately assess pregnancy status using the techniques employed in their study. To ascertain whether progesterone metabolites were an inherently poor indicator of pregnancy status, or alternatively if failure to detect pregnancy in a single sample was a consequence of the  $P_4$  EIA employed, we conducted a similar study using an alternative immunoassay. Unlike the previous report (White et al., 1995), the present data indicated that progesterone immunoreactivity provided a reliable single-sample pregnancy test (Fig. 1). However, the fecal  $P_4$  concentrations from two pregnant animals fell within the extreme upper tail of the distribution for nonpregnant animals, indicating a narrow range for potentially ambiguous pregnancy diagnosis.

Selecting a critical  $P_4$  concentration for pregnancy discrimination is a subjective decision and may depend on whether false

negatives or false positives are more critical to the objectives of a given study. We suggest that when employing the methodology of the present study, only late gestation samples (March–April) should be used, with animals having fecal  $P_4$  concentrations  $<0.9 \mu\text{g/gm}$  considered nonpregnant and those with concentrations  $\geq 1.0 \mu\text{g/gm}$  considered pregnant. Discrimination for samples with concentrations between  $0.90\text{--}0.99 \mu\text{g/gm}$  should be regarded inconclusive. Of 105 late-gestation fecal samples collected from radio-collared cow elk in Montana (USA) and Wyoming in separate studies only three fell within the narrow range of  $P_4$  concentrations where diagnosis may be considered uncertain, suggesting that potential ambiguities in diagnosis will be uncommon in field situations. However, we caution that the results of this study and the discrimination criteria suggested should be independently validated before the procedure is routinely applied for pregnancy diagnosis.

Although detailed descriptions of immunoassay methodologies is beyond the scope of this manuscript, the observation that pregnancy diagnosis efficiency varied using two different immunoassays for the same class of hormone warrants discussion. Progesterone, the most biologically potent progestin, can be routinely quantified in blood using any number of specific antisera which are reported to exhibit little or no crossreactivity with other structurally-similar steroids found in blood. However, when progesterone is metabolized and excreted, it is rarely excreted in its native form. Rather, progesterone is generally metabolized to numerous structurally similar steroid molecules, most of which are rather esoteric in strict physiological terms since their structural forms render them biologically impotent. These byproducts of progesterone are primarily important as indirect measures of progesterone in blood circulation. Steroid metabolites can be excreted as conjugated or unconjugated forms; conjugate side-chain moieties such as glucuronides or sulfates are often added during metabolism

to facilitate water solubility in the aqueous environments of bile or urine (Lasley and Kirkpatrick, 1991; Schwarzenberger et al., 1996). The situation is further complicated by the fact that the number and distribution of these metabolites vary between species.

Although fecal hormone monitoring techniques are now widely accepted for wildlife research, there remains a great deal of misunderstanding about these methods. For the wildlife biologist, it is critical to understand that all immunoassays are not created equal. The backbone of any immunoassay is the antiserum employed. Each antiserum has a characteristic specificity, or ability to recognize minute three-dimensional structural differences among similar classes of antigens (i.e., steroid hormones). Because antibody production is based upon polyclonal or monoclonal lymphocyte production, no two clones of lymphocytes are likely to produce antibodies with identical specificities. Whether employed in an EIA or RIA, it is the unique specificity of the antibody, and not whether the ligand is enzyme- or radio-labeled, that fundamentally dictates assay performance. Thus, a  $P_4$  immunoassay used by one laboratory will not necessarily recognize the same 'family' of progestagens as an immunoassay used by another laboratory. One  $P_4$  immunoassay might be highly specific (i.e., recognize progesterone and only progesterone), whereas another  $P_4$  immunoassay might be relatively nonspecific (i.e., recognize progesterone and a host of other structurally-related progestins). Furthermore, some antibodies only recognize unconjugated steroids, whereas others recognize only conjugated forms, or both conjugated and unconjugated forms. Thus, when one laboratory reports that their particular immunoassay does not detect elevated steroid during pregnancy, it does not, therefore, follow that another laboratory's immunoassay will provide a similar answer.

Despite the fact that the antibodies utilized in the present study, and by White et

al. (1995), both presumably are specific for the same class of progesterone metabolites (i.e., 20-oxo-pregnanes), our antiserum appears to provide improved discrimination for pregnancy detection in elk. Clearly, these differences cannot be attributed to the type of immunoassay employed (EIA versus RIA). However, one explanation for this discrepancy is that our antibody may be more specific for major elk pregnancy progestagens than the antiserum employed by White et al. (1995). Alternatively, the extraction method (boiling in ethanol) utilized in the present study may be superior for efficiently extracting elk progestagens compared to the previously employed method (buffer solubilization). In conclusion, it is critical to carefully scrutinize the endocrine methods employed, even when using extraction procedures and/or immunoassays that have been previously reported to be effective for documenting fecal steroid metabolites.

#### ACKNOWLEDGMENTS

Financial support for this study was provided by Montana State University-Bozeman, University of Wisconsin-Madison, the National Park Service, and Montana Department of Fish, Wildlife, and Parks. Financial and logistical support for the captive animals was provided by the National Council of the Paper Industry for Air and Stream Improvement, Northwest Forest Resource Council, Boise Cascade Corporation, U.S. Forest Service Pacific Northwest Forest and Range Experiment Station, Rocky Mountain Elk Foundation, National Fish and Wildlife Foundation, and Oregon Department of Fish and Wildlife. J. McDonald and A. Pils assisted in sample collections in Yellowstone; captive animal maintenance and sample collection was performed by C. Bowers, S. Cerini, and J. Schweitzer. We thank National Park Service personnel R. Abegglen, R. Lindstrum, J. Mack, and J. Varley and Montana Fish, Wildlife, and Parks personnel Kieth Aune, Michelle Kastler, and John Vore for their assistance, encouragement and logistic support. All research activities were carried out in accordance with approved U.S. Forest Service (Wisdom et al. 1993) and Montana State University animal welfare guidelines (protocol #585).

#### LITERATURE CITED

- BORJESSON, D. L., W. M. BOYCE, I. A. GARDNER, J. DEFORGE, AND B. LASLEY. 1996. Pregnancy detection in bighorn sheep (*Ovis canadensis*) using fecal-based enzyme immunoassay. *Journal of Wildlife Diseases* 32: 67-74.
- BROWN, J. L., S. K. WASSER, D. E. WILDT, AND L. H. GRAHAM. 1994. Comparative aspects of steroid hormone metabolism and ovarian activity in felids, measured noninvasively in feces. *Biology of Reproduction* 51: 776-786.
- COOK, J. G., L. J. QUINLAN, L. L. IRWIN, L. D. BRYANT, R. A. RIGGS, AND J. W. THOMAS. 1996. Nutrition-growth relations of elk calves during late summer and fall. *The Journal of Wildlife Management* 60: 528-541.
- DESAULNIERS, D. M., A. K. GOFF, K. J. BETTERIDGE, J. ROWELL, AND P. F. FLOOD. 1989. Reproductive hormone concentrations in faeces during the oestrous cycle and pregnancy in cattle (*Bos taurus*) and muskoxen (*Ovibos moschatus*). *Canadian Journal of Zoology* 67: 1148-1154.
- GARROTT, R. A., P. J. WHITE, D. B. VAGNONI, AND D. M. HEISEY. 1996. Purine derivatives in snow-urine as a dietary index for free-ranging elk. *The Journal of Wildlife Management* 60: 735-743.
- HEIN, R. G., J. L. MUSSER, AND E. F. BRACKEN. 1991. Serologic, parasitic and pregnancy survey of the Colockum elk herd in Washington. *Northwest Science* 65: 217-222.
- KIRKPATRICK, J. F., S. E. SHIDELER, AND J. W. TURNER, JR. 1990. Pregnancy determination in uncaptured feral horses based on free steroids in feces and steroid metabolites in urine-soaked snow. *Canadian Journal of Zoology* 68: 2576-2579.
- , D. F. GUDERMUTH, R. L. FLAGEN, J. C. MCCARTHY, AND B. L. LASLEY. 1993. Remote monitoring of ovulation and pregnancy of Yellowstone bison. *The Journal of Wildlife Management* 57: 407-412.
- LASLEY, B. L., AND J. F. KIRKPATRICK. 1991. Monitoring ovarian function in captive and free-ranging wildlife by means of urinary and fecal steroids. *The Journal of Zoo and Wildlife Medicine* 22: 23-31.
- LUCAS, Z., J. I. RAESIDE, AND K. J. BETTERIDGE. 1991. Non-invasive assessment of the incidences of pregnancy and pregnancy loss in the feral horses of Sable Island. *Journal of Reproduction and Fertility (Supplement 44)*: 479-488.
- MESSIER, F., D. M. DESAULNIERS, A. K. GOFF, R. NAULT, R. PATENAUE, AND M. CRETE. 1990. Caribou pregnancy diagnosis from immunoreactive progestins and estrogens excreted in feces. *The Journal of Wildlife Management* 54: 279-283.
- MONFORT, S. L., C. C. SCHWARTZ, AND S. K. WASSER. 1993. Monitoring reproduction in moose

- using urinary and fecal steroid metabolites. *The Journal of Wildlife Management* 57: 400–407.
- SASSER, R. G., C. A. RUDER, K. A. IVANI, J. E. BUTLER, AND W. C. HAMILTON. 1986. Detection of pregnancy by radioimmunoassay of a novel pregnancy-specific protein in serum of cows and a profile of serum concentrations during gestation. *Biology of Reproduction* 35: 936–942.
- SCHWARTZ, C. C., S. L. MONFORT, P. DENNIS, AND K. J. HUNDERTMARK. 1995. Fecal progesterone concentration as an indicator of the estrous cycle and pregnancy in moose. *The Journal of Wildlife Management* 59: 583–590.
- SCHWARZENBERGER, F., E. MOSTL, R. PALME, AND E. BAMBERG. 1996. Faecal steroid analysis for non-invasive monitoring of reproductive status in farm, wild and zoo animals. *Animal Reproduction Science* 42: 515–526.
- WASSER, S. K., S. L. MONFORT, AND D. E. WILDT. 1991. Rapid extraction of steroids for measuring reproductive cyclicity and early pregnancy in free-ranging yellow baboons (*Papio cynocephalus cynocephalus*). *Journal of Reproduction and Fertility* 92: 415–423.
- , ———, J. SOUTHERS, AND D. E. WILDT. 1994. Excretion rates and metabolites of oestradiol and progesterone in baboon (*Papio cynocephalus cynocephalus*) faeces. *Journal of Reproduction and Fertility* 101: 213–220.
- WHITE, P. J., R. A. GARROTT, J. F. KIRKPATRICK, AND E. V. BERKELEY. 1995. Diagnosing pregnancy in free-ranging elk using fecal steroid metabolites. *Journal of Wildlife Diseases* 31: 514–522.
- WISDOM, M. J., J. G. COOK, M. M. ROWLAND, AND J. H. NOYES, J. H. 1993. Protocols for care and handling of deer and elk at the Starkey Experimental Forest and Range. United States Forest Service, General Technical Report PNW-GTR-311, U.S. Forest Service, Portland, Oregon, USA, 49 pp.

*Received for publication 1 February 1997.*