

Use of Western Immunoblot Analysis for Testing Moose Serum for *Brucella suis* biovar 4 Specific Antibodies

Authors: Edmonds, Matthew D., Ward, Felicia M., O'Hara, Todd M., and Elzer, Philip H.

Source: Journal of Wildlife Diseases, 35(3) : 591-595

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-35.3.591>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Use of Western Immunoblot Analysis for Testing Moose Serum for *Brucella suis* biovar 4 Specific Antibodies

Matthew D. Edmonds,¹ Felicia M. Ward,¹ Todd M. O'Hara,² Philip H. Elzer,^{1,3,4} ¹ Department of Veterinary Microbiology and Parasitology, Louisiana State University School of Veterinary Medicine, Baton Rouge, Louisiana 70803, USA; ² Department of Wildlife Management, North Slope Borough, Barrow, Alaska 99723, USA; ³ Department of Veterinary Science, Louisiana State University Agricultural Center, Baton Rouge, Louisiana 70803, USA; and ⁴ Corresponding author (e-mail: pelzer@unix1.sncc.lsu.edu).

ABSTRACT: To determine if 12 moose (*Alces alces*) from northern Alaska with agglutinating antibodies specific for *Brucella* spp. had been exposed to either *B. suis* biovar 4 or *B. abortus* biovar 1, western immunoblot serologic analysis was performed. Differential serologic responses to strain specific A and M antigenic variances of the lipopolysaccharide O-polysaccharide sugar allowed strain identification. Prior to examination, test sera were absorbed with killed whole cells from either *B. abortus* biovar 1, containing predominately A antigen (A⁺ M⁻); *B. melitensis* biovar 1, containing essentially M antigen (A⁻ M⁺); or *B. suis* biovar 4, containing both antigenic types (A⁺ M⁺). The resulting sera were then examined by western immunoblot for recognition of either *B. abortus* biovar 1, *B. melitensis* biovar 1, or *B. suis* biovar 4 cell lysates. The results of this study indicate that these moose were exposed to *B. suis* biovar 4, a known pathogen of caribou (*Rangifer tarandus*) from arctic Alaska.

Key words: *Alces alces*, *Brucella suis* biovar 4, moose, serology, survey, western immunoblot.

Brucellosis is a serious, debilitating disease characterized by a variety of signs, including abortion, infertility, arthritis, osteomyelitis and mortality in domestic and wildlife species (Enright, 1990). Brucellosis is caused by the gram negative facultative, intracellular pathogens of the genus *Brucella*, including *Brucella abortus*, *B. melitensis*, *B. suis*, *B. canis*, *B. neotomae*, and *B. ovis*. Both bison (*Bison bison*) and elk (*Cervus elaphus*) in Yellowstone and Grand Teton National Parks (Wyoming, USA) are known to be infected with *B. abortus* biovar 1 (Davis, 1990). Free-ranging caribou (*Rangifer tarandus*) in Alaska are infected with *B. suis* biovar 4 (Davis, 1990). Moose (*Alces alces*) share the same habitat with these infected species and possibly come into contact with infected,

aborted fetuses and other tissues; however, few reports exist of moose infected with either *B. abortus* or *B. suis*.

The few documented cases of free-ranging moose with *B. abortus* infections describe severely debilitated animals exhibiting symptoms including fibrinous myocarditis, pericarditis, and peritonitis (Davis, 1990). Honour and Hickling (1993) reported a weak moose from the Mackenzie River (Northwest Territories, Canada) naturally infected with *B. suis* biovar 4, showing large fluctuant carpal masses with bilateral bursitis and osteomyelitis. A survey by Zarnke (1983) found one of 39 moose in Alaska had a weak serologic reaction for *Brucella* sp. specific antibodies. However, studies in British Columbia, Alberta, and Quebec (Canada) of a combined 356 moose herd failed to find any animals with significant levels of *Brucella* sp. specific antibodies (Hudson et al., 1980; Bourque and Higgins, 1984; Zarnke and Yuill, 1981). Experimental inoculation of moose with *B. suis* biovar 4 or *B. abortus* biovar 1 has been shown to produce a strong antibody response and severe disease in some animals (Dieterich et al., 1991; Forbes et al., 1996). Due to the severity of symptoms and limited number of documented moose with brucellosis, it has been postulated that dissemination throughout the herd may be self-limited (Zarnke, 1983; Forbes et al., 1996). However, six northern free-ranging moose cows in Alaska with strong serologic reactions for *Brucella* sp. specific antibodies have survived in the field for more than 1 yr (O'Hara et al., 1998). Also, two of these moose raised calves that were still alive 1 yr following parturition.

A key component used to differentiate the different *Brucella* spp. is the structure of the O-polysaccharide chain found on the lipopolysaccharide (LPS). The O-polysaccharide of *B. abortus* biovar 1, causative agent of brucellosis in bison, elk, and cattle in the continental United States, consist primarily of a linear homopolymer of α 1,2-linked 4,6-dideoxy-4-formamido-D-manno-pyranosyl residues, commonly referred to as the A antigen or A⁺ M⁻ (Meikle et al., 1989). In contrast, *B. melitensis* biovar 1, a serious cause of human and caprine brucellosis throughout the Middle East, contains O-polysaccharide made up of a predominately linear pentasaccharide repeating unit composed of four α 1,2-linked and one α 1,3-linked sugars, commonly referred to as the M antigen or A⁻ M⁺. The A and M antigens elicit antibodies of different specificity, allowing differentiation based on serology. Unlike *B. abortus* biovar 1 type strain 2308 (S2308) or *B. melitensis* biovar 1 type strain 16M (S16M), *B. suis* biovar 4 elicits the production of both A and M antigen specific antibodies (A⁺ M⁺) from the host. This is because the O-polysaccharide of *B. suis* biovar 4 has a 1:7 ratio of α 1,3-linked to α 1,2-linked sugars as compared to the 1:4 ratio of S16M. Therefore, animals infected with *B. suis* biovar 4 should elicit the production of both A and M antigen specific antibodies unlike S2308 (A⁺ M⁻) or S16M (A⁻ M⁺) infected animals (Meikle et al., 1989).

Here we report the identification of both A and M antigen specific antibodies by western immunoblot in serum collected from free-ranging northern Alaskan moose with a higher prevalence of antibody to *Brucella* spp. than expected (O'Hara et al., 1998). Data from this study suggests that these moose from northern Alaska were exposed to *B. suis* biovar 4.

Sera were obtained from moose from the Colville River and associated drainages (near Umiat, North Slope Borough, Alaska; 69°–70°N, 153°–151°W; O'Hara et al., 1998). All samples were obtained by venous jugular puncture on anesthetized an-

imals as described previously (O'Hara et al., 1998). A combined total of 54 moose serum samples were obtained in two studies from 1996 and 1997. Six animals from the 1996 study were positive on the standard card test (Alton et al., 1975) for agglutinating antibodies to *B. abortus* biovar 1. Four of the six 1996 card test positive animals remained positive for agglutinating antibodies to *B. abortus* biovar 1 in 1997. Also, an additional two animals from the 1997 study were found to possess agglutinating antibodies. All 54 samples, including the 12 card test positive samples (representing 8 animals) along with 12 negative and 15 positive controls, were submitted for analysis by western immunoblot. A blind study was done with the results being recorded qualitatively to identify negative and positive animals. The western immunoblot was used only as a research tool and no attempt was made to verify it as an official diagnostic test.

Sera were again analyzed by the card test (Alton et al., 1975) and western immunoblot (Schurig et al., 1991). These two tests were selected because the card test is an accepted field agglutination test, and the western immunoblot is a sensitive direct binding assay able to detect as little as 1 ng of protein (Harlow and Lane, 1988). Western immunoblot analysis was performed using cell lysates from *B. abortus* S2308, *B. melitensis* S16M, and *B. suis* biovar 4. Cell lysates were obtained by sonication and dilution in Laemmli sample buffer (Schurig et al., 1991). Electrophoresis and transfer to nitrocellulose were performed according to Schurig et al. (1991). Briefly, electrophoresis was performed with a BioRad Mini-Protean II unit (Hercules, California, USA) utilizing the Laemmli discontinuous SDS-PAGE method with 15 μ l of cell lysate per 20 μ l well. Nitrocellulose transfer was performed for 1 hr with 150V in a Mini Trans-Blot Electrophoretic Transfer Cell (BioRad). Following transfer the blots were blocked with a 5% nonfat milk buffer for 30 min followed by washing with tris-buff-

ered saline containing 0.5% Tween-20. Three adjoining membrane lanes containing the three cells lysates were then incubated for 1 hr at 37 C with a 1:40 dilution of the test serum. Subsequently, the blots were washed and incubated with rabbit anti-bovine IgG horseradish peroxidase conjugate (Sigma Chemical Company, Saint Louis, Missouri, USA) at a dilution of 1:800 for 30 min at room temperature. After washing, the substrate 4-chloro-1-naphthol (Sigma) was added for color development.

To perform the absorption assay, whole killed cells of S2308, S16M, and *B. suis* biovar 4 were obtained by formalin fixing a fresh culture of each strain. Serum from each animal was diluted 1:9 in tris-buffered saline and divided into four 1 ml aliquots. To three 1 ml samples, 10 mg of whole killed cells from either S2308, S16M, or *B. suis* biovar 4 were added. As a control, no cells were added to the fourth aliquot. The diluted serum was then incubated for 1 hr at 37 C. Following incubation, the samples were centrifuged at 14,000 rpm, and the aqueous unbound serum removed from the cellular debris. Serum was then analyzed by western immunoblot, as described above, at concentrations of 1:40 and 1:400 per sample.

Serum absorbed with each strain of brucellae was compared to the unabsorbed control following a five min development. Typically, *Brucella* spp. O-polysaccharide from the LPS produces a nondiscrete, ladder-like profile from 29 to 68 kDa when run on a SDS-PAGE gel (Allen et. al., 1998). The binding of anti-O-polysaccharide antibodies to this protein ladder during a western immunoblot produces a dark smear from 29 to 68 kDa based on protein markers ran concurrently on the same gel (Allen et. al., 1998). Positive reactions were qualitatively recorded as either faint, medium, or dark. A faint reaction was characterized by a slight color halo along the ladder that was light enough to allow visualization of the many different protein bands present from 29 to 68 kDa. A me-

dium reaction was dark enough to prevent the visualization of some protein bands, and a dark reaction caused the complete coverage of every protein band along the smear. As positive controls, sera from six cattle experimentally infected with *B. abortus* S2308, six goats infected with *B. melitensis* S16M, and three pigs infected with *B. suis* biovar 4 also were tested along with 12 known negative controls.

Western immunoblot analysis found the 12 card test positive sera samples (representing 8 individual animals) to have antibodies specific for both A and M antigen O-polysaccharide. Furthermore, an additional animal that was card test negative for agglutinating antibodies against *B. abortus* was found to have antibodies against both A and M antigens by western immunoblot. All 13 western immunoblot positive moose samples (representing 9 individuals) with antibodies for *Brucella* sp. were from the Colville River area as described by O'Hara et al. (1998).

The sera from the 13 western immunoblot positive samples, 12 negative controls, and three experimentally infected pigs were then analyzed by absorption with whole killed cells from S2308, S16M, and *B. suis* biovar 4 prior to western immunoblot analysis. Absorption of the sera with S2308 whole killed cells, containing a majority of A antigen O-polysaccharide, caused a decrease in only two of the 13 test samples. However, sera absorption with S16M containing predominately M antigen O-polysaccharide caused an appreciable decrease in western immunoblot reactivity to both *B. suis* biovar 4 and S16M cell lysates. Absorption with *B. suis* biovar 4 led to a decrease in reaction intensity to cell lysates from both predominately A⁺ 2308 and M⁺ S16M indicating the presence of antibodies against both antigens in the sera. The results of two representative serum samples are shown in Table 1. These results were unaffected whether the serum was diluted 1:40 or 1:400. Similar results were obtained from

TABLE 1. Western immunoblot analysis of two Alaskan moose serum samples using cell lysates from *B. abortus* biovar 1 type strain S2308 (S2308), *B. melitensis* biovar 1 type strain 16M (S16M), and *B. suis* biovar 4.

Animal number	No absorption			Absorbed with S2308			Absorbed with S16M			Absorbed with <i>B. suis</i> biovar 4		
	A ^a	B	C	A	B	C	A	B	C	A	B	C
M20 ^b	3 ^c	1	2	1	2	3	2	0	0	2	0	0
M703 ^b	3	1	2	3	2	2	2	1	1	2	1	1
+control ^d	3	1	2	1	2	3	2	0	0	2	0	0
-control ^e	0	0	0	0	0	0	0	0	0	0	0	0

^a A = S2308 cell lysate, B = S16M cell lysate, C = *B. suis* biovar 4 cell lysate.

^b Moose card test positive for *Brucella* sp. specific antibodies.

^c 0 = no reaction with cell lysate by western immunoblot, 1 = faint reaction, 2 = medium reaction, 3 = dark reaction (See text for explanation).

^d Pig experimentally infected with *B. suis* biovar 4.

^e Moose card test negative for *Brucella* sp. specific antibodies.

the three pigs experimentally infected with *B. suis* biovar 4 (Table 1).

Meikle et al. (1989) showed that while *B. abortus* biovar 1 contains a predominance of the α 1,2-linked O-polysaccharide, A antigen, it also has a small number of α 1,3-linked O-polysaccharide, M antigen, epitopes at a ratio of 18 A antigens to 2 M antigens. Furthermore, *B. melitensis* biovar 1 contains a small number of A antigens at a ratio of one A antigen per 19 M antigens. These strains have been designated typical A and M strains with each epitope composed of five sugar units. Animals infected with *B. abortus* biovar 1 will make antibodies primarily directed against A antigen, and *B. melitensis* biovar 1 infected animals make antibodies primarily against the M antigen. For *B. suis* biovar 4, which expresses both A and M antigens, epitopes were found to consist of seven continuous sugar units as compared to the typical five (Meikle et al., 1989). Hence, animals with *B. suis* biovar 4 infections produce an abundance of antibodies against both the M and A antigens. However, a dominance of M antigens in *B. suis* biovar 4 explains the fact that absorption of the reactive moose sera with whole killed cells of S16M, but not S2308, caused a decrease in reactivity to *B. suis* biovar 4 cell lysate by western immunoblot. The dominance of the M antigen also may have caused the moose to produce a higher proportion of

M antigen specific antibodies. This may explain why absorption of the test sera with S2308 whole killed cells caused a decrease in only two of the 13 samples tested. However, the data does indicate that the moose were producing antibodies to both the A and M epitopes, characteristic of *B. suis* biovar 4 infections.

Data from this study suggest that these northern Alaskan moose from the Colville River area were exposed to *B. suis* biovar 4, a pathogen known to occur at varying annual rates in Arctic Alaskan caribou herds and other species.

We wish to thank the many individuals involved in obtaining the sera and for providing us with samples. We specifically thank members of the Alaska Department of Fish and Game, including G. Carroll, L. Anne Ayres, and R. Zarnke. We thank J. Bevins and J. Blake for initial serologic testing and consultation. Technical assistance in the field was provided by C. Hanns, C. Willetto, and K. Burek.

LITERATURE CITED

- ALLEN, C. A., L. G. ADAMS, AND T. A. FICHT. 1998. Transposon-derived *Brucella abortus* rough mutants are attenuated and exhibit reduced intracellular survival. *Infection and Immunity* 66: 1008-1016.
- ALTON, G. G., L. M. JONES, AND D. E. PIETZ. 1975. Laboratory techniques in brucellosis. World Health Organization Press, Geneva, Switzerland, pp. 1-163.

- BOURQUE, M., AND R. HIGGINS. 1984. Serologic studies on brucellosis, leptospirosis, and tularemia in moose (*Alces alces*) in Quebec. *Journal of Wildlife Diseases* 20: 95–99.
- DAVIS, D. S. 1990. Brucellosis in wildlife. In *Animal brucellosis*, K. Nielsen and J. R. Duncan (eds.). CRC Press Inc., Boca Raton, Florida, pp. 321–334.
- DIETERICH, R. A., J. K. MORTON, AND R. L. ZARNKE. 1991. Experimental *Brucella suis* biovar 4 infection in a moose. *Journal of Wildlife Diseases* 27: 470–472.
- ENRIGHT, F. M. 1990. The pathogenesis and pathobiology of *Brucella* infection in domestic animals. In *Animal brucellosis*, K. Nielsen and J. R. Duncan (eds.). CRC Press Inc., Boca Raton, Florida, pp. 301–320.
- FORBES, L. B., S. V. TESSARO, AND W. LEES. 1996. Experimental studies on *Brucella abortus* in moose (*Alces alces*). *Journal of Wildlife Diseases* 32: 94–104.
- HARLOW, E., AND D. LANE. 1988. Immunoblotting. Antibodies, a laboratory manual. Cold Spring Harbor Laboratory 12: 421–470.
- HONOUR, S., AND K. M. H. HICKLING. 1993. Naturally occurring *Brucella suis* biovar 4 infection in a moose (*Alces alces*). *Journal of Wildlife Diseases* 29: 596–598.
- HUDSON, M., K. N. CHILD, D. F. HATLER, K. K. FUJINO, AND K. A. HODSON. 1980. Brucellosis in moose (*Alces alces*). A serological survey in an open range cattle area of north central British Columbia recently infected with bovine brucellosis. *Canadian Veterinary Journal* 21: 47–49.
- MEIKLE, P. J., M. B. PERRY, J. W. CHERWONOGRODZKY, AND D. R. BUNDLE. 1989. Fine Structure of A and M antigens from *Brucella* biovars. *Infection and Immunity* 57: 2820–2828.
- O'HARA, T. M., J. DAU, G. CARROLL, J. BEVINS, AND R. L. ZARNKE. 1998. Evidence of exposure to *Brucella suis* biovar 4 in northern Alaska moose. *Alces* 34: 31–40.
- SCHURIG, G. G., R. M. ROOP, T. BAGCHI, S. BOYLE, D. BUHRMAN, AND N. SRIRANGANATHAN. 1991. Biological properties of RB51; A stable rough strain of *Brucella abortus*. *Veterinary Microbiology* 28: 171–188.
- ZARNKE, R. L., AND T. M. YUILL. 1981. Serologic survey for selected microbial agents in mammals from Alberta, 1976. *Journal of Wildlife Diseases* 17: 453–461.
- . 1983. Serologic survey for selected microbial pathogens in Alaskan wildlife. *Journal of Wildlife Diseases* 19: 324–329.

Received for publication 3 September 1998.