



Absence of Antibodies Against Canine Distemper Virus in Free-ranging Populations of the Eurasian Badger in Great Britain

Authors: Delahay, Richard, and Frölich, Kai

Source: Journal of Wildlife Diseases, 36(3) : 576-579

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-36.3.576>

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Absence of Antibodies Against Canine Distemper Virus in Free-ranging Populations of the Eurasian Badger in Great Britain

Richard Delahay^{1,3} and Kai Frölich² ¹ Central Science Laboratory, Sand Hutton, York, YO4 1LZ, United Kingdom; ² Institut für Zoo und Wildtierforschung, Alfred Kowalke Str. 17, D-10315, Berlin, Germany; ³ Corresponding author (e-mail: r.delahay@csl.gov.uk).

ABSTRACT: Canine distemper virus (CDV) is a serious disease of wild carnivores throughout the world. In Europe, infection has been detected in several carnivores including the Eurasian badger (*Meles meles*). In the present study 182 badger blood samples were collected from an intensively studied population of wild badgers in southwestern England (January–July, 1997), and a further 286 from throughout southern Britain (June 1996–November 1998). A neutralizing peroxidase-linked antibody test was used for the detection of antibodies against CDV. All the samples were negative for CDV antibodies, suggesting that in contrast to mainland Europe, the disease may be either absent or maintained at low levels in British badgers.

Key words: Canine distemper virus, Eurasian badger, *Meles meles*, sero-survey.

Canine distemper virus (CDV) is classified in the Morbillivirus genus of the family Paramyxoviridae and is the causative agent of a serious infectious disease in domestic and free-living carnivores throughout the world (Appel, 1987). Canine distemper is an acute or subacute highly contagious febrile disease that may be manifested by signs of generalized infection, respiratory disease, hyperkeratosis, disruption of the central nervous system or a combination of these (Budd, 1981; Appel and Summers, 1995).

Canine distemper virus infection has been diagnosed in wild-living members of the Canidae, Mustelidae, Hyaenidae, Procyonidae, Ailuridae, Ailuropodidae, Viverridae, Ursidae, and Felidae (Montali et al., 1987). Epizootics have been recorded in several species including gray foxes (*Urocyon cinereoargenteus*), raccoon dogs (*Nyctereutes procyonoides*), raccoons (*Procyon lotor*), and striped skunks (*Mephitis mephitis*) (Appel, 1987; Machida et al., 1993). Canine distemper virus affects susceptible animals of all ages. Morbidity and

mortality rates vary among species and can be high in susceptible host populations (e.g., Machida et al., 1993; Appel and Summers, 1995). However, other wildlife species show little evidence of an effect at the population level despite widespread exposure to CDV (Creel et al., 1997). In Germany an interaction between wildlife and domestic dogs has been suggested as the prevalence of CDV in martens (*Martes* spp.) was positively correlated with that in domestic dogs (Steinhagen and Nebel, 1985).

In the last two decades several dramatic and unexpected distemper episodes have drawn attention to the importance of CDV in wildlife species (Appel and Summers, 1995). The first confirmed case of CDV in a Eurasian badger (*Meles meles*) was reported by Armstrong and Anthony (1942). Fischer (1965) described lesions of the CNS in four free-ranging badgers from Switzerland, in which CDV was suggested as a possible cause. Canine distemper virus infection has also been detected in free-ranging badgers in Austria (Kolbl et al., 1990) and southwestern Germany (van Moll et al., 1995). In the British Isles badger numbers have increased substantially in recent years (Wilson et al., 1997). Furthermore, the intimate communal behaviour of high density badger populations would facilitate spread of CDV within social groups by inhalation (Appel, 1987), and the cool, dark conditions in their underground burrow systems (setts) might allow the virus to persist outside the host (Shen and Gorham, 1980). However, there is currently no published information on CDV in wild badgers in the British Isles. Therefore, our objective was to determine if free-ranging badgers in Britain were naturally infected with CDV.

has not experienced widespread exposure to CDV. The sample provided broad coverage of age ($n = 67$ cubs, 12 yearlings and 103 adults) and sex classes ($n = 75$ males, 107 females). In this high density population the aggregation of individual animals into territorial social groups could mitigate against the rapid dissemination of infectious disease throughout the population and contribute to the persistent concentration of infection within certain groups (Cheeseman et al., 1988). However, the blood samples obtained in the present study came from badgers resident in 28 contiguous social groups, thus providing broad coverage. Results from the smaller number of samples obtained from elsewhere in southern Britain, provide less compelling evidence for the absence of infection (particularly if it were highly aggregated), but are consistent with either absence or relatively low levels of exposure. Furthermore, the absence of positive results from badgers in Great Britain is in stark contrast to those obtained from the limited sampling of badgers carried out on the European mainland (e.g., Kolbl et al., 1990; van Moll et al., 1995).

In Britain the CDV status of wildlife species is unknown, but in domestic dogs infection is rare owing to a widespread and successful vaccination program (H. Thompson, pers. comm.). Nevertheless, the potential for transmission between unvaccinated domestic dogs and wildlife may exist in areas where interactions are most frequent (e.g., suburbs). Unfortunately, in the present study we could not explicitly test for aggregations of CDV infection in badgers in such areas, as samples were predominantly rural in origin.

We thank the Veterinary Laboratories Agency (Surrey, UK) and W. Pressling in particular, for providing blood samples from badger culling operations. Thanks also to P. Spyvee, D. Handoll, L. Rogers and P. Mallinson for the collection of blood samples at Woodchester Park, and the farmers and landowners in the study area for their cooperation during badger

trapping. We are also grateful to B. Kirsch for technical assistance and to S. Chalmers, B. Rima and H. Thompson for their comments on the manuscript.

LITERATURE CITED

- APPEL, M. J. G. 1987. Canine distemper virus. *In* Virus infections of carnivores, Vol. 1, M. J. G. Appel (ed.). Elsevier Science Publishers, Amsterdam, Netherlands, pp. 133–159.
- , AND D. S. ROBSON. 1973. A microneutralization test for canine distemper virus. *American Journal of Veterinary Research* 34: 1459–1463.
- , AND B. A. SUMMERS. 1995. Pathogenicity of morbilliviruses for terrestrial carnivores. *Veterinary Microbiology* 44: 187–191.
- ARMSTRONG, H. W., AND C. H. ANTHONY. 1942. An epizootic of canine distemper in a zoological park. *Cornell Veterinarian* 32: 286–288.
- BUDD, J. 1981. Distemper. *In* Infectious diseases of wild mammals, J. W. Davis, L. H. Karstad and D. O. Trainer (eds.). The Iowa State University Press, Ames, Iowa, pp. 31–44.
- CHEESEMAN, C. L., AND P. J. MALLINSON. 1979. Radio tracking in the study of bovine tuberculosis in badgers. *In* A Handbook on biotelemetry and radio tracking, C. J. Amlaner Jr. and D. W. Macdonald (eds.). Pergamon Press, New York, pp. 649–656.
- , AND J. W. WILESMITH, F. A. STUART, AND P. J. MALLINSON. 1988. Dynamics of tuberculosis in a naturally infected badger population. *Mammal Review* 18: 61–72.
- CREEL, S., N. M. CREEL, L. MUNSON, D. SANDERLIN, AND M. J. G. APPEL. 1997. Serosurvey for selected viral diseases and demography of African wild dogs in Tanzania. *Journal of Wildlife Diseases* 33: 823–832.
- FISCHER, K. 1965. Staupe-Enzephalitis bei Dachsen. *Schweizer Archiv für Tierheilkunde* 107: 87–91.
- GRAHAM, R. C., U. LUNDHOLM, AND M. J. KARNOVSKY. 1965. Cytochemical demonstration of peroxidase activity with 3-amino-9-ethyl carbazole. *Journal of Histochemistry and Cytochemistry* 13: 150–152.
- HAIG, D. A. 1948. Preliminary note on the cultivation of green's distemperoid virus in fertile hen eggs. *Onderstepoort Journal of Veterinary Science and Animal Industry* 23: 149–155.
- KOLBL, S., H. SCHNABEL, AND M. MIKULA. 1990. Staupeinfektion als Todesursache bei Dachsen in Österreich. *Tierärztliche Praxis* 13: 81–84.
- MACHIDA, N., K. KIRYU, K. OH-ISHI, E. KANDA, N. IZUMISAWA, AND T. NAKAMURA. 1993. Pathology and epidemiology of canine distemper in raccoon dogs (*Nyctereutes procyonoides*). *Journal of Comparative Pathology* 108: 383–392.
- MACKINTOSH, C. G., J. A. MACARTHUR, T. W. A. LITTLE, AND P. STUART. 1976. The immobilization

- of the badger (*Meles meles*). British Veterinary Record 132: 609–614.
- MONTALI, R. J., C. R. BARTZ, AND M. BUSH. 1987. Canine distemper virus. In *Virus infections of carnivores*, M. Appel (ed.). Elsevier Science Publisher B. V., Amsterdam, Netherlands, pp. 437–443.
- SHEN, D. T., AND J. R. GORHAM. 1980. Survival of pathogenic virus at 5° C and 25° C. *Veterinary Medicine/Small Animal Clinician* 75: 69–70.
- STEINHAGEN, P., AND W. NEBEL. 1985. Staupe beim Steinmarder (*Martes foina*) in Schleswig-Holstein—Beitrag zur Epidemiologie der Staupe. *Deutsch Tierärztliche Wochenschrift* 92: 178–181.
- VAN MOLL, P., S. ALLDINGER, W. BAUMGARTNER, AND M. ADAMI. 1995. Distemper in wild carnivores: An epidemiological, histological and immunocytochemical study. *Veterinary Microbiology* 44: 193–199.
- WILSON, G., S. HARRIS, AND G. MCLAREN. 1997. Changes in the British badger population, 1988 to 1997. Peoples Trust for Endangered Species, London, United Kingdom, pp. 7–9.
- ZAGHAWA, A., B. LIESS, AND H. R. FREY. 1990. Antiserum raised in pigs against canine distemper virus and its utility in diagnostic procedures for Morbillivirus infections (canine distemper, phocine distemper, rinderpest). *Journal of Veterinary Medicine* 37: 353–362.

Received for publication 15 July 1999.