

Abbreviata terrapenis (Nematoda: Physalopteridae): An Accidental Parasite of the Banded Rock Rattlesnake (Crotalus lepidus klauberi)

Authors: Goldberg, Stephen R., Bursey, Charles R., and Holycross, Andrew T.

Source: Journal of Wildlife Diseases, 38(2) : 453-456

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-38.2.453>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Abbreviata terrapenis* (Nematoda: Physalopteridae): An Accidental Parasite of the Banded Rock Rattlesnake (*Crotalus lepidus klauberi*)**

Stephen R. Goldberg,^{1,4} Charles R. Bursey,² and Andrew T. Holycross³ ¹ Department of Biology, Whittier College, Whittier, California 90608, USA; ² Department of Biology, Pennsylvania State University, Shenango Campus, Sharon, Pennsylvania 16146, USA; ³ Department of Biology, Arizona State University, Tempe, Arizona 85287, USA; ⁴ Corresponding author (email: sgoldberg@whittier.edu).

ABSTRACT: The nematode, *Abbreviata terrapenis* (Physalopteridae) was found in 16 (6%) of 267 banded rock rattlesnakes (*Crotalus lepidus klauberi*) from Arizona and New Mexico. *Abbreviata terrapenis* in *C. lepidus* represents an accidental parasite in that “infection” was acquired by the ingestion of lizard prey. Feeding captive snakes on wild-caught lizards poses a risk of introducing nematodes to the snakes.

Key words: *Abbreviata terrapenis*, accidental parasite, banded rock rattlesnake, *Crotalus lepidus klauberi*, nematode.

During a study of diet, nematodes fitting the description of *Abbreviata terrapenis* were collected from the digestive tract of the banded rock rattlesnake (*Crotalus lepidus klauberi*) from Arizona and New Mexico. *Abbreviata terrapenis*, originally described from the ornate box tortoise (*Terrapene ornata*), infects painted turtle (*Chrysemys picta*), snapping turtle (*Chelydra serpentina*), Blanding’s turtle (*Emydoidea blandingii*), Sonoran spotted whiptail (*Cnemidophorus sonora*), western whiptail (*Cnemidophorus tigris*), and mountain spiny lizard (*Sceloporus jarrovi*) (Hill, 1941; Morgan, 1945; Goldberg et al., 1995, 1996, 1997). Because *A. terrapenis* is a heteroxenous helminth with an indirect life cycle involving an insect intermediate host (Anderson, 2000), its occurrence in insectivores, but not carnivores, might be expected.

The rock rattlesnake (*Crotalus lepidus*), a mountain rock dweller found on rocky ridges, hillsides, streambeds, and gorges in arid and semiarid habitats from 300–2,930 m, occurs from southeastern Arizona, southern New Mexico, and west Texas to Jalisco, Mexico (Stebbins, 1985). Four subspecies are recognized: both banded rock rattlesnake (*C. lepidus klauberi*) and mottled rock rattlesnake (*C. lepidus lepi-*

lus) are found in the United States and Mexico; Durangan rock rattlesnake (*C. lepidus maculosus*) and Tamaulipan rock rattlesnake (*C. lepidus morulus*) are restricted to Mexico (Liner, 1994; Crother et al., 2000). Analysis of a large sample suggests that banded rock rattlesnake diet consists primarily of lizards and centipedes but small rodents, birds, and snakes are also consumed (Holycross, unpubl. data). Although McCrystal et al. (1996) reported captive neonatal snakes to eat crickets and Conant (1955) reported a grasshopper and caterpillars along with a lizard in the stomach of a wild-caught mottled rock rattlesnake, insects are not typical rattlesnake prey. Thus the occurrence of *A. terrapenis* in banded rock rattlesnakes from Arizona and New Mexico.

Two hundred sixty-seven banded rock rattlesnakes were borrowed from museums; 16 (6%) were found to harbor *A. terrapenis*. The body size (mean snout-vent length) of infected snakes was 474 ± 84 mm SD (range = 318–667 mm). These snakes were from the following museum collections: California Academy of Sciences (CAS 01814, 48022, 48024–48026, 48029, 48030, 48032, 84131); Museum of Southwestern Biology (MSB 4115, 32319, 49551); San Diego Society of Natural History (SDSNH 2160, 3123); University of Kansas (KU 6648); and University of Michigan (UMMZ 54019). Of the infected snakes, 14 were from Cochise County, Arizona and two were from Hidalgo County, New Mexico. The other snakes were from Arizona, New Mexico, and Chihuahua and Sonora, Mexico. The body cavity was opened by an incision along the ven-

TABLE 1. Number and location of *Abbreviata terrapenis* and prey found in banded rock rattlesnakes.

Snake identification	<i>Abbreviata terrapenis</i>	Helminth site	Prey	Prey site
CAS 01814	3 female	stomach	<i>Sceloporus jarrovii</i>	stomach
CAS 48022	1 female	stomach	<i>Sceloporus</i> sp.	intestine
CAS 48024	4 (2 male, 2 female)	stomach and intestine	none	—
CAS 48025	1 male	stomach	<i>Sceloporus jarrovii</i>	intestine
CAS 48026	6 (3 male, 3 female)	stomach	<i>Sceloporus jarrovii</i>	intestine
CAS 48029	2 female	stomach	none	—
CAS 48030	4 female	stomach	<i>Sceloporus jarrovii</i>	stomach
CAS 48032	10 (3 male, 7 female)	stomach	none	—
CAS 48131	1 male	stomach	<i>Sceloporus jarrovii</i>	stomach
KU 6648	1 male	colon	<i>Sceloporus</i> sp.	intestine
MSB 4115	39 (15 male, 24 female)	stomach	<i>Sceloporus</i> sp.	intestine
MSB 32319	4 3rd stage encapsulated	peritoneum	none	—
MSB 49551	24 (10 male, 14 female)	stomach	none	—
SDSNH 2160	4 4th stage larvae	stomach	none	—
SDSNH 3123	4 female	stomach	none	—
UMMZ 54019	2 (1 male, 1 female)	stomach	<i>Sceloporus jarrovii</i>	stomach
7F7B10752C	2 female	feces	<i>Cnemidophorus</i> sp.	feces
7F7B177600	1 female	feces	<i>Cnemidophorus</i> sp.	feces
7F7B180432	1 female	feces	<i>Sceloporus jarrovii</i>	feces

tral surface and the contents of the stomach and distal 3 cm of the intestine were examined. All prey remains and parasites were removed. In addition, three fecal samples palpated from live banded rock rattlesnakes (catch and release, Peloncillo Mountains [31°31'N, 109°03'W] Hidalgo County, New Mexico, A.T.H. field numbers 7F7B10752C, 7F7B177600, 7F7B180432) contained *A. terrapenis* (Table 1). Prey and parasites were initially identified using a dissecting microscope. Nematodes were cleared in a drop of glycerol and identified using a compound microscope.

There was evidence of lizard consumption in 12 (5%) banded rock rattlesnakes (*Cnemidophorus* in two snakes; *Sceloporus* in 10 snakes) (Table 1). With some variation, the morphology of mature specimens from banded rock rattlesnakes agreed with the description of *A. terrapenis* as presented by Hill (1941). The nematodes possessed symmetrical lips, each supporting a single median external tooth, a single median internal tooth, and two doubled teeth located toward the angles of the mouth. Total length of females ranged from 27–38 mm. The vulva was situated 6–8 mm

from the anterior end. The vagina was long, slender, and ran anteriad for a short distance, then posteriad to an enlargement which narrowed before dividing dichotomously into four uteri. Males were shorter and ranged from 20–26 mm in length. Spicules were very unequal in size and shape with the left spicule about eight times the length of the right. There were four pairs of para-anal pedunculated papillae, a median sessile papilla, one pair of preanal papillae, and five pairs of post anal sessile papillae. Selected nematodes were deposited in the United States National Parasite Collection, (USNPC), Beltsville, Maryland as USNPC 91254.

Babero and Emmerson (1974) reported *Thubunaea cnemidophorus*, normally a parasite of teiid lizards (Goldberg et al., 1997), from three rattlesnake species collected in Nevada: sidewinder (*Crotalus cerastes*), speckled rattlesnake (*C. mitchellii*), and Mojave rattlesnake (*C. scutulatus*). Sidewinders, speckled rattlesnakes, and young Mojave rattlesnakes also feed on lizards (Lowe et al., 1986). *Abbreviata terrapenis* has been previously found in both lizard genera, *Cnemidophorus* and *Scelo-*

porus (Goldberg et al., 1995, 1996, 1997) on which banded rock rattlesnakes feed. Thus, the presence of *A. terrapenis* in banded rock rattlesnakes would appear to be the fourth example of accidental parasitism in North American rattlesnakes. Babero and Emmerson (1974) reported the sidewinder, the speckled rattlesnake, and the Mojave rattlesnake as new host records, a status we believe must be slightly modified.

We have chosen the term "accidental" for this occurrence because of the textbook definitions of parasitism, the term most accurately describes our observations (see Roberts and Janovy, 2000): *A. terrapenis* has entered a host different from its normal, definitive host. We base this supposition on the fact that in snake MSB 32319, four third stage larvae were found encapsulated in the peritoneum. Jones (1995) has reported encapsulated larvae of species of *Abbreviata* in gastric tissues of small snakes and lizards preyed upon by large Australian lizards which in turn serve as definitive hosts of *Abbreviata*. Because third stage larvae of *Abbreviata* had become encapsulated in a rattlesnake, we believe the snake would not be a definitive host, but rather, a potential paratenic host. Fourth stage and mature nematodes were found in the lumen of the digestive tract in a posture similar to that seen in lizard stomachs. How long gravid female nematodes might survive and whether growth and maturity of immature nematodes would occur remains an open question. Because mature nematodes enter with lizard prey, we consider rattlesnakes to be accidental hosts. We would apply the same terminology to the nematodes in the report of Babero and Emmerson (1974).

These observations suggest that feeding captive snakes wild-caught lizards presents an opportunity for transmission of nematodes to the snakes. In addition, when reporting the presence of nematodes in snakes, one must consider the possibility that the nematode entered in a prey item.

We thank J. Vindum (California Acade-

my of Sciences, San Francisco), C. W. Painter (Museum of Southwestern Biology, Albuquerque, New Mexico), B. D. Hollingsworth (San Diego Society of Natural History, San Diego, California), J. Simmons (University of Kansas, Lawrence), and G. Schneider (University of Michigan, Ann Arbor) for permission to examine banded rock rattlesnakes. The Wallace Research Foundation (Tucson, Arizona) provided funding support to A.T.H. Banded rock rattlesnakes were collected under permits issued by Arizona Game and Fish Department (SP841338) and the New Mexico Department of Fish and Game (2824). The study was approved by the Arizona State University Institutional Animal Care and Use Committee (ASU 93-280R).

LITERATURE CITED

- ANDERSON, R. C. 2000. Nematode parasites of vertebrates: Their development and transmission, 2nd Edition. CABI Publishing, CAB International, Wallingford, Oxon, U.K., 650 pp.
- BABERO, B. B., AND F. H. EMMERSON. 1974. *Thubunaea cnemidophorus* in Nevada rattlesnakes. *Journal of Parasitology* 60: 595.
- CONANT, R. 1955. Notes on three Texas reptiles, including an addition to the fauna of the state. *American Museum Novitates* (1726): 1-6.
- CROTHER, B. I., J. BOUNDY, J. A. CAMPBELL, K. DE QUEIROZ, D. R. FROST, R. HIGHTON, J. B. IVERSON, P. A. MEYLAN, T. W. REEDER, M. E. SEIDEL, J. W. SITES, JR., T. W. TAGGART, S. G. TILLEY, AND D. B. WAKE. 2000. Scientific and standard English names of amphibians and reptiles of North America north of Mexico, with comments regarding confidence in our understanding. Society for the Study of Amphibians and Reptiles, Herpetological Circular No. 29, Shoreview, Minnesota, 82 pp.
- GOLDBERG, S. R., C. R. BURSEY, AND R. L. BEZY. 1995. Helminths of isolated montane populations of Yarrow's spiny lizard, *Sceloporus jarrovi* (Phrynosomatidae). *Southwestern Naturalist* 40: 330-333.
- , ———, AND ———. 1996. Gastrointestinal helminths of Yarrow's spiny lizard, *Sceloporus jarrovi* (Phrynosomatidae) in Mexico. *American Midland Naturalist* 135: 299-309.
- , ———, AND H. CHEAM. 1997. Helminths from the Sonoran spotted whiptail, *Cnemidophorus sonorae*, and the western whiptail, *Cnemidophorus tigris* (Sauria: Teiidae), from southern Arizona with comments on *Abbreviata terrapenis*

- (Nematoda: Physalopteridae). Great Basin Naturalist 57: 273–277.
- HILL, W. C. 1941. *Physaloptera terrapenis*, a new nematode from a tortoise. Transactions of the American Microscopical Society 60: 59–64.
- JONES, H. J. 1995. Pathology associated with physalopterid larvae (Nematoda: Spirurida) in the gastric tissues of Australian reptiles. Journal of Wildlife Diseases 31: 299–306.
- LINER, E. A. 1994. Scientific and common names for the amphibians and reptiles of Mexico in English and Spanish. Society for the Study of Amphibians and Reptiles, Herpetological Circular No. 23, Lawrence, Kansas, 113 pp.
- LOWE, C. H., C. R. SCHWALBE, AND T. B. JOHNSON. 1986. The venomous reptiles of Arizona. Arizona Game and Fish Department, Phoenix, Arizona, 115 pp.
- MCCRISTAL, H. K., C. R. SCHWALBE, AND D. F. RETES. 1996. Selected aspects of the ecology of the Arizona ridge-nosed rattlesnake (*Crotalus willardi willardi*) and the banded rock rattlesnake (*Crotalus lepidus klauberi*) in Arizona. Unpublished Report, Arizona Game and Fish Department, Phoenix, 34 pp.
- MORGAN, B. B. 1945. The nematode genus *Abbreviata* (Travassos, 1920) Schulz, 1927. American Midland Naturalist 34: 485–490.
- ROBERTS, L. S., AND J. JANOVY, JR. 2000. Gerald D. Schmidt & Larry S. Roberts' Foundations of Parasitology, 6th Edition. McGraw Hill, Boston, Massachusetts, 670 pp.
- STEBBINS, R. C. 1985. A field guide to western reptiles and amphibians. Houghton Mifflin Company, Boston, Massachusetts, 336 pp.

Received for publication 25 April 2001.