

## CORRECTION

Source: Journal of Wildlife Diseases, 39(1) : 246

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-39.1.246>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## CORRECTION

Southey et al., 2002. Sulfadimethoxine and rhodamine B as oral biomarkers for European badgers.

On page 380 of our manuscript we wrote:

“All of the badgers trapped over the course of the study were anesthetized with 0.1 mg/kg ketamine hydrochloride (Vetalar, Pharmacia and Upjohn Ltd., Crawley, UK) and 0.1 mg/kg medetomidine hydrochloride (Domitor, Orion Corporation, Espoo, Finland), administered by intramuscular injection.”

We wish to correct this by stating that the dose of ketamine hydrochloride used to anesthetize badgers should have read as 10 mg/kg.