



INFECTIOUS DISEASE SURVEY OF GEMSBOK IN NEW MEXICO

Authors: Bender, Louis C., Li, Hong, Thompson, Bruce C., Morrow, Patrick C., and Valdez, Raul

Source: Journal of Wildlife Diseases, 39(4) : 772-778

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-39.4.772>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

INFECTIOUS DISEASE SURVEY OF GEMSBOK IN NEW MEXICO

Louis C. Bender,^{1,5} Hong Li,² Bruce C. Thompson,¹ Patrick C. Morrow,³ and Raul Valdez⁴

¹U.S. Geological Survey, New Mexico Cooperative Fish and Wildlife Research Unit, P.O. Box 30003 MSC 4901, Las Cruces, New Mexico 88003, USA

²Animal Diseases Research Unit, USDA/ARS, Washington State University, 3003ADBF, Pullman, Washington 99164, USA

³U.S. Army White Sands Missile Range, Environmental Stewardship Division, CSTE-DTC-ES, WSMR, New Mexico 88002, USA

⁴Department of Fishery and Wildlife Sciences, New Mexico State University, Las Cruces, New Mexico 88003, USA

⁵Corresponding author (email: lbender@nmsu.edu)

ABSTRACT: Exotic wildlife can introduce new diseases or act as reservoirs of endemic diseases. On White Sands Missile Range, New Mexico (USA), significant declines in populations of native ungulates generally correspond to increases in range and population density of the exotic gemsbok (*Oryx gazella gazella*), introduced beginning in 1969. We surveyed gemsbok in 2001 for exposure to a variety of diseases potentially important for native ungulates. High seroprevalence was found for malignant catarrhal fever virus (49 [98%] of 50 sera; 43 [96%] of 45 plasma samples), blue-tongue virus (48 [96%] of 50), bovine respiratory syncytial virus (33 [66%] of 50), and parainfluenza-3 virus (10 [20%] of 50). Low numbers of *Nematodirus* spp. eggs in a few individuals were the only parasites detected in gemsbok. Exposure to the above diseases in gemsbok is of interest to managers because of potential implications for recovery of desert bighorn sheep (*Ovis canadensis mexicana*) and desert mule deer (*Odocoileus hemionus crooki*) in the White Sands area because each has been implicated in mortality in these species either in the White Sands area or elsewhere in the western/southwestern United States.

Key words: Bluetongue, bovine respiratory syncytial virus, desert bighorn sheep, gemsbok, malignant catarrhal fever, mule deer, New Mexico, parainfluenza-3, pasteurellosis.

INTRODUCTION

Gemsbok (*Oryx gazella gazella*) were introduced onto White Sands Missile Range (WSMR, New Mexico, USA) by the New Mexico Department of Game and Fish (NMDGF) to provide a huntable ungulate in an area of the state providing limited big game hunting opportunities (Wood et al., 1970). Native to arid habitats throughout southern Africa, 93 gemsbok obtained from captive-bred stock were released from 1969–77 and have since greatly increased their range and population density throughout WSMR and surrounding areas (Burkett et al., 2002). Gemsbok population estimates currently range from 3,000–6,000 (L. Bender, unpubl. data), and are still increasing despite hunting harvests in excess of 600–700 annually (Burkett et al., 2002). The current extent of gemsbok range is unknown, but is in excess of 15,000 km² in southern New Mexico. The significant increases in range, population size, and population density illustrate how well gemsbok have adapted

to habitat conditions of the Chihuahuan Desert, which are markedly similar to their native range in southern Africa (Wood et al., 1970). In the ~30 yr since their introduction, gemsbok have become the most numerous ungulate in the Chihuahuan Desert system of New Mexico (Burkett et al., 2002).

Concurrent with expansion of gemsbok was a decline in native ungulates, particularly desert mule deer (*Odocoileus hemionus crooki*) and the state-designated endangered desert bighorn sheep (*Ovis canadensis mexicana*) (Burkett et al., 2002). Neither proximate nor ultimate causes of decline of these species have been identified, although speculation for desert bighorn has focused on disease, particularly scabies, despite the absence of documented mortalities (Lange et al., 1980; Hoban, 1990). Anecdotal evidence suggests that dramatic population declines coincided with large increases in gemsbok population densities and expansion of gemsbok range. Gemsbok possess a number of characteristics that make them superior com-

petitors to mule deer and desert bighorn in the arid relatively unproductive habitats of the Chihuahuan Desert. Among these are a broader diet breadth (Smith et al., 1998), impressive physiologic and behavioral specializations to conserve water (Taylor, 1969; Hamilton et al., 1977; Burkett et al., 2002), and aggressive behavior (Walther, 1980). Gemsbok are also less vulnerable to native Chihuahuan Desert predators (Burkett et al., 2002).

Gemsbok are an introduced exotic in the Chihuahuan Desert. As well as competing with native species, introduced species have the potential to introduce new diseases, as well as to serve as a new reservoir for existing diseases (Clark et al., 1970; Miller, 2001; Rupprecht et al., 2001). A recent example of the former include introduced reindeer (*Rangifer tarandus tarandus*) that brought the parasite *Elaphostrongylus rangiferi* from Norway to Newfoundland (Canada); this exotic parasite has caused epizootics of cerebrospinal elaphostrongylosis in native caribou (*R. tarandus caribou*) (Ball et al., 2001). Similarly, movement of ranched elk (*Cervus elaphus*) has been implicated in spread of chronic wasting disease (CWD) among game farms (Williams et al., 2001).

Despite the potential of introduced diseases accompanying introduced wildlife, disease surveillance has been sporadic for gemsbok in New Mexico, considering their widespread distribution and large population size. Previous work identified exposure of gemsbok to viral pathogens including bluetongue virus (BTV), epizootic hemorrhagic disease virus (EHDV), bovine viral diarrhea (BVD) virus, parainfluenza-3 (PI3) virus, bovine respiratory syncytial virus (BRSV), equine viral rhinopneumonitis virus, western equine encephalitis virus (WEEV), vesicular stomatitis virus, infectious bovine rhinotracheitis (IBR) virus, and Malpai Springs virus (MSV) (Clark et al., 1988; White Sands Missile Range, 1991; Calisher and Taylor, 1993). Calisher and Taylor (1993) speculated on the role of gemsbok as a reservoir or an amplifying

host of viral diseases or viruses (BTV, EHDV, WEEV, MSV) on WSMR based upon long-term high seroprevalence. However, most of this preliminary work focused on longitudinal studies of select diseases, not potential implications to wild ungulate populations, because gemsbok numbers and distribution were still relatively limited during these early studies (600–800; Clark et al., 1988) and population declines of native ungulates were not yet widespread.

Since the initial disease work in the 1980s, gemsbok numbers and distribution have increased dramatically, and populations of native ungulates have declined significantly and show no signs of recovery. Thus, our goal was to survey free-ranging gemsbok for exposure to selected diseases and parasites, emphasizing those associated with exotic ungulates or previously identified with gemsbok in New Mexico which could pose potential threats to native ungulates (desert mule deer, desert bighorn sheep, pronghorn [*Antilocapra americana*]) in New Mexico.

MATERIALS AND METHODS

White Sands Missile Range (approximately 32°50'N, 106°30'W) encompasses about 8,800 km² (approximately 165 km north to south and 64 km east to west) and includes portions of the Jornada del Muerto, the Tularosa Basin, and the San Andres and several other mountain ranges of southcentral New Mexico. Major vegetation communities on WSMR include semidesert grassland, Chihuahuan desertscrub, and Great Basin conifer woodland (Burkett et al., 2002). Physical, climatic, and vegetation characteristics of WSMR and surrounding areas are detailed in Burkett et al. (2002).

We collected blood and fecal samples from 101 gemsbok captured by aerial darting on WSMR incidental to a large-scale capture operation during February and April 2001. Gemsbok were captured over approximately 700,000 ha and included all sex and age classes except those <6 mo old. We collected blood samples by jugular puncture and fecal samples from the rectum. Blood samples were divided into serum and ethylenediaminetetraacetic acid tubes and placed in a cooler immediately after collection. Serum tubes were spun (3,200 × G; 8–10 min) and separated the evening following

TABLE 1. Results of serology for evidence of exposure to *Leptospira interrogans* (*Leptospira*; serovars bratislava, canicola, grippityphosa, hardjo, icterohemorrhagiae, and pomona), *Brucella abortus*, *Brucella ovis*, *Mycobacterium avium* subsp. *paratuberculosis*, *Anaplasma* spp., bovine viral diarrhea (BVD) virus, infectious bovine rhinotracheitis virus (bovine herpesvirus-1; IBR), bluetongue virus (BRV), bovine respiratory syncytial virus (BRSV), parainfluenza-3 (PI-3) virus, and malignant catarrhal fever (MCF) virus in free-ranging gemsbok from White Sands Missile Range, New Mexico.

| Agent | Number tested/number positive | Test (threshold) for positive reaction) ^a | Reference |
|---|-------------------------------|--|------------------------------|
| <i>Leptospira</i> | 1/50 | MAT (1:100) | Gochenour et al., 1958 |
| <i>Brucella abortus</i> | 0/50 | Card ^b | Alton et al., 1985 |
| <i>Brucella ovis</i> | 1/50 | ELISA ^c | Lee et al., 1985 |
| <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> | 0/50 | ELISA ^d | Shulaw et al., 1986 |
| <i>Anaplasma</i> spp. | 0/50 | CF (1:5) | Hidalgo and Dimopoulos, 1967 |
| BVD virus | 2/50 | VN (1:4) | Carbrey et al., 1971 |
| IBR virus | 0/50 | VN (1:4) | Carbrey et al., 1971 |
| BTB | 48/50 | cCELISA ^e | Pearson et al., 1992 |
| BRSV | 33/50 | VN (1:4) | Westenbrink et al., 1985 |
| PI-3 virus | 10/50 | VN (1:4) | Carbrey et al., 1971 |
| MCF virus (serum) | 49/50 | CI-ELISA ^f | Li et al., 1994, 2001 |
| MCF virus (plasma) | 43/45 | CI-ELISA ^f | Li et al., 1994, 2001 |

^a Tests were: MAT=microscopic agglutination test; CF=complement fixation; ELISA=enzyme-linked immunoabsorbent assay (c=competitive, CI=competitive inhibition); VN=virus neutralization. Titer in parentheses is threshold value for positive test.

^b No agglutination on BAPA card=negative; reactors on BAPA then cross-checked by BALA (latex agglutination).

^c Any optical density greater than the National Veterinary Services Laboratory (Ames, Iowa) standard positive at 1:10 by complement fixation.

^d Ratio of antibody detected in serum sample compared to a high positive standard serum (S/P)>0.25=positive; S/P 0.200–0.249=suspect.

^e >50% inhibition of negative controls=positive.

^f >25% inhibition based on a panel of MCF virus-free sheep sera.

collection. Fecal samples were refrigerated within 30–60 min after collection until analysis.

We randomly selected 50 individual blood samples for disease screening. We selected health tests (Table 1) based on potential significance to gemsbok or other ungulates in the Chihuahuan Desert system. We tested for malignant catarrhal fever (MCF) virus antibodies using the competitive inhibition enzyme-linked immunosorbent assay (Li et al., 1994, 2001) on serum and an additional 45 plasma samples following positive results from serum (not all serum and plasma samples were from the same individuals). Parasites were identified and quantified using either fecal flotation (Foreyt, 2001) or the modified Baermann test for lungworms (Forrester and Lankester, 1997).

RESULTS

Exposure of gemsbok to a variety of disease agents was shown, of which three were of immediate relevance for native ungulates on WSMR (Table 1). First, 98% of serum samples and 96% of plasma sam-

ples from gemsbok were positive for MCF virus antibody. Secondly, 66% of gemsbok were positive for BRSV antibodies (titers 1:4–1:128) and 20% for PI3 virus (titers 1:4–1:128) antibodies. Lastly, 96% of gemsbok were positive for BTB antibody. Small numbers of animals were positive for BVD virus antibodies and antibodies against *Brucella ovis* and *Leptospira interrogans* subspecies bratislava (titer 1:100). Antibodies were not detected for other agents (Table 1). The only parasite found in fecal floatations was *Nematodirus* spp. (2–5 eggs/g) in three (6%) of 50 gemsbok.

DISCUSSION

Malignant catarrhal fever is a sporadic, infectious disease of domestic and wild ruminants (Heuschele and Reid, 2001). It is often fatal in aberrant hosts, with deer in particular being especially vulnerable.

Classic MCF was first attributed to either alcelaphine herpesvirus-1 (AIHV-1) the virus first identified in wildebeest (*Connochaetes taurinus*), or ovine herpesvirus-2 (OvHV-2), the natural host of which is domestic sheep (Heuschele and Reid, 2001). Since identification of these viruses, at least four additional herpesviruses in the MCF group have been recognized (Li et al., 2000, 2001; Heuschele and Reid, 2001). Antibody cross-reactions to AIHV-1 in a variety of other African ungulates suggested that other unidentified MCF-group herpesviruses exist (Plowright, 1981). Although transmission of MCF is not completely understood, transmission of AIHV-1 to other species usually occurs from newborns, who shed virus in tears and nasal secretions until ~3–4 mo old, when they develop adequate immunity and cease shedding virus (Heuschele and Reid, 2001).

Polymerase chain reaction sequencing identified two new herpesviruses from gemsbok on WSMR: one within the MCF virus group that is closely related to AIHV-1 and another within the non-MCF lymphotropic herpesvirus group (Li et al., 2003). Currently, no data are available as to whether these viruses are pathogenic to other species. However, presence of an MCF-group herpesvirus similar to AIHV-1 warrants further investigation due to the possibility that this virus may cause MCF in desert mule deer and other North American ungulates.

Antibodies against two bovine respiratory viruses implicated as predisposing agents to pasteurellosis, BRSV and PI3 virus (Miller, 2001), were found in gemsbok on WSMR. Pneumonic pasteurellosis can cause high mortality in bighorn sheep populations (Miller, 2001). Heavy infestation with lungworms (i.e., *Protostrongylus* spp., *Muellerius* spp.) may also predispose bighorn sheep to pneumonic pasteurellosis (Spraker and Hibler, 1982; Miller, 2001). It has been hypothesized that abundance and distribution of bighorn sheep popula-

tions is limited by recurrent pasteurellosis epidemics (Hobbs and Miller, 1992).

Gemsbok occur throughout desert bighorn range adjacent to WSMR, and because the San Andres range is a priority desert bighorn recovery area in New Mexico, the potential for disease to limit success of future transplant efforts needs to be evaluated. This is especially true because desert bighorn apparently declined from epizootics of unknown origin several times in the past in this area (Hoban, 1990). Gemsbok should be sampled to determine whether BRSV and PI3 virus are being maintained in the population, or whether antibodies detected were due to exposure to bighorn sheep and/or livestock with these viruses. This is particularly relevant given that recent surveillance resulted in isolation of *Pasteurella trehalosi* from gemsbok on San Andres National Wildlife Refuge (SANWR), adjacent to WSMR (Bender and Weisenberger, unpubl. data). Thus, if BRSV and/or PI3 virus are common in gemsbok there could be an impact on successful desert bighorn sheep recovery in areas occupied by gemsbok.

Bluetongue is an arthropod-vector-borne infectious but noncontagious disease of wild and domestic ruminants (Ramsay et al., 1985; Howerth et al., 2001). Among ruminants present in the gemsbok range, bluetongue has caused clinical disease in mule deer, elk, bison (*Bison bison*), pronghorn, and desert bighorn sheep (Howerth et al., 2001). Bluetongue can cause extensive wildlife mortalities over large areas (Thorne, 1982; Howerth et al., 2001), low-level annual mortality (Hoff et al., 1974), and limit reproduction following outbreaks due to fetal resorption, abortion, and/or congenital physical or nervous system deformities without causing clinical disease in adults (Thorne, 1982; Ramsay et al., 1985). The presence of extensive outbreaks of bluetongue in domestic sheep and subsequent bluetongue in desert bighorn sheep was implicated in elimination of desert bighorn from Texas (Robinson et al., 1967). Mortality from bluetongue also

was documented in desert bighorn from SANWR in New Mexico (Lange et al., 1980; Hoban, 1990).

Perinatal infection with BTV has been reported in an aborted gemsbok calf (Ramsay et al., 1985), though no cases of clinical disease in adults have been reported (Howerth et al., 2001). However, based on high levels of exposure to BTV detected during long-term serologic surveillance, gemsbok may serve as amplifying hosts of BTV in the WSMR vicinity (Calisher and Taylor, 1993). If so, presence of BTV in gemsbok has implications for native ungulates in gemsbok range, particularly mule deer and desert bighorn sheep, both of which are susceptible to bluetongue and present in extremely low numbers.

Management implications

Behavioral, physiologic, and physical characteristics make gemsbok a superior competitor to native ungulates in the Chihuahuan Desert system, and thus they potentially limit population size of native ungulates (Burkett et al., 2002). Gemsbok also may have introduced new disease agents (i.e., MCF) and/or serve as a reservoir for existing disease agents to which they are resistant but native ungulates susceptible. Combined impacts of competition and disease may make restoration of viable populations of native species such as desert bighorn sheep difficult or impossible in areas dominated by gemsbok. Further, popular game species such as desert mule deer also may remain depressed.

In New Mexico, gemsbok are highly prized by the hunting public as a trophy of great value; gemsbok hunts on WSMR are once in a lifetime licenses, with >4,000 hunters applying for >800 licenses annually. Any effort to eliminate or limit gemsbok numbers because of perceived detriment to desert bighorn sheep and desert mule deer would have significant recreational and economic impacts. Greater investigation is therefore needed on how disease and/or competition may impact pop-

ulations of native ungulates. Without an understanding of the exact nature of disease relationships between gemsbok and native ungulates, it is unlikely that there can be successful management or recovery of gemsbok or native ungulates in the Chihuahuan Desert system in New Mexico.

ACKNOWLEDGMENTS

We thank all U.S. Military, U.S. National Park Service, New Mexico Department of Game and Fish, New Mexico State University, and U.S. Department of Agriculture employees who assisted in collection or evaluation of this data. Funding for this project was provided by the U.S. Army White Sands Missile Range (Approved for public release—Distribution is unlimited. OPSEC review conducted 13 August 2002), New Mexico Department of Game and Fish Department, and the U.S. Geological Survey. The Agricultural Experiment Station at New Mexico State University provided additional financial support.

LITERATURE CITED

- ALTON, G. G., L. M. JONES, R. D. ANGUS, AND J. M. VERGER. 1988. Techniques for the brucellosis laboratory. Institut National de la Recherche Agronomique, Paris, France, 190 pp.
- BALL, M. C., M. W. LANKESTER, AND S. P. MAHONEY. 2001. Factors affecting the distribution and transmission of *Elaphostrongylus rangiferi* (Protostrongylidae) in caribou (*Rangifer tarandus caribou*) of Newfoundland, Canada. *Canadian Journal of Zoology* 79: 1265–1277.
- BURKETT, D. W., R. VALDEZ, B. C. THOMPSON, AND K. G. BOYKIN. 2002. Gemsbok: The management challenge of an exotic ungulate in the American Southwest. *In Sustainable utilization—conservation in practice*, H. Ebedes, B. Reilly, W. van Hoven and B. Penzhorn (eds.). Proceedings of the Fifth Wildlife Ranching Symposium, South African Game Rancher's Organisation, Pretoria, South Africa, pp. 166–171.
- CALISHER, C. H., AND D. E. TAYLOR. 1993. Serologic evidence that bluetongue, epizootic hemorrhagic disease, and Malpais Springs viruses are persistently transmitted to wild and feral deer, oryx, and pronghorns in central New Mexico. *In Arbovirus research in Australia*, Proceedings Sixth Symposium, M. F. Ureh and B. H. Kay (eds.). Q.I.M.R., Brisbane, Australia, pp. 223–226.
- CARBREY, E. A., L. N. BROWN, T. L. CHOW, R. F. KAHRS, D. G. MCKERCHER, L. K. SMITHIES, AND T. W. TAMOGLIA. 1971. Recommended standard laboratory techniques for diagnosing in-

- fectious bovine rhinotracheitis, bovine viral diarrhoea and shipping fever (parainfluenza-3). Proceedings of the United States Animal Health Association 75: 629–648.
- CLARK, G. C., C. H. CALISHER, C. L. CRABBS, K. M. CANESTORP, R. B. TESH, R. A. BOWEN, AND D. E. TAYLOR. 1988. Malpais Spring virus: A new vesiculovirus from mosquitoes collected in New Mexico and evidence of infected indigenous and exotic ungulates. American Journal of Tropical Medicine and Hygiene 39: 586–592.
- CLARK, K. A., R. M. ROBINSON, R. G. MARBURGER, L. P. JONES, AND J. H. ORCHARD. 1970. Malignant catarrhal fever in Texas cervids. Journal of Wildlife Diseases 6: 376–383.
- FOREYT, W. J. 2001. Veterinary parasitology reference manual. Iowa State University Press, Ames, Iowa, 235 pp.
- FORRESTER, S. G., AND M. W. LANKESTER. 1997. Extracting protostrongylid nematode larvae from ungulate feces. Journal of Wildlife Diseases 33: 511–516.
- GOCHENOUR, W. S., C. A. GLEISER, AND N. K. WARD. 1958. Laboratory diagnosis of leptospirosis. Annals of the New York Academy of Sciences 70: 421–426.
- HAMILTON, W. J., III, R. BUSKIRK, AND W. H. BUSKIRK. 1977. Intersexual dominance and differential mortality of gemsbok *Oryx gazella* at Namib desert waterholes. Madoqua 10: 5–19.
- HEUSCHELE, W. P., AND H. W. REID. 2001. Malignant catarrhal fever. In Infectious diseases of wild mammals, E. S. Williams and I. K. Barker (eds.). Iowa State University, Ames, Iowa, pp. 157–164.
- HIDALGO, R. J., AND G. T. DIMOPOULLOS. 1967. Complement-fixation microprocedures for anaplasmosis. American Journal of Veterinary Research 28: 245–251.
- HOBAN, P. A. 1990. A review of desert bighorn sheep in the San Andres Mountains, New Mexico. Transactions of the Desert Bighorn Council 34: 14–22.
- HOBBS, N. T., AND M. W. MILLER. 1992. Interactions between pathogens and hosts: Simulation of pasteurellosis epizootics in bighorn sheep populations. In Wildlife 2001: Populations, D. R. McCullough and R. H. Barrett (eds.). Elsevier Science, London, England, pp. 997–1007.
- HOFF, G. L., D. O. TRAINER, AND M. M. JOCHIM. 1974. Bluetongue virus and white-tailed deer in an enzootic area of Texas. Journal of Wildlife Disease 10: 158–163.
- HOWERTH, E. W., D. E. STALLKNECHT, AND P. D. KIRKLAND. 2001. Bluetongue, epizootic hemorrhagic disease, and other orbivirus-related diseases. In Infectious diseases of wild mammals, E. S. Williams and I. K. Barker (eds.). Iowa State University, Ames, Iowa, pp. 77–97.
- LANGE, R. E., A. V. SANDOVAL, AND W. P. MELENEY. 1980. Psoroptic scabies in bighorn sheep (*Ovis canadensis mexicana*) in New Mexico. Journal of Wildlife Diseases 16: 77–82.
- LEE, K., C. CARGILL, AND H. ATKINSON. 1985. Evaluation of an enzyme linked-immunoabsorbent assay for the diagnosis of *Brucella ovis* infection in rams. Australian Veterinary Journal 62: 91–93.
- LI, H., D. T. SHEN, D. P. KNOWLES, J. R. GORHAM, AND T. B. CRAWFORD. 1994. Competitive inhibition enzyme-linked immunosorbent assay for antibody in sheep and other ruminants to a conserved epitope of malignant catarrhal fever virus. Journal of Clinical Microbiology 32: 1674–1679.
- , N. DYER, J. KELLER, AND T. B. CRAWFORD. 2000. Newly recognized herpesvirus causing malignant catarrhal fever in white-tailed deer (*Odocoileus virginianus*). Journal of Clinical Microbiology 38: 1313–1318.
- , J. KELLER, D. P. KNOWLES, AND T. B. CRAWFORD. 2001. Recognition of another member of the malignant catarrhal fever virus group: An endemic gammaherpesvirus in domestic goats. Journal of General Virology 82: 227–232.
- , K. GAILBREATH, L. C. BENDER, K. WEST, J. KELLER, AND T. B. CRAWFORD. 2003. Evidence of three new members of the malignant catarrhal fever virus group in muskox (*Ovibos moschatus*), Nubian ibex (*Capra nubiana*), and gemsbok (*Oryx gazella*). Journal of Wildlife Diseases 39: 875–880.
- MILLER, M. W. 2001. Pasteurellosis. In Infectious diseases of wild mammals, E. S. Williams and I. K. Barker (eds.). Iowa State University, Ames, Iowa, pp. 330–339.
- PEARSON, J. E., G. A. GUSTAFSON, A. L. SHAFER, AND A. D. ALSTAD. 1992. Diagnosis of bluetongue and epizootic hemorrhagic disease. In Bluetongue, African horse sickness, and other related orbiviruses, T. E. Walton and B. I. Osburn (eds.). CRC Press, Boca Raton, Florida, pp. 533–546.
- PLOWRIGHT, W. 1981. Herpesviruses of wild ungulates, including malignant catarrhal fever virus. In Infectious diseases of wild mammals, J. W. Davis, L. H. Karstad and D. O. Trainer (eds.). Iowa State University, Ames, Iowa, pp. 126–146.
- RAMSAY, E. C., S. J. RODGERS, A. E. CASTRO, E. L. STAIR, AND B. M. BAUMEISTER. 1985. Perinatal bluetongue viral infection in exotic ruminants. Journal of the American Veterinary Medical Association 187: 1249–1251.
- ROBINSON, R. M., T. L. HAILEY, C. W. LIVINGSTON, AND J. W. THOMAS. 1967. Bluetongue in the desert bighorn sheep. Journal of Wildlife Management 31: 165–168.
- RUPPRECHT, C. E., K. STOHR, AND C. MEREDITH. 2001. Rabies. In Infectious diseases of wild mammals, E. S. Williams and I. K. Barker (eds.). Iowa State University, Ames, Iowa, pp. 3–36.
- SHULAW, W. P., J. C. GORDON, S. BECH-NIELSEN, C.

- I. PRETZMAN, AND G. F. HOFFSIS. 1986. Evidence of paratuberculosis in Ohio's white-tailed deer as determined by an enzyme-linked immunosorbent assay. *American Journal of Veterinary Research* 47: 2539–2542.
- SMITH, J. C., R. VALDEZ, J. L. HOLECHEK, P. J. SWANK, AND M. CARDENAS. 1998. Diets of native and non-native ungulates in southcentral New Mexico. *Southwestern Naturalist* 43: 163–169.
- SPRAKER, T. R., AND C. P. HIBLER. 1982. An overview of the clinical signs, gross and histological lesions of the pneumonia complex of bighorn sheep. *Biennial Symposium of the Northern Wild Sheep and Goat Council* 3: 163–172.
- TAYLOR, C. R. 1969. The eland and the oryx. *Scientific American* 20: 89–95.
- THORNE, E. T. 1982. Bluetongue. *In* *Disease of wildlife in Wyoming*, E. T. Thorne, N. Kingston, W. R. Jolley and R. C. Bergstrom (eds.). Wyoming Game and Fish Department, Laramie, Wyoming, pp. 5–9.
- WALTHER, F. R. 1980. Aggressive behavior of oryx antelope at water-holes in the Etosha National Park. *Madoqua* 11: 271–302.
- WESTENBRINK, J. M., M. A. BRINKHOF, P. J. STRAV-
ER, J. QUAK, AND P. W. DE LEEUW. 1985. Comparison of a newly developed enzyme-linked immunosorbent assay with complement fixation and neutralisation tests for serology of bovine respiratory syncytial virus infections. *Research in Veterinary Science* 38: 334–340.
- WHITE SANDS MISSILE RANGE. 1991. Viral analysis of collected wildlife sera. White Sands Missile Range Environmental Services report number 200-8. White Sands Range Environment and Safety Directorate, White Sands, New Mexico, 6 pp.
- WILLIAMS, E. S., J. K. KIRKWOOD, AND M. W. MILLER. 2001. Transmissible spongiform encephalopathies. *In* *Infectious diseases of wild mammals*, E. S. Williams and I. K. Barker (eds.). Iowa State University, Ames, Iowa, pp. 292–302.
- WOOD, J. E., R. J. WHITE, AND J. L. DURHAM. 1970. Investigations preliminary to the release of exotic ungulates in New Mexico. *Bulletin No. 13*, New Mexico Department of Game and Fish, Sante Fe, New Mexico, 64 pp.

Received for publication 30 October 2002.