



A Survey for West Nile Virus in Bats from Illinois

Authors: Bunde, Jennifer M., Heske, Edward J., Mateus-Pinilla, Nohra E., Hofmann, Joyce E., and Novak, Robert J.

Source: Journal of Wildlife Diseases, 42(2) : 455-458

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.2.455>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A Survey for West Nile Virus in Bats from Illinois

Jennifer M. Bunde,¹ Edward J. Heske,^{2,4} Nohra E. Mateus-Pinilla,² Joyce E. Hofmann,² and Robert J. Novak³ ¹ Champaign County Forest Preserve District, Environmental Education Center, 2573 S. Homer Lake Road, Homer, Illinois 61849, USA; ² Center for Wildlife and Plant Ecology, Illinois Natural History Survey, 1816 S. Oak Street, Champaign, Illinois 61820, USA; ³ Center for Ecological Entomology, Illinois Natural History Survey, 1816 S. Oak Street, Champaign, Illinois 61820, USA; ⁴ Corresponding author (email: eheske@uiuc.edu)

ABSTRACT: A blocking enzyme-linked immunosorbent assay was used to test 97 serum samples from big brown bats (*Eptesicus fuscus*) captured in six counties in Illinois between May 2002 and February 2004 for West Nile virus (WNV) antibodies. One female big brown bat tested positive for WNV antibodies. Samples of kidney, liver, and heart tissue were collected from 312 bats of seven species that were submitted to the Illinois (USA) Department of Public Health or the Illinois Department of Agriculture diagnostic laboratories between January 2001 and December 2003. Tissue samples were tested for WNV using TaqMan reverse transcriptase polymerase chain reaction and all were negative. Prevalence of WNV antibodies in the bats (1%) was lower than previously reported for other flaviviruses, but similar to the prevalence (2%) of WNV antibodies reported in bats from New Jersey and New York, USA. Additional research is needed to determine potential impact of WNV infections on bats and to determine whether they play a role in the WNV transmission cycle.

Key words: Antibodies, bats, Chiroptera, Illinois, survey, West Nile Virus.

Since 1999, when West Nile virus (WNV) was introduced into the northeastern USA, it has spread across the continental USA and has caused >14,000 human cases (CDC, 2003). In Illinois, USA, WNV was first detected in 2001, and the number of human cases peaked in 2002 (Illinois Department of Public Health, 2005). Mosquitoes are the primary vectors for WNV, and wild birds are the principal hosts (Hubalek and Halouzka, 1999; Campbell et al., 2002; McLean et al., 2002). Evidence of WNV infections in a variety of vertebrates (Steele et al., 2000; Komar et al., 2001; Ludwig et al., 2002; Lichtensteiger et al., 2003; Steinman et al., 2003; Heinz-Taheny et al., 2004) suggests that WNV is widespread in wildlife, and other vertebrate hosts might serve as an

overwintering and maintenance reservoir for WNV.

A role for bats in the maintenance, dispersal, and natural history of arboviruses has been suggested (Fontanelli et al., 1989; Geevarghese and Banjeree, 1990). In Israel, 8% of surveyed fruit bats (*Rousettus aegypticus*) tested positive for WNV antibodies (Akov and Goldwasser, 1966). WNV infection was reported in four live big brown bats (*Eptesicus fuscus*) and two dead little brown bats (*Myotis lucifugus*) in New York in 2000 (CDC, 2000). One little brown bat and one northern long-eared bat (*M. septentrionalis*) tested positive for WNV antibodies in New Jersey, USA, in 2002 (Pilipski et al., 2004). Other flaviviruses closely related to WNV also infect bats (Constantine, 1970; Geevarghese and Banjeree, 1990). In Ohio, nearly 10% of the big brown bats and little brown bats tested were positive for Saint Louis encephalitis virus (SLEV) antibodies (Herbold et al., 1983).

When bats hibernate, their metabolic rate drops considerably, reducing the rate at which bats can produce antibodies (Sulkin et al., 1963; McNab, 1982). Hibernating bats may remain viremic with flaviviruses such as SLEV during their dormant season (Herbold et al., 1983) and may serve as a source of virus transmission and amplification in the spring. For this reason, Sulkin and Allen (1974) suggested that hibernating and migrating bats might be involved in arbovirus overwintering or reintroduction into a particular area.

Blood samples were collected from live big brown bats in Illinois to estimate the prevalence of antibodies to WNV. We selected big brown bats as the focus of our

study because they often live in proximity to humans (Kurta and Baker, 1990), they hibernate in Illinois (Hoffmeister, 1989), and their large size allows for the collection of a sufficient volume of blood for testing without posing a risk to the health of the bat. To increase our sample size, we also collected tissue samples from bats that were submitted for rabies testing to the Illinois Department of Public Health (IDPH) and Illinois Department of Agriculture (IDA) for WNV.

Blood was collected from 97 live big brown bats that were either captured by hand in hibernacula or with mist nets. Bats were collected in Champaign (40.385°N, 87.975°W; $n=6$) and Vermillion (39.970°N, 87.579°W; $n=3$) counties during the summers of 2002 and 2003; in Edgar County (39.611°N, 87.696°W) during July 2002 ($n=11$) and July 2003 ($n=7$); at the LeRoy Oakes Forest Preserve (41.927°N, 88.347°W; $n=7$) and the Paul Wolff Forest Preserve (42.067°N, 88.367°W; $n=2$), Kane County, once each in July 2003; from a privately owned barn in Momence, Kane County (41.167°N, 87.663°W; $n=17$), and from Guthrie Cave, Union County (37.567°N, 89.220°W; $n=4$), in November 2002; and from Magazine Mine, UNIMIN Corporation, Alexander County (37.327°N, 89.259°W; $n=40$), in February 2004. Additional details and descriptions of sites are in Bunde (2004).

Species, sex, age class (juvenile or adult), and weight were recorded for each bat and a 50–100 μ l blood sample was then taken from a vein in the interfemoral membrane; bats were not anesthetized. Blood was collected in a 100- μ l heparinized capillary tube, immediately transferred to a centrifuge tube, and placed in a cooler on dry ice, followed by centrifugation to separate the sera. Samples were stored at -80 C until analysis. Bats sampled from hibernacula were placed in a bag with hand warmers to rouse them from torpor before blood was collected. All bats were released at their capture

sites after bleeding had stopped and their condition appeared stable.

Sera were tested for WNV antibodies using blocking enzyme-linked immunosorbent assays (ELISAs), using a 1:10 serum dilution and monoclonal antibodies 2B2 and 6B6C-1 following Blitvich et al. (2003). The 2B2 flavivirus-specific monoclonal antibody (MAb) reacts with WNV and Koutango virus (Blitvich et al., 2003). The 6B6C-1 MAb reacts broadly with many flaviviruses, including WNV and SLE virus (Blitvich et al., 2003). We used parallel testing to provide a more sensitive diagnostic strategy resulting in a higher negative predictive value (Smith, 1995). Bats that tested negative in both tests were considered seronegative. Only one bat, an adult female from the maternity roost at LeRoy Oakes Forest Preserve in Kane County, tested positive for antibodies to WNV. This bat tested positive to both the 6B6C-1 MAb and the 2B2 Mab. We did not test for virus in these samples.

We necropsied 312 bats submitted to the IDPH and the IDA from 35 counties between 9 January 2001 and 10 December 2003 (Bunde, 2004). Most samples came from Cook County ($n=85$ bats), followed by Winnebago ($n=73$), Will ($n=27$ bats), and McHenry ($n=21$) counties. From 1 to 15 bats were available from other counties. Seven species of bats were included in this analysis: 258 big brown bats, 27 red bats (*Lasiurus borealis*), 20 silver-haired bats (*Lasionycteris noctivagans*), four hoary bats (*Lasiurus cinereus*), one little brown bat, one evening bat (*Nycticeus humeralis*), and one eastern pipistrelle (*Pipistrellus subflavus*).

Samples of kidney, liver, and heart tissue were individually tested for WNV by TaqMan reverse transcriptase-polymerase chain reaction (RT-PCR) following Lanciotti et al. (2000). A North American WNV strain (NY99) was used as a positive control. Brain tissue, which was removed as part of rabies testing, was not available for this study, but kidney, liver, and heart tissue have been used to test bats for

arboviruses in previous studies (Rueger et al., 1966; Sulkin and Allen, 1974; Herbold et al., 1983). All bat tissues tested negative for WNV.

Bats may be involved in the epidemiology of WNV either as amplifying hosts or as a virus reservoir allowing for WNV maintenance during the winter. Reduced metabolic activity during hibernation may prolong the period of WNV infection. The clustering behavior of bats in maternity colonies also has been hypothesized to be conducive to transmission of viruses via arthropod vectors (Main, 1979). However, in this study, antibodies were detected in only one bat, and all attempts to detect WNV by RT-PCR were negative. Herbold et al. (1983) reported antibodies to SLEV in 9% of 390 live bats tested, but, similar to results from this study, SLEV antigen was not detected from >1,000 tissue samples.

Although WNV antibodies can be detected from bats using cELISA methods, the sensitivity and specificity of these tests for bats is unknown (Blitvich, pers. comm.). However, the low prevalence of antibodies detected in this study are consistent with previous reports from bats. Akov and Goldwasser (1966) used a hemagglutination inhibition test in their survey that reported three of 37 (8%) fruit bats tested positive for WNV in Israel, although none of six pipistrelles (*Pipistrellus* sp.) tested positive in their study. Pilipski et al. (2004) reported that only two of 83 (2%) bats from New York and New Jersey tested positive for WNV antibodies by plaque-reduction neutralization test.

The number of bats sampled in hibernacula in this study ($n=44$) was too small to critically test hypotheses about the role of hibernating bats in the WNV transmission cycle, and our results should be considered as preliminary. Because big brown bats hibernate in buildings in urban areas and because our survey was conducted during years when WNV was entering Illinois, additional work is recommended.

We thank the Illinois Department of Public Health, the Illinois Department of Agriculture, and Daniel Brault of All Seasons Wild Animal Control for assistance with sample collection. This project was supported by a grant from the US Centers for Disease Control and Prevention, PHS U50 CCU52051, and in part by the State of Illinois, Waste Tire Fund, Illinois Department of Natural Resources, to R.J.N.

LITERATURE CITED

- AKOV, Y., AND R. GOLDWASSER. 1966. Prevalence of antibodies to arboviruses in various animals in Israel. *Bulletin of the World Health Organization* 34: 901–909.
- BLITVICH, B. J., N. L. MARLENEE, R. A. HALL, C. H. CALISHER, R. A. BOWEN, J. T. ROEHRIG, N. KOMAR, S. A. LANGEVIN, AND B. J. BEATY. 2003. Epitope-blocking enzyme-linked immunosorbent assays for the detection of serum antibodies to West Nile virus in multiple avian species. *Journal of Clinical Microbiology* 41: 1041–1047.
- BUNDE, J. M. 2004. Do bats in Illinois have West Nile Virus? Unpublished MS Thesis, University of Illinois Urbana-Champaign, 49 pp.
- CAMPBELL, G. L., A. A. MARFIN, R. S. LANCIOTTI, AND D. J. GUBLER. 2002. West Nile Virus. *The Lancet Infectious Diseases* 2: 519–529.
- CDC. 2000. West Nile Virus. *Morbidity and Mortality Weekly Report* 49: 820–822.
- . 2003. West Nile Virus activity—United States, November 20–25, 2003. *Morbidity and Mortality Weekly Report* 52: 1160.
- CONSTANTINE, D. J. 1970. Bats in relation to the health, welfare, and economy of man. *In* *Biology of bats*, Vol. II, W. Wimsatt (ed.). Academic Press, New York, New York, pp. 320–420.
- FONTANELLI, D., F. RODHAIN, J. P. DIGOUTTE, C. MATHIOT, J. MORVAN, AND P. COULANGES. 1989. Transmission cycle of West Nile virus in Madagascar (Indian Ocean). *Annales de la Societe Belge de Medecin Tropicale* 69: 242–243.
- GEEVARGHESE, G., AND K. BANERJEE. 1990. The role of bats in the natural cycle of arbovirus. *Current Science* 59: 26–31.
- HEINZ-TAHENY, K. M., J. J. ANDREWS, M. J. KINSEL, A. P. PESSIER, M. E. PINKERTON, K. Y. LEMBERGER, R. J. NOVAK, G. DIZIKES, E. EDWARDS, AND N. KOMAR. 2004. West Nile Virus infection in free-ranging squirrels in Illinois. *Journal of Veterinary Diagnostic Investigation* 16: 186–190.
- HERBOLD, J. R., W. P. HEUSCHELE, R. L. BERRY, AND M. A. PARSONS. 1983. Reservoir of St. Louis

- encephalitis virus in Ohio bats. *American Journal of Veterinary Research* 44: 1889–1893.
- HOFFMEISTER, D. F. 1989. *Mammals of Illinois*. University of Illinois Press, Urbana and Chicago, Illinois, 348 pp.
- HUBALEK, Z., AND J. HALOUZKA. 1999. West Nile Fever—A reemerging mosquito-borne viral disease in Europe. *Emerging Infectious Diseases* 5: 644–648.
- ILLINOIS DEPARTMENT OF PUBLIC HEALTH. 2005. West Nile Virus. <http://www.idph.state.il.us/envhealth/wnv.htm>. Accessed 24 August 2005.
- KOMAR, N., N. A. PANELLA, AND E. BOYCE. 2001a. Exposure of domestic animals to West Nile Virus during and outbreak of human encephalitis, New York City, 1999. *Emerging Infectious Diseases* 7: 736–737.
- KURTA, A., AND R. H. BAKER. 1990. *Eptesicus fuscus*. *Mammalian Species* 356: 1–10.
- LANCIOTTI, R. S., A. J. KERST, R. S. NASCI, M. S. GODSEY, C. J. MITCHELL, H. M. SAVAGE, N. KOMAR, N. A. PANELLA, B. C. ALLEN, K. E. VOLPE, B. S. DAVIS, AND J. T. ROEHRIG. 2000. Rapid detection of West Nile Virus from human clinical specimens, field-collected mosquitoes, and avian samples by a TaqMan reverse transcriptase-PCR assay. *Journal of Clinical Microbiology* 38: 4066–4071.
- LICHTENSTEIGER, C. A., K. HEINZ-TAHENY, T. S. OSBORNE, R. J. NOVAK, B. A. LEWIS, AND M. L. FIRTH. 2003. Fatal West Nile Virus encephalitis and myocarditis in two canids (wolf and dog). *Emerging Infectious Diseases* 9: 1303–1306.
- LUDWIG, G. V., P. P. CALLE, J. A. MANGIAFICO, B. L. RAPHAEL, D. K. DANNER, J. A. HILE, T. L. CLIPPINGER, J. F. SMITH, R. A. COOK, AND T. MCNAMARA. 2002. An outbreak of West Nile Virus in a New York City captive wildlife population. *American Journal of Tropical Medicine and Hygiene* 67: 67–75.
- MAIN, A. J. 1979. Eastern equine encephalomyelitis virus in experimentally infected bats. *Journal of Wildlife Diseases* 15: 467–477.
- MCLEAN, R. G., S. R. UBICO, D. BOURNE, AND N. KOMAR. 2002. West Nile virus in livestock and wildlife. *Current Topics in Microbiological Immunology* 267: 271–308.
- MCNAB, B. K. 1982. Evolutionary alternatives in the physiological ecology of bats. In *Ecology of bats*, T. H. Kunz (ed.). Plenum Press, New York, New York, pp. 151–200.
- PILIPSKI, J. D., L. M. PILIPSKI, AND L. S. RISELEY. 2004. West Nile Virus antibodies in bats from New Jersey and New York. *Journal of Wildlife Diseases* 40: 335–337.
- RUEGER, M. E., T. A. OLSON, AND R. D. PRICE. 1966. Studies of potential avian, arthropod and mammalian hosts of mosquito-borne arboviruses in the Minnesota area. *American Journal of Epidemiology* 83: 33–37.
- SMITH, R. D. 1995. *Veterinary clinical epidemiology: A problem oriented approach*. 2nd Edition. CRC Press, Ann Arbor, Michigan, 279 pp.
- STEELE, K. E., M. J. LINN, R. J. SCHOPP, N. KOMAR, T. W. GEISBERT, R. M. MANDUCA, P. P. CALLE, B. L. RAPHAEL, T. L. CLIPPINGER, T. LARSEN, J. SMITH, R. S. LANCIOTTI, N. A. PANELLA, AND T. S. MCNAMARA. 2000. Pathology of fatal West Nile Virus infections in native and exotic birds during the 1999 outbreak in New York City, New York. *Veterinary Pathology* 37: 208–224.
- STEINMAN, A., C. BARNET-NOACH, S. TAL, O. LEVI, L. SIMANOV, S. PERK, M. MALKINSON, AND N. SHPIGEL. 2003. West Nile Virus infection in crocodiles. *Emerging Infectious Diseases* 9: 887–888.
- SULKIN, E. S., AND R. ALLEN. 1974. Virus infection in bats. *Monographs in Virology* 8: 1–103.
- , ———, AND R. SIMS. 1963. Studies of arthropod-borne virus infection in Chiroptera: I. Susceptibility of insectivorous species to experimental infection with Japanese B and St. Louis Encephalitis viruses. *The American Journal of Tropical Medicine and Hygiene* 12: 800–814.

Received for publication 16 September 2004.