

PREVALENCE AND DIVERSITY OF AVIAN HEMATOZOAN PARASITES IN ASIA: A REGIONAL SURVEY

Authors: Ishtiaq, Farah, Gering, Eben, Rappole, Jon H., Rahmani, Asad R., Jhala, Yadvendradev V., et al.

Source: Journal of Wildlife Diseases, 43(3) : 382-398

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-43.3.382>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PREVALENCE AND DIVERSITY OF AVIAN HEMATOZOAN PARASITES IN ASIA: A REGIONAL SURVEY

Farah Ishtiaq,^{1,7,8} Eben Gering,¹ Jon H. Rappole,² Asad R. Rahmani,³ Yadvendradev V. Jhala,⁴ Carla J. Dove,⁵ Chris Milensky,⁵ Storrs L. Olson,⁵ Mike A. Peirce,⁶ and Robert C. Fleischer¹

¹ Genetics Program, National Zoological Park, National Museum of Natural History, Smithsonian Institution, 3001 Connecticut Ave. NW, Washington, D.C. 20008, USA.

² Conservation and Research Center, Front Royal, Virginia 22630, USA

³ Bombay Natural History Society, Hornbill House, S. B. Singh Road, Mumbai 400023, India

⁴ Wildlife Institute of India, PO Box 18, Chandrabani, Dehra Dun 248001, Uttaranchal, India

⁵ Division of Birds, National Museum of Natural History, Smithsonian Institution, Room E 607, MRC 116, Washington, D.C. 20013-7012, USA

⁶ 6 Normandale House, Normandale, Bexhill-on-Sea, East Sussex, TN39 3NZ, UK

⁷ Current address: Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford, South Parks Road, Oxford, OX1 3PS, UK.

⁸ Corresponding author (email: farah.ishtiaq@zoo.ox.ac.uk)

ABSTRACT: Tissue samples from 699 birds from three regions of Asia (Myanmar, India, and South Korea) were screened for evidence of infection by avian parasites in the genera *Plasmodium* and *Haemoproteus*. Samples were collected from November 1994 to October 2004. We identified 241 infected birds (34.0%). Base-on-sequence data for the cytochrome *b* gene from 221 positive samples, 34 distinct lineages of *Plasmodium*, and 41 of *Haemoproteus* were detected. Parasite diversity was highest in Myanmar followed by India and South Korea. Parasite prevalence differed among regions but not among host families. There were four lineages of *Plasmodium* and one of *Haemoproteus* shared between Myanmar and India and only one lineage of *Plasmodium* shared between Myanmar and South Korea. No lineages were shared between India and South Korea, although an equal number of distinct lineages were recovered from each region. Migratory birds in South Korea and India originate from two different migratory flyways; therefore cross-transmission of parasite lineages may be less likely. India and Myanmar shared more host species and habitat types compared to South Korea. Comparison between low-elevation habitat in India and Myanmar showed a difference in prevalence of haematozoans.

Key words: Avian malaria, hematozoan parasites, India, Myanmar, South Korea.

INTRODUCTION

The avian hematozoan parasites *Plasmodium* and *Haemoproteus* are two well-studied and globally distributed genera of blood parasites (Atkinson and Van Riper 1991). Recent molecular studies using DNA sequencing to identify and characterize parasite lineages (Bensch et al., 2000; Ricklefs and Fallon, 2002; Fallon et al., 2003; Beadell et al., 2004) have revealed more detailed information than was provided by morphology alone (Perkins and Schall, 2002). Studies have been conducted on the morphologic classification of these two parasite genera (Atkinson and Van Riper, 1991), host specificity (Bensch et al., 2000; Ricklefs and Fallon, 2002; Waldenström et al., 2002; Beadell et al., 2004), and geographical distribution (Fallon et al., 2005; Durrant et al., 2006).

Molecular techniques also have been used to differentiate between native and introduced parasite lineages in native and introduced populations of common myna (*Acridotheres tristis*; Ishtiaq et al., 2006) and to detect cross-species transmissions between migratory and resident birds (Waldenström et al., 2002).

Each year billions of migratory birds migrate between tropical wintering areas to temperate summer breeding grounds (McClure, 1974b). Although Asia is a wintering and staging ground for many migratory birds, and a major source of exotic species that have been introduced into many other regions, it has remained the least-explored region as far as the genetic diversity of avian blood parasites is concerned. The global movements of diverse avian species that may be infected with blood parasites should be a cause of concern

given the potential of migrants and introduced species to spread novel diseases worldwide (McClure, 1974a; Rappole et al., 2000). This may be especially relevant to isolated regions where native species may have a reduced immune defense against such parasites (Van Riper et al., 1986).

Hawaii is an excellent example of biological invasion where the introductions of avian malaria (*Plasmodium relictum*) and avian pox into highly susceptible native populations led to the endangerment and possible extinction of many native birds (Van Riper et al., 1986). About 125 bird species were introduced to the Hawaiian Islands in the late nineteenth and early twentieth centuries (Caum, 1933; Berger, 1981; Pratt, 1994); the majority of these originated from India and Southeast Asia. As part of a global survey to determine the origin of *Plasmodium relictum* in Hawaii, India has been included as a region of interest as at least 12 avian species were introduced from Hawaii from this source (Long, 1981). In this study, we examined the genetic diversity of hematozoan parasites of India, Myanmar, and South Korea. Although McClure et al. (1978) conducted a large-scale microscopy-based survey of the hematozoans in birds from eastern and southern Asia, molecular-based surveys have not been previously conducted in Asia. Our goals were to 1) determine the prevalence of infection within and across regions, and in families of host species shared between regions, 2) determine the phylogenetic relationships of the hematozoan parasites found in these regions, 3) estimate the degree of lineage sharing between regions, and 4) determine parasite diversity among regions.

MATERIALS AND METHODS

Sampling

We collected blood samples from 183 birds in India from seven sites (three sites near Pinjore, in Haryana, Meerut in Uttar Pradesh, and three sites near Guwahati in Assam) between May and June 2003 (before the

monsoon during the dry season) that varied from dry deciduous forest to scrubland and marshland (Fig. 1A). Each bird captured was banded for individual identification with rings provided by the Bombay Natural History Society (BNHS), sexed, aged, measured, weighed, and examined for flight or body molt. Approximately 50–100 μ l of whole blood was drawn by humeral venipuncture into a microvette with EDTA anticoagulant and used for the preparation of blood smears and DNA analysis. In South Korea, 181 tissue samples were collected in October 2004 from two airbases (Kunsan and Osan) (Fig. 1B). We examined 335 bird tissue samples (all blood impregnated) collected from four locations in Myanmar between November 1994 and March 2001 (in winter–dry season). Myanmar's climate and vegetation vary greatly from the temperate north and high-altitude zones to equatorial regions in the far south. The four sampling locations (Hkakabo Razi National Park, Taungoo area, Alaungdaw Kathapa National Park, and Chatthin Wildlife Sanctuary) varied from altitudes of 200 to 3,500 m and habitats ranging from degraded forest to alpine forest (Fig. 1C).

Screening of blood smears by microscopy

Thin blood smear slides were obtained from each bird sampled in India. These were fixed with methanol, stained for 1 hr with 6% phosphate buffered Giemsa (pH 7.0) and scanned at 400 \times for 10 min to diagnose malarial infections. Parasite species were identified using morphologic characteristics (Garnham, 1966). A minimum of 50,000 erythrocytes were examined on each slide (Atkinson et al., 1995). The level of parasitemia was based upon results of these examinations. Blood smears were not available for birds sampled in Myanmar and South Korea.

Molecular analysis

Host and parasite DNA was extracted from tissue and blood samples using Qiagen DNeasy kits (Qiagen, Valencia, California USA) following manufacturer guidelines. To ensure the quality of DNA extractions, we amplified a small fragment (347 bp) of *cyt b* DNA of host using primers cytb-1 and cytb2 following the methods described in Kocher et al. (1989). These PCRs were successful in all cases. Two primers sets were used for parasite screening of samples; these included primer set F2/R2 (91bp, cytochrome b gene [*cyt b*]) and primer set 850F/1024R (167 bp; Beadell et al., 2004; COIII gene). For positive samples, we attempted to amplify a larger fragment




FIGURE 1. Sampling sites in A) India, B) South Korea, and C) Myanmar.

(533 bp) of the *cyt b* sequence using primers 3760F/4292RW2 (Beadell et al., 2004). Because it was not always possible to amplify the larger 533 bp fragments, we attempted to amplify smaller fragments of either 256 bp using primers 3760F with R1 (5'-CTT TTT AAG GTT GGG TCA CTT-3') or 295 bp using primers F3 (5'-CCA GGA CTT GTT TCA TGG AT-3') with 4292RW2. When these primer sets failed to amplify DNA, additional reactions using F3 with R3 (5'-GCA TAT CTA

TCT ACT GTA ATT GC-3'; 313 bp) and FIFI (5'-GGG TCA AAT GAG TTT CTGG-3') with 4292RW2 (351 bp) were used. Details of the PCR protocols are reported in Beadell et al. (2004) and Ishtiaq et al. (2006). Negative and positive controls were included in all reactions. Parasite DNA (*P. relictum*) from an infected Hawaii Amakihi (*Hemignathus virens*) was used as a positive control. Samples from India were screened and amplified in WII's conservation genetics lab, and Myanmar

TABLE 1. Prevalence of *Plasmodium* and *Haemoproteus* based on polymerase chain reaction (PCR) detection for sampling sites in Myanmar and India.

Sampling site ^a	Date	Altitude (m)	Habitat	Total samples	<i>Plasmodium</i>	<i>Haemoproteus</i>
Myanmar						
Hkakabo Razi NP	February–March 2001	>3,500	Alpine	178	29	1
Taungoo Area	November 1994	200–500	Degraded	13	0	1
Alaungdaw Kathapa NP	December 1994, 1997	200–500	Lowland deciduous	23	4	1
Chatthin WLS	November–December 1994	500–1,000	Lowland evergreen	121	35	8
India						
Vulture (H) center	May–June 2003	<200	Dry deciduous	31	7	14
Pinjore (H)	May–June 2003	<200	City	16	3	4
Nagaon (A)	May–June 2003	<200	City	27	5	0
Meerut (UP)	May–June 2003	<200	Scrubland	64	18	11
Guwahati (A)	May–June 2003	<200	City	13	0	1
Deobali Jalah (A)	May–June 2003	<200	Marshland	17	4	0
Ambala (H)	May–June 2003	<200	City	15	8	3

^a H = Haryana, A = Assam, UP = Uttar Pradesh.

and South Korea samples were analyzed at the genetics program at the Smithsonian Institution.

All samples with positive amplification for larger fragments were purified using Qiaquick kits (Qiagen) and sequenced bidirectionally on an ABI 3100 Sequencer (Applied Biosystems, Inc., Foster City, California, USA). Sequences were edited and aligned using the program SEQUENCHER version 4.1 and are available through Genbank (accession numbers EF380111–EF380209).

Phylogenetic analysis

Phylogenetic relationships were evaluated using samples for which we had at least 256 bp of cyt *b* sequence. Based on the phylogeny in Perkins and Schall (2002), the phylogenetic trees were rooted using either lizard *Plasmodium* (Genbank accession numbers AY099061, AY099060) or *Haemoproteus* (Genbank accession numbers AY099062, AY099057) sequences as outgroups. We used MODELTEST version 3.6 (Posada and Crandall, 1998) to determine the most appropriate evolutionary model for our data. The hierarchical likelihood ratio test using Akaike Information Criterion (AIC) selected the general time-reversible model (GTR+I+G) for both *Plasmodium* and *Haemoproteus* phylogenies separately. For *Plasmodium*, the proportion of invariable sites equaled 0.6407, and the

gamma shape parameter equaled 1.0639; for *Haemoproteus* the invariable sites equaled 0.4514, and the gamma shape parameter equaled 0.4899. These settings were implemented in a maximum likelihood analysis to estimate the topology of the avian malaria parasites. We used 100 replicates and the “fast” heuristic in PAUP* (Swofford, 1999) to estimate bootstrap support.

Statistical analysis

To assess differences in the prevalence of *Plasmodium* spp. and *Haemoproteus* spp. between different regions, we used contingency table analyses in SAS to compare frequency differences using χ^2 or Fisher exact tests. To assess whether prevalence of parasites differed across host families and regions, we performed analysis of variance (ANOVA) on arcsine square-root transformed prevalence rates at the host species level (Scheuerlein and Ricklefs, 2004). We estimated generalized linear models (SAS institute) with prevalence as the dependent variable and region and family as independent variables. Avian families were included in analyses only if they occurred in two or more regions and included ≥ 5 individuals per regions (Table 1). Individual contribution of variables was summarized as *F*-values and associated probabilities (type III sums of squares). Taxonomy follows the *Handbook of the Birds of the World* (<http://>

www.hbw.com/ibc/phtml/familia.phtml). EstimateS version 7.0 (2004) was used to calculate Shannon diversity ($\text{mean} \pm \text{s.d.}$) within regions for both parasite lineages and host species.

RESULTS

Prevalence of parasites

We screened a total of 699 birds from Myanmar, India, and South Korea. Of the 335 samples tested in Myanmar, the prevalence of infection was 122 (37.3%). Of these, 113 individuals were positive for *Haemoproteus* spp. (45 [40%]) and *Plasmodium* spp. (68 [60%]). Based on sequence data, nine individuals showed double peaks indicating that these individuals were infected with two parasite lineages. Sequencing results identified 28 unique lineages for *Plasmodium* and 21 for *Haemoproteus*; these were detected in 42 of 133 represented bird species. There was a significant difference in *Plasmodium* spp. ($P=0.012$, Fisher's exact test) but not in *Haemoproteus* spp. ($P=0.079$, Fisher's exact test) prevalence among populations at different study sites in Myanmar (Table 1). Significant differences in prevalence of *Haemoproteus* spp. ($P=0.232$, Fisher's exact Test) or *Plasmodium* spp. ($P=0.065$, Fisher's exact Test) between the pooled low (500–3,500 m) and high (200–500 m) elevation sites were not detected.

In India, of the 183 samples representing 43 species, 84 individuals of 33 species tested positive for *Plasmodium* spp. and *Haemoproteus* spp. Of the positive samples, 51 (28%) were *Plasmodium* spp. and 33 (18%) were *Haemoproteus* spp. infections. There were 11 individuals showing multiple infections; we were not able to determine sequences for 23 positive samples. Sequencing results indicated that positive samples represented 11 *Plasmodium* and 19 *Haemoproteus* lineages. There was no significant difference in *Plasmodium* spp. ($\chi^2=12.2$, $P=0.056$, $\text{df}=6$) prevalence among populations in India, but significant difference for the prevalence of *Haemoproteus* spp. between

sites was detected ($\chi^2=26.65$, $P=0.0002$, $\text{df}=6$).

Of the 181 samples of 46 bird species in South Korea, 76 individuals (42%) representing 34 species tested positive for *Plasmodium* spp. and *Haemoproteus* spp. Of the positive samples, 56 (31%) were *Plasmodium* spp. and 20 (11%) were *Haemoproteus* spp. infections. The positive samples included 10 *Plasmodium* and five *Haemoproteus* lineages.

There was no difference ($\chi^2=0.15$, $P=2.0$, $\text{df}=1$) in the parasite prevalence between host blood (India) and host tissue (Myanmar and South Korea) samples. There were significant differences for *Plasmodium* spp. ($\chi^2=9.27$, $P=0.009$, $\text{df}=2$) and *Haemoproteus* spp. ($\chi^2=29.56$, $P=0.001$, $\text{df}=2$) prevalence among the three regions, with India and South Korea having the higher prevalence for both lineages than Myanmar. Differences in prevalence between Myanmar and India at sites located between 200 and 500 m elevation were not detected for *Plasmodium* spp. ($P=0.48$, Fisher's exact test) but were observed for *Haemoproteus* spp. ($P=0.005$, Fisher's exact test). For lower-sited elevations, prevalence differences between South Korea and Myanmar and India (pooled together) also were not detected for *Plasmodium* spp. ($P=0.06$, Fisher's exact test), but differences were detected for *Haemoproteus* spp. ($P=0.04$, Fisher's exact test).

Of 52 avian families recorded, 17 were common to all three regions, but there was no overlap at the species level between regions (Table 2). Prevalence of *Plasmodium* spp. ($F=1.16$, $\text{df}=16$, $P=0.310$) and *Haemoproteus* spp. ($F=1.08$, $\text{df}=16$, $P=0.383$) did not differ among avian families. Among regions, differences in prevalence for the 17 families shared between regions were detected for both *Plasmodium* spp. ($F=12.94$, $\text{df}=2$, $P=0.001$) and *Haemoproteus* spp. ($F=12.89$, $\text{df}=2$, $P=0.001$). Regional grouping explained 35% and 27% of the total variance for *Plasmodium* spp. and *Haemoproteus*

TABLE 2. Prevalence of *Plasmodium* and *Haemoproteus* based on PCR detection for selected host families from three Asian regions (Myanmar, India, and South Korea).

Family	Region	Host species	Sample size	<i>Plasmodium</i> positive ^c	<i>Haemoproteus</i> positive ^c
Columbidae ^c	Myanmar	Spotted Dove <i>Streptopelia chinensis</i>	2	0	1
		Red Turtle Dove <i>Streptopelia tranquebarica</i>	1	1	0
	India	Red Turtle Dove	1	0	1
		Blue Rock Pigeon <i>Columba livia</i>	5	0	1
	S. Korea	Rufous Turtle Dove <i>Streptopelia orientalis</i>	4	1	0
Corvidae ^c	Total		13	2 (15)	3 (23)
	Myanmar	Collared Treepie <i>Dendrocitta frontalis</i>	2	0	0
	India	Rufous Treepie <i>Dendrocitta vagabunda</i>	5	0	3
	S. Korea	Black-billed Magpie <i>Pica pica</i>	3	1	1
		Jay <i>Garrulus glandarius</i>	5	1	0
Meropidae ^b	Total		15	2 (13) ^f	4 (26)
	Myanmar	Little Green Bee-eater <i>Merops orientalis</i>	5	2	0
	India	Little Green Bee-eater	1	0	0
Motacillidae ^a	Total		6	2 (33)	0
	Myanmar	Black-backed Wagtail <i>Motacilla lugens</i>	1	0	0
	S. Korea	Olive-backed Pipit <i>Anthus hodgsoni</i>	3	2	0
		Pipit <i>Anthus</i> sp.	2	0	0
		Red-throated Pipit <i>Anthus cervinus</i>	3	1	0
White Wagtail <i>Motacilla alba</i>		1	1	0	
Muscicapidae ^c	Total		9	4 (44) ^f	0
	Myanmar	Grey-headed Flycatcher <i>Culicicapa ceylonensis</i>	1	0	0
		Niltava	1	0	1
		Large Niltava <i>Niltava grandis</i>	5	0	0
		Little Forktail <i>Enicurus scouleri</i>	1	0	0
		Oriental Magpie-Robin <i>Cosychus saularis</i>	7	6	0
		White-rumped Shama <i>Copsychus malabaricus</i>	7	2	0
		Orange-flanked Bush Robin <i>Tarsiger cyanurus</i>	2	1	0
		Plumbeous Water Redstart <i>Rhyacornis fuliginosus</i>	1	0	0
		Daurian Redstart <i>Phoenicurus aureus</i>	1	0	0
		Red-breasted Flycatcher <i>Ficedula parva</i>	6	1	4
		Rufous-bellied Niltava <i>Niltava sundara</i>	7	1	0
		Rufous-gorgeted Flycatcher <i>Ficedula strophiatea</i>	5	1	0
		Slaty-backed Forktail <i>Enicurus schistaceus</i>	1	0	0
		Small Nilatava <i>Niltava macgregoriae</i>	1	0	0
		White-gorgeted Flycatcher <i>Ficedula monileger</i>	8	0	0
		White-tailed Robin <i>Cinclidium leucurum</i>	1	1	0
		Blue Flycatcher <i>Cyornis</i> sp.	1	0	0
		Oriental Magpie Robin	5	0	0
		Indian Robin <i>Saxicoloides fulicata</i>	2	0	2
		Brown Rock Chat <i>Cercomela fusca</i>	3	1	0
		S. Korea	Daurian Redstart <i>Phoenicurus aureus</i>	4	1
Blue-and-white Flycatcher <i>Cyanoptila cyanomelana</i>	2		0	1	
Total			75	16 (21) ^f	9 (12) ^f
		Common Stonechat <i>Saxicola torquata</i>	1	1	0
		Mugimaki Flycatcher <i>Ficedula mugimaki</i>	2	0	1

TABLE 2. Continued.

Family	Region	Host species	Sample size	<i>Plasmodium</i> positive ^e	<i>Haemoproteus</i> positive ^e
Nectariniidae ^b	Myanmar	Black-throated Sunbird <i>Aethopyga saturata</i>	3	0	0
		Crimson Sunbird <i>Aethopyga siparaja</i>	2	0	0
		Green-tailed Sunbird <i>Aethopyga nipalensis</i>	1	0	0
		Little Spiderhunter <i>Arachnothera longirostra</i>	1	0	0
		Streaked Spiderhunter <i>Arachnothera magna</i>	2	0	0
		Purple Sunbird <i>Nectarinia sperata</i>	4	0	2
Paridae ^a	Total		14	0	2 (15) ^f
	Myanmar	Great Tit <i>Parus major</i>	1	0	0
		Green-backed Tit <i>Parus monticolus</i>	3	0	0
	S. Korea	Sultan Tit <i>Melanochlora sultanea</i>	1	0	0
		Coal Tit <i>Parus ater</i>	2	1	0
		Great Tit	16	12	1
		Long-tailed Tit <i>Aegithalos caudatus</i>	4	0	0
		Marsh Tit <i>Parus palustris</i>	2	1	0
		Varied Tit <i>Parus varius</i>	7	4	0
	Total		36	18 (50) ^f	1 (3)
Passeridae ^d	India	House Sparrow <i>Passer domesticus</i>	21	4	2
		Tree Sparrow <i>Passer montanus</i>	3	0	0
	S. Korea	Tree Sparrow	2	0	0
	Total		26	4 (15)	2 (8)
Phasianidae ^a	Myanmar	Himalayan Monal <i>Lophophorus impejanus</i>	1	0	0
		Slater Monal <i>Lophophorus slateri</i>	1	0	0
		Kalij Pheasant <i>Lophura leucomelanos</i>	1	0	0
	S. Korea	Japanese Quail <i>Coturnix japonica</i>	1	1	0
		Ring-necked Pheasant <i>Phasianus colchicus</i>	4	1	0
	Total		8	2 (25)	0
	Myanmar	Eurasian Wryneck <i>Jynx torquilla</i>	2	1	0
		Greater Yellownape Woodpecker <i>Picus flavinucha</i>	1	0	0
		Grey-headed Woodpecker <i>Picus flavinucha</i>	12	0	0
		Yellow-crowned Woodpecker <i>Dendrocopos mahrattensis</i>	2	1	0
		White-browed Piculet <i>Sasia ochracea</i>	3	0	0
		Rufous Piculet <i>Sasia abnormis</i>	2	0	0
		Common Goldenback Woodpecker <i>Dinopium javanense</i>	1	0	0
		Grey-headed Woodpecker	2	1	0
Picidae ^a	S. Korea	Japanese Pygmy Woodpecker <i>Picoides kizuki</i>	2	0	0
	Total		27	4 (15)	0
Psittacidae ^b	Myanmar	Rose-ringed Parakeet <i>Psittacula krameri</i>	2	0	0
		Rose-ringed Parakeet <i>Psittacula krameri</i>	1	0	1
	India	Alexandrine Parakeet <i>Psittacula eupatria</i>	2	1	0
	Total		5	1 (20)	1 (20)

TABLE 2. Continued.

Family	Region	Host species	Sample size	<i>Plasmodium</i> positive ^c	<i>Haemoproteus</i> positive ^c
Pycnonotidae ^b	Myanmar	Black-crested Bulbul <i>Pycnonotus melanicterus</i>	3	0	1
		Red-vented Bulbul <i>Pycnonotus cafer</i>	6	0	0
		Grey-eyed Bulbul <i>Iole propinqua</i>	1	0	0
		<i>Alphoixus</i> (<i>Criniger</i> auct.) spp.	4	0	0
	Total	Red-vented Bulbul	10	3	2
Paradoxornithidae ^a	Myanmar	Black-throated Parrotbill <i>Paradoxornis nipalensis</i>	24	3 (12) ^f	3 (12)
		Lesser Rufous-throated Parrotbill <i>Paradoxornis atrosuperciliaris</i>	2	0	0
	S. Korea	Vinous-throated Parrotbill <i>Paradoxornis webbianus</i>	9	1	0
		Total		12	1 (8)
	Sturnidae ^b	Myanmar	Asian Pied Starling <i>Sturnus contra</i>	1	0
Black-collared Starling <i>Sturnus nigricollis</i>			1	0	0
India			Asian Pied Starling	4	0
Total		Common Myna <i>Acridotheres tristis</i>	20	8	2
		Bank Myna <i>Acridotheres gingianianus</i>	4	2	1
	Jungle Myna <i>Acridotheres fuscus</i>	1	0	0	
Sylviidae ^c	Myanmar		31	10 (32)	4 (13)
		Common Tailorbird <i>Orthotomus sutorius</i>	5	1	0
		Mountain Tailorbird <i>Orthotomus cuculatus</i>	2	0	0
		Thick-billed Warbler <i>Acrocephalus aedon</i>	1	1	0
		Rufous-faced Warbler <i>Abroscopus albugularis</i>	1	0	0
		Blyth's Leaf Warbler <i>Phylloscopus reguloides</i>	2	0	0
		Yellow-browed Warbler <i>Phylloscopus inornatus</i>	3	0	1
		Mountain Leaf Warbler <i>Phylloscopus trivirgatus</i>	2	0	0
		Leaf Warbler <i>Phylloscopus</i> spp.	6	0	0
		Grey-cheeked Warbler <i>Seicercus polio-genys</i>	6	1	0
		Jungle Babbler <i>Turdoides striatus</i>	2	0	2
		Black-chinned Babbler <i>Stachyris pyrrhops</i>	2	2	0
		Striated Grassbird <i>Megalurus pryeri</i>	2	2	0
		Bush Warbler <i>Cetti</i> sp.	1	0	0
	India	Common Tailorbird	2	1	0
Yellow-eyed Babbler <i>Chrysomma sinense</i>		4	2	1	
S. Korea		Yellow-browed Warbler	5	0	0
	Arctic Warbler <i>Phylloscopus borealis</i>	1	0	1	
	Oriental Great Reed Warbler <i>Acrocephalus orientalis</i>	1	1	0	
Total	Leaf Warbler	9	2	0	
		57	13 (23) ^f	5 (9)	

TABLE 2. Continued.

Family	Region	Host species	Sample size	<i>Plasmodium</i> positive ^c	<i>Haemoproteus</i> positive ^c
Timaliidae ^b	Myanmar	Coral-billed Scimitar Babbler <i>Pomatorhinus ferruginosus</i>	2	1	0
		Eyebrowed Wren Babbler <i>Napothera epilepidotai</i>	1	0	0
		Long-billed Wren Babbler <i>Rimantor malaoptilus</i>	2	1	0
		Golden Babbler <i>Stachris chrysaea</i>	3	1	0
		Grey-cheeked Fulvetta <i>Alcippe morrisonia</i>	11	1	0
		Rufous-throated Fulvetta <i>Alcippe rufogularis</i>	6	0	0
		Yellow-throated Fulvetta <i>Alcippe cinerea</i>	3	0	0
		Rufous-winged Fulvetta <i>Alcippe castaneiceps</i>	5	0	0
		Grey-throated Babbler <i>Stachyris nigriceps</i>	7	1	0
		Spot-necked Babbler <i>Stachyris striolata</i>	2	1	0
		Rufous-capped Babbler <i>Stachyris ruficeps</i>	4	0	0
		Greater Necklaced Laughingthrush <i>Garrulax pectoralis</i>	3	2	0
		Lesser Necklaced Laughingthrush <i>Garrulax monileger</i>	2	0	0
		Scaly Laughingthrush <i>Garrulax subunicolor</i>	1	0	0
		Striated Laughingthrush <i>Garrulax striatus</i>	1	1	0
		White-crested Laughingthrush <i>Garrulax leucolophus</i>	2	0	0
		Black-faced Laughingthrush <i>Garrulax affinis</i>	1	0	0
		Chestnut-backed Laughingthrush <i>Garrulax nuchalis</i>	2	0	0
		Chestnut-crowned Laughingthrush <i>Garrulax erythrocephalus</i>	4	0	0
		Grey-sided Laughingthrush <i>Garrulax caerulatus</i>	1	0	0
		Long-tailed Sibia <i>Heterophasia picaoides</i>	5	1	0
		Puff-throated Babbler <i>Pellorneum ruficeps</i>	9	1	0
		Pygmy Wren Babbler <i>Pnoepyga pusilla</i>	1	0	0
		Red-faced Liocichla <i>Liocichla phoenicea</i>	1	0	0
		Red-tailed Minla <i>Minla ignotincta</i>	1	0	0
		Silver-eared Mesia <i>Leiothrix argenteauris</i>	3	0	0
		Streaked Wren Babbler <i>Napothera brevicaudata</i>	1	0	0
		Wedge-billed Wren Babbler <i>Sphenocichla humei</i>	1	0	0
		White-bellied Yuhina <i>Yuhina zantholeuca</i>	2	0	0
		Black-chinned Yuhina <i>Yuhina nigrimenta</i>	1	0	0
		White-naped Yuhina <i>Yuhina bakeri</i>	1	0	0
		White-browed Scimitar Babbler <i>Pomatorhinus schisticeps</i>	1	0	0
		White-throated Babbler <i>Turdoides gularis</i>	2	0	0
		Yellow-bellied Warbler <i>Abroscopus superciliaris</i>	1	0	0
		Yellow-eyed Babbler <i>Chrysomma sinense</i>	7	3	0
	Total		35	14 (40)	0

TABLE 2. Continued.

Family	Region	Host species	Sample size	<i>Plasmodium</i> positive ^c	<i>Haemoproteus</i> positive ^c
Turdidae ^a	Myanmar	Long-tailed Thrush <i>Zoothera dixonii</i>	2	0	0
		Lesser Shortwing <i>Brachypteryx leucophrys</i>	2	0	0
		Blue Rock Thrush <i>Monticola solitarius</i>	1	0	0
		Blue Whistling Thrush <i>Myophonus caeruleus</i>	1	0	0
	S. Korea	Grey Thrush <i>Turdus cardis</i>	1	0	1
	Total		7	0	1 (14)

^a Family shared between Myanmar and South Korea.

^b Family shared between Myanmar and India.

^c Family shared among Myanmar, India, and South Korea.

^d Family shared between India and South Korea.

^e *n* (% positive).

^f $P \leq 0.001$, Fisher's exact test (among regions).

spp., respectively. Prevalence of parasites observed in families represented in a single region was analyzed separately from families shared among regions (Table 3). As found for shared families, the prevalence of *Plasmodium* spp. and *Haemoproteus* spp. did not differ significantly among these families. We tested the prevalence of parasites for families shared between two or more regions separately and found significant differences in only a few cases (Table 2).

Microscopy and PCR detection

Of 183 blood smears collected in India, 61 individuals were positive for either *Plasmodium* spp. or *Haemoproteus* spp. Of these, 22 (12%) *Plasmodium* and 39 (21.3%) *Haemoproteus* positive samples were detected by examination of blood smears. Results from blood smear examination and PCR were in agreement for only 98 (54%) of the 183 samples.

Phylogenetics

We identified 241 infected birds and obtained sequence for *cyt b* from 221 of these. Within the 197 samples for which we had at least 256 bp of sequence, 34 lineages of *Plasmodium* and 41 of *Haemo-*

proteus were identified. Within the *Plasmodium* tree (Fig. 2), there were four lineages (LIN 8, LIN 26, LIN 28, LIN 29) shared between India and Myanmar, and one lineage (LIN 15) shared between Myanmar and South Korea. In *Haemoproteus* (Fig. 3), a single lineage (LIN 1) was shared between India and Myanmar. The proportion of parasite lineages sharing identical sequences in more than one host species occurs more frequently ($\chi^2=15.1$, $P=0.035$) in the *Plasmodium* tree than in the *Haemoproteus* tree. This may indicate higher rates of host-switching and reduced host-specificity in *Plasmodium* (see Figs. 2 and 3 for details on host species).

For 20 additional birds, short (91 bp) sequences were obtained from primers F2/R2. These were used to identify parasite genera but were not included in the phylogenetic analysis. These results, however, suggest the presence of the following *Plasmodium* lineages in host species in South Korea: LIN 7—great spotted woodpecker [*Dendrocopos major*], common stonechat [*Saxicola torquata*], red throated pipit [*Anthus cervinus*]; LIN 10—bull-headed shrike [*Lanius bucephalus*], varied tit [*Parus varius*], great tit [*Parus major*], coal tit [*Parus ater*], marsh tit [*Parus*

TABLE 3. Prevalence of hematozoan parasites for avian families recorded within one region.

Region	Host family	Species	Sample size	<i>Plasmodium</i> positive (n)	<i>Haemoproteus</i> positive (n)
Myanmar	Alcedidae	Common Kingfisher <i>Alcedo atthis</i>	6	1	0
	Strigidae	Asian Barred Owlet <i>Glaucidium cuculoides</i>	4	1	0
		Oriental Scops Owl <i>Otus sunia</i>	2	0	0
		Spotted Owlet <i>Athene brama</i>	1	1	0
	Dicruridae	Ashy Drongo <i>Dicrurus leuco-phaeus</i>	2	2	0
		Bronzed Drongo <i>Dicrurus aeneus</i>	1	0	0
		Hair-crested Drongo <i>Dicrurus hottentottus</i>	2	0	0
		Lesser Racket-tailed Drongo <i>Dicrurus remifer</i>	2	0	0
		Common Green Magpie <i>Cissa chinensis</i>	2	0	0
		Gold-billed Magpie <i>Urocissa flavirostris</i>	1	0	0
	Aegithinidae	Common Iora <i>Aegithina tiphia</i>	1	2	0
	Prionopidae	Common Woodshrike <i>Te-phrodornis pondicerianus</i>	1	3	1
		Rosy Minivet <i>Pericrocotus roseus</i>	1	0	0
	Campephagidae	Scarlet Minivet <i>Pericrocotus flammeus</i>	3	0	0
		Small Minivet <i>Pericrocotus cinnamomeus</i>	3	1	0
		White-bellied Minivet <i>Pericrocotus erythropygius</i>	2	1	0
		White-browed Fantail <i>Rhipidura aureola</i>	3	1	0
		White-throated Fantail <i>Rhipidura albicollis</i>	3	1	0
	Sittidae	Beautiful Nuthatch <i>Sitta formosa</i>	2	0	0
		Chestnut-bellied Nuthatch <i>Sitta castanea</i>	5	0	0
India	Cisticolidae	Jungle Prinia <i>Prinia sylvatica</i>	1	0	0
		Plain Prinia <i>Prinia inornata</i>	4	2	2
		Ashy Prinia <i>Prinia socialis</i>	5	1	2
		Grey-breasted Prinia <i>Prinia hodgsonii</i>	1	0	0
		<i>Prinia</i> spp.	2	0	2
		Streaked Fantail Warbler <i>Cisticola juncidis</i>	2	0	0
	Zosteropidae	Oriental White-eye <i>Zosterops palpebrosus</i>	7	2	1
	Estrildidae	Red Munia <i>Estrilda amandava</i>	31	11	0
		Spotted Munia <i>Lonchura punctulata</i>	11	1	0
		White-throated Munia <i>Lonchura malabarica</i>	1	0	0
		Black-headed Munia <i>Lonchura malaca atricapilla</i>	4	1	0

TABLE 3. Continued.

Region	Host family	Species	Sample size	<i>Plasmodium</i> positive (n)	<i>Haemoproteus</i> positive (n)
South Korea	Emberizidae	Black-faced Bunting <i>Emberiza</i> <i>spodocephala</i>	9	4	0
		Chestnut Bunting <i>Emberiza</i> <i>rutila</i>	7	1	3
		Rustic Bunting <i>Emberiza rus-</i> <i>tica</i>	2	0	0
		Tristram's Bunting <i>Emberiza</i> <i>tristrami</i>	2	1	1
		Yellow-throated Bunting <i>Em-</i> <i>beriza elegans</i>	1	1	0
	Laridae	Black-tailed Gull <i>Larus cras-</i> <i>sirostris</i>	5	0	2
	Laniidae	Bull-headed Shrike <i>Lanius</i> <i>bucephalus</i>	10	3	1
	Fringillidae	Common Rosefinch <i>Carpoda-</i> <i>cus erythrinus</i>	1	1	0
		Oriental Greenfinch <i>Carduelis</i> <i>sinica</i>	7	1	1
	Scolopacidae	Dunlin <i>Calidris alpina</i>	5	0	0
		Red-necked Stint <i>Calidris ru-</i> <i>ficollis</i>	5	0	0
	Anatidae	Mallard <i>Anas platyrhynchos</i>	2	0	1
		Spot-billed Duck <i>Anas poeci-</i> <i>lorhyncha</i>	5	0	0

palustris], Daurian redstart [*Phoenicurus auroreus*], leaf warbler [*Phylloscopus* sp.], black-billed magpie [*Pica hudsonia*]; LIN 11—bull-headed shrike [*Lanius bucephalus*], ring-necked pheasant [*Phasianus colchicus*], yellow-browed warbler [*Phylloscopus inornatus*], great tit [*Parus major*]; LIN 12—great tit; LIN 34—vinous-throated parrotbill [*Paradoxornis webbianus*]; LIN 35—great tit. Results suggested the presence of one additional *Haemoproteus* lineage, LIN 41, from a black-billed magpie (*Pica hudsonia*).

Host and parasite diversity

The Shannon diversity indices indicate that host diversity was higher in Myanmar (5.86 ± 0.01) than in India (5.20 ± 0.03) or South Korea (5.20 ± 0.02); parasite diversity was lower in South Korea (3.91 ± 0.01) as compared to Myanmar (4.32 ± 0.02) or India (4.30 ± 0.03). The host and parasite ratio was 82% in India followed by South Korea (75%) and Myanmar (73%).

DISCUSSION

The overall prevalence (*Haemoproteus* or *Plasmodium*) was higher in India (tropical zone); the highest number of unique *Haemoproteus* lineages also was observed among birds sampled in India. The lower parasite prevalence observed for Myanmar could be attributed to differences in habitat as most of the birds from Myanmar were sampled from high altitudes where less exposure to vectors may have occurred. On the other hand, the comparison between low-elevation habitats of India and Myanmar (tropical zone) and South Korea (temperate zone) showed no significant difference in prevalence for either parasite. Our results are not entirely consistent with previous studies that have used PCR-based techniques. Durrant et al. (2006) found that tropical zones had higher parasite prevalence than temperate zones, whereas Ricklefs (1992) showed that temperate




FIGURE 2. Relationships of *Plasmodium* lineages based on maximum likelihood using the model GTR+I+G. Each lineage represents host species and GenBank number.

birds had higher parasite prevalence. These differences may result from seasonal differences in timing of sampling or diagnostic techniques. In this study the

correspondence between PCR and microscopy results for samples from India was poor. This may have resulted from a low-intensity peripheral parasitemia and




FIGURE 3. Relationships of *Haemoproteus* lineages based on maximum likelihood using the model GTR+I+G. Each lineage represents host species and GenBank number.

variation in diagnostic sensitivity and is consistent with results from the previous studies where blood smears were evaluated by both methods (Jarvi et al., 2002; Richard et al., 2002). We are confident that the prevalence results from this study are not biased by testing a combination of

blood and tissue samples; such differences have not been apparent in our previous studies (Ishtiaq et al., 2006; Pagenkopp, 2006) using such samples.

The prevalence of parasites differed at the regional level but not at host family level. This finding is consistent with pre-

vious studies (Hellgren, 2005; Durrant et al., 2006). Temporal and spatial variation in parasite prevalence has been demonstrated (Schall and Marghoob, 1995; Bensch and Akesson, 2003), and it is likely that different parasite lineages are associated with particular vector communities. This factor may explain the differences in the geographic distribution of parasite lineages at different sites. The *Haemoproteus* tree shows regional clades, whereas the *Plasmodium* clades are “scattered” between regions and host species.

Plasmodium and *Haemoproteus* are widely distributed blood parasites that have been reported in many families of birds. Migrants can form a link for the transmission of parasites between tropical and temperate resident birds, especially among taxonomically similar hosts. Some sharing of parasite lineages between resident and migratory songbirds has been observed in Nigeria, indicating parasite transmission between migrants and residents on the wintering grounds (Waldenström et al., 2002). As a major wintering ground for hundreds of avian species, we would expect more shared parasite lineages between India and Myanmar because migrants utilizing these area originate from the same biogeographic region and share similar host families. The *Plasmodium* phylogenetic tree shows four shared lineages between these two regions (Fig. 1A). Birds migrating to South Korea do not appear to use the same flyway as birds migrating to India and Myanmar, and this is reflected by the few shared lineages between these regions; only one *Plasmodium* lineage was shared between Myanmar and South Korea (Fig. 1A). Although we found similar indices of parasite diversity in India and South Korea, the host-parasite ratio was higher in India, indicating more parasites infecting several host species than in South Korea or Myanmar. India had high parasite prevalence and highest number of unique parasite lineages as well.

The phylogenetic tree differentiates *Haemoproteus* and *Plasmodium* lineages; a well-supported topology is not necessary for our analyses, but it clearly defines lineages shared between different regions. There is some evidence of geographic structuring of parasite lineages in the *Plasmodium* tree as indicated in other studies (e.g., Beadell et al., 2006). In our parasite phylogeny (Fig. 2), more lineages of *Plasmodium* are shared between multiple host species than are of *Haemoproteus*, which is indicative of host switching (Bensch et al., 2000; Ricklefs and Fallon, 2002; Waldenström et al., 2002; Beadell et al., 2004) and supports *Haemoproteus* as being relatively more host specific than *Plasmodium* (Atkinson and van Riper, 1991).

Variation in the abundance of dipteran vectors at a particular geographic location is positively correlated with variation in the prevalence of parasites in the avian host community (Bennett and Cameron, 1974) and also may be related to variation in the intensity of parasitic infection. Host habitat choice is known to influence parasite load in host species as it can affect the frequency of contact with flying vectors of hematozoan parasites, which are mainly associated with wet and closed habitats such as the forest canopy (Scott and Edman, 1991). We did find a slight difference in prevalence of *Haemoproteus* in high- compared with low-elevation habitat in Myanmar and India but prevalence of *Plasmodium* remained the same, suggesting more exposure to ceratopogonids and hippoboscids at low elevations. Comparison between low-elevation habitats in India and Myanmar may have been affected by the timing of sampling in two regions. Samples from India were collected between May and June, which is the peak breeding season for lowland resident species, while winter (November–March) was the sampling period in Myanmar and October in South Korea. Parasite load is known to increase in spring–summer owing to the increased abundance of flying vectors at that time (Bennett and Ca-

meron, 1974; Atkinson and Van Riper, 1991) and because higher levels of sexual steroid hormones are circulating in the bird's blood, which in general depresses the immune system and allows parasites to survive (Wedekind and Følstad, 1994; Saino et al., 1995).

The Asian region harbors a diverse community of avian hematozoan lineages that were distributed among 41% of the birds sampled in this study. Prevalence varied between regions but not at the host family level. An in-depth study on the distribution of vector communities and their relationship to hematozoa in these regions would be useful and may provide some insight into the regional distribution of specific hematozoan lineages.

ACKNOWLEDGMENTS

Part of this study was funded by a National Institutes of Health grant (1R01GM063258). We would like to thank the Ministry of Environment and Forests, Government of India; Myanmar Forestry Ministry, Nature and Wildlife Conservation Division (Myanmar Collection and Export Permit 11853-55/96, UYin-Office/TWMC/2303/2002, SI/4697/2004) for permits to catch birds for blood and tissue samples. The Smithsonian expedition to South Korea was supported by a grant from the U.S. Department of Defense—Legacy Resource Management Program (DACA87-03-H-0006). We thank C. McIntosh, K. Durrant, and D. K. Sharma for facilitating laboratory analyses. We would also like to thank R. Jakati, A. Choudhury, R. Bharagava, H. Singha, K. Lahkar, B. P. Lahkar, and S. Barua for help in organizing field camps in Assam, and Cho Sok Hun, 51 Fighter Wing, Osan, Air Base for organizing field work in South Korea.

LITERATURE CITED

- ATKINSON, C. T., AND C. VAN RIPER III. 1991. Pathogenicity and epizootiology of avian haematozoa: *Plasmodium*, *Leucocytozoon*, and *Haemoproteus*. In *Bird-parasite interactions: Ecology, evolution and behaviour*, J. E. Loye and M. Zuk (eds.), Oxford University Press, New York, pp. 19–48.
- , K. L. WOODS, R. J. DUSEK, L. S. SILEO, AND W. M. IKO. 1995. Wildlife disease and conservation in Hawaii: Pathogenicity of avian malaria (*Plasmodium relictum*) in experimentally infected iiwi (*Vestiaria coccinea*). *Parasitology* 111: S59–S69.
- BEADELL, J. S., E. GERING, J. AUSTIN, J. P. DUMBA-CHER, M. A. PEIRCE, T. K. PRATT, C. T. ATKINSON, AND R. C. FLEISCHER. 2004. Prevalence and differential host-specificity of two avian blood parasite genera in the Australo-Papuan region. *Molecular Ecology* 13: 3829–3844.
- , F. ISHTIAQ, R. COVAS, M. MELO, B. H. WARREN, C. T. ATKINSON, S. BENSCH, G. R. GRAVES, Y. V. JHALA, M. A. PEIRCE, A. R. RAHMANI, D. M. FONSECA, AND R. C. FLEISCHER. 2006. Global phylogeographic limits of Hawaii's avian malaria. *Proceeding of Royal Society London B* 273: 2935–2944.
- BENNETT, G. F., AND M. CAMERON. 1974. Seasonal prevalence of avian hematozoa in passeriform birds of Atlantic Canada. *Canadian Journal of Zoology* 52: 1259–1264.
- BENSCH, S., AND S. AKESSON. 2003. Temporal and spatial variation of Haematozoans in Scandinavian willow warblers. *Journal of Parasitology* 89: 388–391.
- , M. STJERNMAN, D. HASSELQUIST, O. OSTMAN, B. HANSSON, H. WESTERDAHL, AND R. T. PINHEIRO. 2000. Host specificity in avian blood parasites: A study of *Plasmodium* and *Haemoproteus* mitochondrial DNA amplified from birds. *Proceedings of the Royal Society of London B* 267: 1583–1589.
- BERGER, A. J. 1981. *Hawaiian birdlife*. 2nd Edition. University Hawaii Press, Honolulu, Hawaii, 260 pp.
- CAUM, E. L. 1933. The exotic birds of Hawaii. *Occasional Papers of the Bernice P. Bishop Museum, Honolulu, Hawaii* 10: 1–55.
- DURRANT, K. L., J. S. BEADELL, F. ISHTIAQ, G. R. GRAVES, S. L. OLSON, E. GERING, M. PEIRCE, C. ATKINSON, C. M. MILENSKY, B. K. SCHMIDT, C. GEBHARD, AND R. C. FLEISCHER. 2006. Avian malaria in South America: A comparison of temperate and tropical zones. *Ornithological Monographs* 60: 98–111.
- FALLON, S. M., R. E. RICKLEFS, B. L. SWANSON, AND E. BERMINGHAM. 2003. Detecting avian malaria: An improved PCR diagnostic. *Journal of Parasitology* 89: 1044–1047.
- , E. BERMINGHAM, AND R. E. RICKLEFS. 2005. Host specialization and geographic localization of avian malaria parasites: A regional analysis in the Lesser Antilles. *American Naturalist* 165: 466–480.
- GARNHAM, P. C. C. 1966. *Malaria parasites and other haemosporidia*. Blackwell Scientific, Oxford, UK, 1114 pp.
- HELLGREN, O. 2005. The occurrence of haemosporidian parasites in the Fennoscandian bluethroat (*Luscinia svecica*) population. *Journal of Ornithology* 146: 55–60.
- ISHTIAQ, F., J. S. BEADELL, A. J. BAKER, A. R.

- RAHMANI, Y. V., JHALA, AND R. C. FLEISCHER. 2006. Prevalence and evolutionary relationships of haematozoan parasites in native versus and introduced populations of common myna *Acridotheres tristis*. *Proceedings of the Royal Society of London B* 273: 587–594.
- JARVI, S. I., M. E. M. FARIAS, H. BAKER, H. B. FRIEFELD, P. E. BAKER, E. VAN GELDER, J. G. MASSEY, AND C. T. ATKINSON. 2003. Detection of avian malaria (*Plasmodium* spp.) in native land birds of American Samoa. *Conservation Genetics* 4: 629–637.
- , J. J. SCHULTZ, AND C. T. ATKINSON. 2002. PCR diagnostics underestimate the prevalence of avian malaria (*Plasmodium relictum*) in experimentally-infected passerines. *Journal of Parasitology* 88: 153–158.
- KOCHER, T. D., W. K. THOMAS, A. MEYER, S. V. EDWARDS, S. PAABO, AND A. C. WILSON. 1989. Dynamics of mt DNA evolution in animals: Amplification and sequencing with conserved primers. *Proceedings of the National Academy of Science USA* 86: 6196–6200.
- LONG, J. L. 1981. *Introduced birds of the World. The worldwide history, distribution and influence of birds introduced to new environments.* Reed, Sydney, Australia.
- MCCLURE, H. E. 1974a. *Migration and survival of the birds of Asia.* U. S. Army Component, SEATO Medical Research Lab, Bangkok, Thailand.
- . 1974b. *Migration and survival of the birds of Asia.* Applied Science Research. Corporation. Thailand, Bangkok.
- , P. POONSWAD, E. C. GREINER, AND M. LAIRD. 1978. *Haematozoa in the birds of eastern and southern Asia.* Memorial University of Newfoundland, St. John's, Canada.
- PAGENKOPP, K. 2006. Using parasite lineages to track the migratory patterns of the common yellowthroat (*Geothlypis trichas*). MS Thesis, American University, Washington, D.C., USA. 49 pp.
- PERKINS, S. L., AND J. J. SCHALL. 2002. A molecular phylogeny of malarial parasites recovered from cytochrome *b* gene sequences. *Journal of Parasitology* 88: 972–978.
- POSADA, D., AND K. A. CRANDALL. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14 (9): 817–818.
- PRATT, H. D. 1994. Avifaunal change in the Hawaiian Islands. *Studies in Avian Biology* No. 15: 103–118.
- RAPPOLE, J. H., S. R. DERRICKSON, AND Z. HUBALEK. 2000. Migratory birds and spread of West Nile Virus in the western hemisphere. *Emerging Infectious Diseases* 6: 319–328.
- RICHARD, F. A., R. N. M. SEHGAL, H. I. JONES, AND T. B. SMITH. 2002. A comparative analysis of PCR-based detection methods for avian malaria. *Journal of Parasitology* 88 (4): 819–822.
- RICKLEFS, R. E. 1992. Embryonic development period and the prevalence of avian blood parasites. *Proceedings of the National Academy of Science USA* 89: 4722–4725.
- , AND S. M. FALLON. 2002. Diversification and host switching in avian malaria parasites. *Proceedings of the Royal Society of London B* 269: 885–892.
- SAINO, N., A. P. MØLLER, AND A. M. BOLZERN. 1995. Testosterone effects on the immune system and parasite infections in the barn swallow (*Hirundo rustica*): An experimental test of the immunocompetence handicap. *Behavioural Ecology* 6: 397–404.
- SCHALL, J. J., AND A. B. MARGHOOB. 1995. Prevalence of malarial parasite over time and space: *Plasmodium mexicanum* in its vertebrate host, the western fence lizard *Sceloporus occidentalis*. *Journal of Animal Ecology* 64: 177–185.
- SCHEUERLEIN, A., AND R. E. RICKLEFS. 2004. Prevalence of blood parasites in European passeriform birds. *Proceedings of the Royal Society of London B* 271: 1363–1370.
- SCOTT, T. W., AND J. D. EDMAN. 1991. Effects of avian host age and arbovirus infection on mosquito attraction and blood-feeding success. In *Bird-parasite interactions*, J. E. Loye and M. Zuk (eds.), Oxford University Press, Oxford, UK, pp. 179–204.
- SWOFFORD, D. L. 1999. PAUP*: Phylogenetic analysis using parsimony (*and other methods). Sinauer, Sunderland, Massachusetts.
- VAN RIPER, III, C., S. G. VAN RIPER, M. L. GOFF, AND M. LAIRD. 1986. The epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecological Monographs* 56: 327–344.
- WALDENSTRÖM, J., S. BENSCH, S. KIBOI, D. HASSELQUIST, AND U. OTTOSSON. 2002. Cross-species infection of blood parasites between resident and migratory songbirds in Africa. *Molecular Ecology* 11: 1545–1554.
- WEDEKIND, C., AND I. FØLSTAD. 1994. Adaptive or nonadaptive immunosuppression by sex hormones? *American Naturalist* 143: 936–938.

Received for publication 17 August 2005.