

NEUROLOGIC DISEASE IN A WHITE-TAILED DEER MASSIVELY INFECTED WITH MENINGEAL WORM (*Pneumoststrongylus tenuis*) *

Author: PRESTWOOD, ANNIE K.

Source: Journal of Wildlife Diseases, 6(1) : 84-86

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.1.84>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NEUROLOGIC DISEASE IN A WHITE-TAILED DEER MASSIVELY INFECTED WITH MENINGEAL WORM

(*Pneumostrongylus tenuis*)*

Pneumostrongylus tenuis generally is not regarded as an important cause of central nervous disease in its natural host, the white-tailed deer (*Odocoileus virginianus*). The parasite develops in the neural parenchyma of the central nervous system and matures in the subdural space and venous sinuses primarily within the cranium. Infected deer usually are

not infected with large numbers of worms and rarely display neurologic signs. This report indicates, however, that abnormally heavy infections of *P. tenuis* in the subdural space and the venous sinuses of the cranium of white-tailed deer occur and can result in severe neurologic signs and death.

Materials and Methods

The deer under consideration was a pregnant three-year old doe (65 lbs. and in poor condition) from the Lake Russell Wildlife Management Area, Habersham County, Georgia. This animal was

brought to the laboratory for clinical and pathological study because of her poor condition and obvious locomotor difficulties.

Results

Clinical observations

The doe was circling to the right when found. Her gait was unsteady and she tended to fall. The animal's head was held to the right, and she demonstrated a moderate nystagmus. The deer was killed for necropsy by giving an intramuscular injection of 50 mg succinylcholine chloride.

Necropsy findings

The most significant findings were in the cranium. Masses of adult *Pneumostrongylus tenuis* were found in the subdural space near the posterior choroid plexus and immediately behind the pituitary. Fewer worms were found elsewhere in the subdural space. Worms also were extremely numerous in the intercavernous and dorsal sagittal sinuses. Eighty adult *P. tenuis* were recovered from the subdural space and an estimated 20-40

worms were entangled within the sinuses. The dura mater was dark brown in color. A salient feature was the orange-brown color of the leptomeninges extending posteriorly as far as the medulla oblongata.

Histological findings

The findings in the dura mater were similar to those reported by Alibasoglu et al. (1961, Cornell Vet. 51: 431-441) and Anderson (1963, Can. J. Zool. 41: 775-792). The dura mater was thickened and in many areas eggs and larvae of *P. tenuis* were found surrounded by connective tissue and frequently by giant cells. Focal accumulations of lymphocytes as well as numerous eosinophils were found throughout the dura.

Dr. T. J. Hulland, Department of Pathology, Ontario Veterinary College, kindly provided the following notes on

*This study was supported by an appropriation from the Congress of the United States. Funds were administered and Research coordinated under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract Nos. 14-16-0008-676 and 14-16-0008-777, Bureau of Sport Fisheries and Wildlife, U.S. Department of the Interior.



FIGURE 1. *Pigment deposition adjacent to vessels within the cerebral cortex.*
H & E x 100.




FIGURE 2. *Detail of pigment deposition associated with vessels of cerebral cortex.*
H & E x 450.

the leptomeninges and the brain. "The cerebral meningeal surfaces show evidence of intracellular and extracellular pigment granules in the leptomeninges and also in the outer zone of the cortex. In hematoxylin and eosin preparations, most of the pigment, both in the meninges and the brain parenchyma, appears to be in histiocytes which have small, eccentric nuclei. The pigment is golden-brown but in concentration appears almost black. Some collections of free pigment appear to have been extruded by degenerating macrophages. Granule diameter varies up to 2 μ . An unusual location for the pigment is the elastica and collagen of medium sized meningeal vessels. In this location it is present in relatively large quantities in some vessels. The pigment is Peridic Acid Schiff-negative, Aldehyde

fuchsin-negative, Phloxine-negative, but strongly iron-positive as determined by Perle's Prussian blue reaction. One can only conclude that the pigment is a product of red blood cell breakdown by macrophages, although no erythrocyte extravasation or intermediate stages of breakdown are visible" (Fig. 1, 2).

"A separate lesion is visible adjacent to the median sagittal sinus of the cerebral meninges. In this tissue is a focal area containing several larvae and many multinucleate giant cells. Several larvae are ensheathed by two or more giant cells and some giant cells contain remnants of dead larvae. Giant cells contain up to two hundred nuclei and are mixed with a few pigment-laden mononuclear cells. The pigment present here is similar to that described earlier."

Discussion

On the basis of the material available for study, there is little evidence to suggest that the abnormal neurologic signs displayed by the deer were due directly to trauma of the neural parenchyma by wandering *P. tenuis*. This is in contrast to damage produced by *P. tenuis* in moose (*Alces a. americana*), which consists of extensive trauma to the spinal cord (Anderson, 1964. *Path. Vet.* 1: 289-322; Anderson, 1965. *Can. J. Zool.* 43: 635-639). The presence of pigment in the meninges and cerebral cortex suggests, however, that the massive infection of worms in the cranium had induced some circulatory abnormality.

Possibly large masses of worms in the venous sinuses and the reaction they provoked had interfered mechanically with the blood flow.

It is worth mentioning that of approximately 2500 deer examined for *P. tenuis* (Prestwood and Smith, 1969. *J. Parasit.* 55: 720-725), this was the only animal with signs of neurologic disturbance attributed to meningeal worms. The orange-brown discoloration of the leptomeninges and cerebral cortex also has been noted in other deer with large numbers of *P. tenuis*. Possibly circulatory disturbances are not uncommon in heavy infections of meningeal worm in deer.

Acknowledgements

The author wishes to extend sincere appreciation to Dr. Roy C. Anderson, University of Guelph, for his criticisms and comments on the manuscript and for photomicrography. Gratitude also is expressed to Dr. T. J. Hulland, University of Guelph, for examining histological sections of meninges and brain tissue and providing comments on this material.

ANNIE K. PRESTWOOD

*Southeastern Cooperative Wildlife
Disease Study
Department of Pathology and
Parasitology
School of Veterinary Medicine
University of Georgia
Athens, Georgia*

December 4, 1969