



Development of Third-stage Physaloptera Larvae from *Crotalus viridis* Rafinesque, 1818 in Cats with Notes on Pathology of the Larvae in the Reptile. (Nematoda, Spiruroidea) *

Author: WIDMER, E. A.

Source: Journal of Wildlife Diseases, 6(2) : 89-93

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.2.89>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Development of Third-stage *Physaloptera* Larvae from *Crotalus viridis* Rafinesque, 1818 in Cats with Notes on Pathology of the Larvae in the Reptile. (Nematoda, Spiruroidea)*

E. A. WIDMER

Department of Environmental Health, School of Public Health,
Loma Linda University, Loma Linda, California 92354.

Abstract

The prairie rattlesnake, *Crotalus viridis*, was found to be commonly infected with third-stage *Physaloptera* larvae. A total of 112 larvae were fed to 3 laboratory raised cats. Adult worms recovered 42 and 59 days postinfection were identified as *P. rara*. Observations were made on the pathology of the larvae in the snake.

Introduction

Considerable descriptive taxonomic information is available for the genus *Physaloptera* but few life cycles have been determined. Partially complete cycles are available for *P. turgida* by Alicata¹ and *P. maxillaris* by Hobmaier.⁵ Complete life cycles have been studied for *P. rara* and *P. praeputialis* by Petri and Ameel⁶ and *P. hispida* by Schell.⁹ These studies utilized orthopteran and coleopteran insects as intermediate hosts.

Natural infections of third-stage physalopterid larvae have been reported from both vertebrate and invertebrate hosts. The earliest report of a natural infection in an invertebrate was the earwig, *Labidura repara*, by Basir.² Zago Filho^{11,12} found the field cricket, *Gryllulus assimilis*, harboring infective *P. praeputialis* larvae and showed that the grasshoppers, *Orphulella punctata*, *Eutryxalis filata filata* and *Dichroplus punctulatus*, were

natural intermediate hosts for *P. praeputialis* and *P. turgida*.

Dixon and Roberson⁴ in their report of *Physaloptera* larvae in the quail, *Colinus virginianus*, reviewed previous reports of infections in birds but omitted the observation of encysted *Physaloptera* larvae in the ruffed grouse, *Bonasa umbellus*, by Boughton.³ The observation of third-stage *Physaloptera* larvae in the stomach of the prairie rattlesnake, *Crotalus viridis*, by Widmer¹⁰ was a new host record.

The presence of third-stage larvae in these vertebrates suggests the possibility of a paratenic host in the life cycle for at least some species of this genus.

The purpose of this paper is to report studies on species identification of the third-stage *Physaloptera* larvae obtained from *C. viridis* and to suggest a paratenic role for this host in the life cycle of *P. rara*.

*Supported under the graduate study leave policy of Loma Linda University, La Sierra, California.

Material and Methods

The snakes used in these experiments were captured in Weld County, Colorado. Techniques in killing and examination were described by Widmer.¹⁰

Microdissection scissors and minuten teasing needles aided in mechanical removal of the larvae. Parasites not used for experimental purposes were killed and fixed in hot 70% alcohol, cleared slowly in glycerine-alcohol and mounted in glycerine.

Sections of the stomach taken from points of larval attachment, were fixed in Bouin's solution and stained in hematoxylin and eosin.

Laboratory raised cats were used in the experiment and controls were estab-

lished by using 2 control animals (#4 and 5). Ether-formalin concentration technique of the feces 30 days before exposure and during the experimental period was routine for all experimental cats. Infections were obtained by force-feeding of the larvae.

Specific infection and necropsy procedures for the 3 experimental cats (#1, 2, and 3) varied as follows: cat #1, fed 16 larvae in a single dose, was killed 7 days after infection; cat #2 was given 40 larvae, 10 initially and 10 at weekly intervals extending over a 3 week period, — necropsy performed 42 days after the initial feeding; cat #3, fed 56 larvae in a single dose, was killed 59 days after exposure.

Results

Host reaction to third-stage Physaloptera larvae.

Mechanical removal of the third-stage *Physaloptera* larvae from the stomach mucosa left small macroscopic lesions. Sections of attachment sites showed a cephalic penetration to the muscularis mucosae (327.6 μ). Lymphocytes, monocytes and plasma cells are located above and below the muscularis mucosae and a concentration of these cells exists near vessels in the submucosa (Fig. 1). Oral glands (Fig. 2) are apparently used in the liquefaction of host tissue prior to ingestion (Figs. 3, 4).

Development in the definitive host.

Third-stage larvae when fed to cats develop to maturity as noted in Table 1.

Mean lengths and widths of worms varied with the duration of infection. Female and male juvenile worms from cat #1 had a mean length of 8.5 and 7.4 mm respectively as compared to 8.0 and 6.2 mm for the third-stage larvae. The sex of these juveniles could be easily determined due to the increase in size of the genital primordia.

In cat #1 the 12 worms were located in the stomach while in cats #2 and 3, 8 and 6 worms respectively were recovered from the duodenum. Control cats #4 and 5 were negative for infection.

For cat #3 eggs of the genus *Physaloptera* were recovered from the feces 35 days after feeding. In cat #2 no eggs were recovered before necropsy, but

TABLE 1. Results of feeding third-stage *Physaloptera* larvae to laboratory raised cats. All measurements are given in mm.

Identification of Host	No. of Larvae Fed	Length of Infection in Days	No. of Worms Recovered		Mean Size			
			Females	Males	Length		Width at Base of Esophagus	
					Females	Males	Females	Males
#1	16	7	7	5	8.5	7.4	.236	.203
#2	40	42	8	3	27.8	23.2	.599	.489
#3	56	59	10	11	33.8	24.0	.808	.613




FIGURE 1. *Third-stage Physaloptera larva attached to stomach mucosa. x 120.* (Photographs shown in Figures 1-4 are of longitudinal sections).



FIGURE 2. *Third-stage Physaloptera larva at edge of tubular mucosal lesion. Note oral glands (arrow) anterior to the nerve ring (N). x 425.*



FIGURE 3. *Third-stage Physaloptera larva at base of tubular lesion. Zone of liquefaction at point of attachment (Z). x 600.*




FIGURE 4. *Tubular lesion, without larva, extending to muscularis mucosae. Zone of liquefaction shows some cellularity. x 425.*

examination of the worms showed mature reproductive organs. Mature eggs were observed in the uteri of all female worms. Additional work needs to be done

to determine a more precise prepatent period.

Mature worms from cats #1 and 2 were identified as *P. rara*.

Discussion

The role of arthropods as intermediate hosts for *Physaloptera turgida*, *P. maxillaris*, *P. rara*, *P. praeputialis*, and *P. hispida* have been reviewed by Lee⁹ and Zago Filho.¹¹ In the original studies no mention was made of a possible paratenic host for any species of this genus. Dixon and Roberson⁷ suggest that *C. virginiana* is not a likely normal intermediate host of *Physaloptera* and consider their findings as a case of an aberrant host.

The high incidence of third-stage larvae in *C. viridis* reported by Widmer¹⁰ does not suggest an aberrant host role for this vertebrate. *P. rara* is a common parasite of cats, dogs, and coyotes.⁷ The apparent ease of infection of cats with third-stage larvae from *C. viridis* suggests the role of a paratenic host of the snake for *P. rara*.

These observations suggest the necessity of further work to determine the role of these larvae in other vertebrate hosts.

Acknowledgments

The author wishes to express his gratitude to Mr. J. Scott Altenbach for collection of the snakes, and to Dr. Leo H. Petri for the identification of the adult worms.

Literature Cited

1. ALICATA, J. E. 1937. Larval development of the spirurid nematode, *Physaloptera turgida*, in the cockroach *Blattella germanica*. Papers on Helm. 30 yr. Jubileum K. I. Skrjabin pp. 11-14.
2. BASIR, M. A. 1948. On a *Physaloptera* larva from an insect. Can. J. Res. D, 26: 197-200.
3. BOUGHTON, R. V. 1935. Endoparasitic infestations in grouse, their pathogenicity and correlation with meteorological conditions. Ph.D. Thesis. Univ. of Minnesota, Minneapolis. 50 p.
4. DIXON, J. M., and J. H. ROBERSON. 1967. Case report: Aberrant larvae of *Physaloptera* sp. in a quail (*Colinus virginianus*). Avian Dis. 11: 41-44.
5. HOBMAIER, M. 1941. Extramammalian phase of *Physaloptera maxillaris* Molin, 1860 (Nematoda). J. Parasit. 27: 233-235.
6. LEE, S. H. 1955. The mode of egg dispersal in *Physaloptera phrynosoma* Ortlepp (Nematoda: Spiruroidea), a gastric nematode of Texas horned toads, *Phrynosoma cornutum*. J. Parasit. 41: 70-74.
7. LEVINE, N. D. 1968. Nematode parasites of domestic animals and of man. Burgess Publishing Co., Minneapolis. 600 p.
8. PETRI, L. H. and D. J. AMEEL. 1950. Studies on the life cycle of *Physaloptera rara* Hall and Wigdor, 1918 and *Physaloptera praeputialis* Linstow 1889. J. Parasit. 36: (6, Suppl.): 40.
9. SCHELL, S. C. 1952. Studies on the life cycle of *Physaloptera hispida* Schell (Nematoda: Spiruroidea) a parasite of the cotton rat (*Sigmodon hispidus littoralis* Chapman). J. Parasit. 38: 462-472.
10. WIDMER, E. A. 1967. Helminth parasites of the prairie rattlesnake, *Crotalus viridis* Rafinesque, 1818, in Weld County, Colorado. J. Parasit. 53: 362-363.
11. ZAGO FILHO, H. 1957. Contribuição para o conhecimento de hospedeiros intermediários e definitivos da *Physaloptera praeputialis* Linstow, 1889 (Nematoda, Spiruroidea). Rev. Brazil Biol. 17: 513-520.
12. ZAGO FILHO, H. 1959. Os gafanhotos como hospedeiros intermediários de *Physaloptera praeputialis* Linst., 1889 e da *Turgida turgida* (Rud. 1819) Trav., 1920 (Nematoda, Spiruroidea). Rev. Brazil Biol. 19: 9-12.