

A Preliminary Report on the Incidence of Lymphocystis Disease in the Fish of the Sapelo Island, Georgia, Area

Author: SMITH, FREDERICK G.

Source: Journal of Wildlife Diseases, 6(4) : 469-471

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.4.469>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A Preliminary Report on the Incidence of Lymphocystis Disease in the Fish of the Sapelo Island, Georgia, Area

FREDERICK G. SMITH

*University of Georgia
Marine Institute
Sapelo Island, Georgia*

Introduction

The observations reported here are a part of a year long survey of diseases of estuarine and near shore fishes being conducted in the Sapelo Island, Georgia, area. This area is located near the middle of the Georgia coast and about five miles to the north of the mouth of the Altamaha River. A variety of diseases and parasitic infections have been found so far, but it appears lymphocystis disease will be one of the most interesting of these problems to study.

Lymphocystis disease is a viral disease that affects both marine and fresh-water fish during the warm months of the year.⁷ It is caused by a DNA virus⁸ that infects

the connective tissue cells of the spleen, ovary, gut wall or the skin of the fish.^{5,7} The principal effect on these fibroblasts is to cause extreme hypertrophy and the secretion of a hyaline membrane around the infected cells.^{1,4,5,8} These greatly enlarged fibroblasts, singly or in tumorous masses, are often macroscopic and constitute the usual means of detecting the disease in infected fish. The virus particles appear in the cytoplasm and escape when the hyaline capsule is ruptured^{4,8} or the fish dies and decomposes.⁷

Much of the information known about this disease is summarized by Weisenberg (1965).⁹

Materials and Methods

Fish for this survey were collected by the use of a ten foot Otter trawl in the Duplin River once a week and by cast net, beach seine and hook-and-line fishing at other locations around the island. All fish were preserved in 10 per cent neutral formalin and given a post mortem examination when returned to the laboratory. An illuminated five inch magnifier

was used for the examination and a dissecting microscope when needed. Tissues for histological preparation were removed from observed lesions. The tissues were embedded in Paraplast[®] and sectioned at eight microns. Haematoxylin - Eosin was used for routine staining.⁸

Results

From Table 1 it can be seen that all cases of lymphocystis disease observed during the first four months of the survey were found in *Stellifer lanceolatus* or

Cynoscion regalis. In October, 1154 fish were examined. Of these, 575 were *S. lanceolatus*. Eight cases of lymphocystis disease were found in *S. lanceolatus*. This

□ Fisher Scientific Co., Pittsburgh, Pennsylvania.

indicates an infection rate of about 1.5%. Two of these cases were visceral infections and the other six were cutaneous infections.

There were fifty *C. regalis* examined in October and one of these had a cutaneous infection of lymphocystis disease; an infection rate of 2%.

In November, 1192 fish were examined. Of these, 793 were *S. lanceolatus*. Three cases of cutaneous lymphocystis disease were found, indicating an infection rate

of about 1.5%. One case of cutaneous lymphocystis disease in the 22 *C. regalis* examined in November indicates an infection rate of about 4.5%.

There were no cases of lymphocystis disease observed in December or January, although 610 and 1252 fish were examined in those months, respectively. In December, 365 *S. lanceolatus* and 6 *C. regalis* were examined. In January 615 and 5 specimens of these fish were examined.

TABLE 1. Incidence of lymphocystis disease near Sapelo Island, Georgia.

Month and Total Fish	Case Number	Genus and Species	Incidence <i>Stellifer</i>	Incidence <i>Cynoscion</i>
October 1154	003	<i>S. lanceolatus</i>	8/575 = 1.4%	1/50 = 2%
	014	<i>S. lanceolatus</i>		
	016	<i>S. lanceolatus</i>		
	030	<i>S. lanceolatus</i>		
	045	<i>S. lanceolatus</i>		
	046	<i>S. lanceolatus</i>		
	057	<i>C. regalis</i>		
	059	<i>S. lanceolatus</i>		
060	<i>S. lanceolatus</i>			
November 1192	090	<i>S. lanceolatus</i>	3/793 = 0.4%	1/22 = 4.6%
	092	<i>S. lanceolatus</i>		
	112	<i>C. regalis</i>		
	119	<i>S. lanceolatus</i>		
December 610			0/364	0/6
January 1252			0/615	0/5

Discussion

Lymphocystis disease has not been reported in the two species of fish which it infects in the Sapelo Island area. Both of these fish are members of the family Sciaenidae and one, *Cynoscion*, is an economically important fish. The cases reported here have all been taken in the Duplin River, a river draining the marshes along Sapelo Island and connecting with Dobby Sound to the south. This does not necessarily mean that the

disease is confined to this one river. It will be necessary to sample more extensively to determine the distribution of infected fish.

From Table 1, it will be noted that the disease was not found in December and January when the water temperature was lower. Enough *Stellifer* and *Cynoscion* were caught during these months to make detection of the disease likely if it was present. The disappearance of the visible

lesions during the colder months has been reported before⁵ and my observations support this. Whether it is water temperature acting along or in conjunction with salinity, photoperiod, dissolved nutrients, planktonic organisms, etc., is yet to be determined. The fish may be more resistant to infection at lower temperatures, or the virus may be less infective.

Templeman⁷ considers lymphocystis disease to be "usually not fatal." I have not found any fish in my samples that appeared to have died of the disease, but several of the heavily infected individuals must have been impaired by the presence of the tumorous masses. This probably increases the susceptibility of these fish to predation.

The lesions found on *Cynoscion* were milder than on *Stellifer* but would have been sufficient to keep the fish out of the food market. The incidence of this disease in market fish should be studied

even though the disease apparently presents no danger to the human consumer.⁷

It has been indicated in the literature that infection with the lymphocystis virus takes place through the gills or by entering the fish through breaks in the skin.⁷ Ectoparasites have been cited as vectors.⁷ There is no indication in this survey what the means of infection in *Stellifer* and *Cynoscion* may be. Both fish are abundant and both fish have been caught with surface abrasions evident. Ectoparasites are found on both fish. *Cynoscion* is a carnivore and *Stellifer* is a filter feeder.² The visceral infections in *Stellifer* may be associated with the ingestion of planktonic organisms carrying the virus or may be the result of viremic transfer.

This summer, studies will be started on the manner of spread of the disease in the two populations of fish in the Sapelo Island area. Additional studies will be started on the species specificity of the virus.

Acknowledgments

This study has been supported by the Department of Anatomy and Histology and the Institute of Comparative Medicine of the College of Veterinary Medicine of the University of Georgia. The staff and facilities of the University of Georgia Marine Institute, Sapelo Island, have made the research possible. University of Georgia Marine Institute Contribution Number 206.

Literature Cited

1. ALEXANDROWICZ, J. S. 1951. Lymphocystis tumors in the red mullet (*Mullus surmuletus* L.). J. Mar. Biol. Assoc. U. K. 30: 315-332.
2. BREDER, C. M. 1948. *Field book of marine fishes of the Atlantic coast from Labrador to Texas*. G. P. Putnam's Sons, New York.
3. LUNA, L. G. (Ed.). 1968. *Manual of histologic staining methods of the Armed Forces Institute of Pathology*. Third edition. McGraw-Hill Book Co., New York.
4. NIGRELLI, R. F., and G. D. RUGGIERI. 1965. Studies on virus diseases of fishes. Spontaneous and experimentally induced cellular hypertrophy (lymphocystis disease) in fishes of the New York Aquarium, with a report of new cases and an annotated bibliography (1874-1965). Zoo. 50: 83-96.
5. NIGRELLI, R. F., and G. M. SMITH. 1939. Studies on lymphocystis disease in the orange filefish, *Ceratacanthus schoepfii* (Walbaum), from Sandy Hook Bay, N. J. Zoo. 24: 255-264.
6. SINDERMAN, C. J. 1970. *Principal diseases of marine fish and shellfish*. Academic Press, Inc., New York.
7. TEMPLEMAN, W. 1965. Lymphocystis disease in American plaice of the eastern Grand Bank. J. Fish. Res. Bd. Canada. 22: 1345-1356.
8. WEISENBERG, R. 1965. Fifty years of research on the lymphocystis virus disease of fishes (1914-1964). Ann. N.Y. Acad. Sci. 126: 362-374.
9. WOLF, K., M. GRAVELL, and R. G. MALSBERGER. 1966. Lymphocystis virus: Isolation and propagation in Centrarchid fish cell lines. Science 151: 1004-1005.