

First Detection of Chronic Wasting Disease in a Wild Red Deer (*Cervus elaphus*) in Europe

Authors: Vikøren, Turid, Våge, Jørn, Madslien, Knut I., Røed, Knut H., Rolandsen, Christer M., et al.

Source: Journal of Wildlife Diseases, 55(4) : 970-972

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/2018-10-262>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First Detection of Chronic Wasting Disease in a Wild Red Deer (*Cervus elaphus*) in Europe

Turid Vikøren,^{1,4} Jørn Våge,¹ Knut I. Madslien,¹ Knut H. Røed,² Christer M. Rolandsen,³ Linh Tran,¹ Petter Hopp,¹ Vebjørn Veiberg,³ Marianne Heum,¹ Torfinn Moldal,¹ Carlos G. das Neves,¹ Kjell Handeland,¹ Bjørnar Ytrehus,³ Øyvør Kolbjørnsen,¹ Helene Wisløff,¹ Randi Terland,¹ Britt Saure,¹ Kine M. Dessen,¹ Solveig Gjerden Svendsen,¹ Brit S. Nordvik,¹ and Sylvie L. Benestad¹ ¹Norwegian Veterinary Institute, PO Box 750 Sentrum, NO-0106 Oslo, Norway; ²Faculty of Veterinary Medicine, Norwegian University of Life Sciences, PO Box 369 Sentrum, N-0102 Oslo, Norway; ³Norwegian Institute for Nature Research, PO Box 5685 Torgarden, NO-7485 Trondheim, Norway; ⁴Corresponding author (email: turid.vikoren@vetinst.no)

ABSTRACT: Chronic wasting disease (CWD) is a fatal contagious prion disease naturally occurring in cervids in North America. In 2016, CWD was detected in wild reindeer (*Rangifer tarandus*) and moose (*Alces alces*) in Norway. Here, we report the first known naturally infected wild Norwegian red deer (*Cervus elaphus*).

Chronic wasting disease (CWD) is a fatal neurodegenerative disorder that affects wild and farmed cervids. It is a prion disease belonging to the transmissible spongiform encephalopathies. The main distribution is North America, and natural infections occur in mule deer (*Odocoileus hemionus*), white-tailed deer (*Odocoileus virginianus*), elk (*Cervus canadensis*), and less frequently in moose (*Alces alces*; Williams 2005; Baeten et al. 2007). Recently, CWD was detected in Europe in wild reindeer (*Rangifer tarandus*) (Benestad et al. 2016) and moose (Pirisinu et al. 2018) from Norway and in one moose from Finland (Finnish Food Safety Authority Evira 2018). In captive red deer (*Cervus elaphus*), natural CWD has been reported in a single animal in Minnesota, US (Schwablander et al. 2013) and recently in a farm in Quebec, Canada (Canadian Food Inspection Agency 2018), respectively. Experimentally, CWD has been transmitted to various cervids, including red deer (Balachandran et al. 2010; Haley and Hoover 2015).

Norway has an estimated winter population of 335,000 wild red deer, moose, roe deer (*Capreolus capreolus*), and reindeer, in total (Austrheim et al. 2011), and approximately 120 deer farms, keeping mainly red deer and some fallow deer (*Dama dama*). The red deer inhabit southern Norway up to 66°N, with the

main population along the west coast, and 42,541 were harvested by hunters in 2017 (Statistics Norway 2018). Here, we report the first case of CWD in a wild red deer worldwide.

An adult female red deer was shot by a hunter in October 2017 in Gjemnes municipality in western Norway (Fig. 1). Before being shot, she was grazing alone on farmland with no signs of disease. She was field dressed, and the head and carcass were delivered to a game slaughterhouse. A spoon sample of the medulla oblongata (MO) was submitted to the Norwegian Veterinary Institute (Oslo, Norway) as part of a national CWD surveillance program. After the initial positive CWD result by routine TeSeE Short Assay Protocol (SAP) enzyme-linked immunosorbent assay (ELISA) (Bio-Rad, Hercules, California, USA) for detection of abnormal prion protein (PrP^{res}), the available remains of the animal were examined at Norwegian Veterinary Institute.

The deer was estimated to be 16 yr old based on cementum annuli in the first incisor root (Hamlin et al. 2000). Necropsy of the head, carcass, and hide revealed good body condition and no gross pathologic lesions. Half of the brain tissue remaining after the spoon sample, half of various lymph nodes (LN), and the tonsils were frozen, and the other halves were fixed in 10% buffered formalin. Samples of brain, LN, and tonsil were processed routinely for immunohistochemical and histologic examination.

The MO and five LNs (retropharyngeal, mandibular, parotid, prescapular, popliteal) were analyzed using TeSeE SAP ELISA. Both the initial sample and second sample of MO at



FIGURE 1. Map of Norway showing the location of wild cervids found positive for chronic wasting disease (CWD) in the period from 2016 through 29 October 2018. The CWD-positive red deer (*Cervus elaphus*) was in Gjemnes municipality (blue filled circle). The CWD-positive moose (*Alces alces*) were in the municipalities of Selbu, 2016 ($n=2$) and Lierne, 2017 ($n=1$; green filled circles). The Nordfjella reindeer area is functionally divided into two zones, and all ($n=19$) the CWD-positive wild reindeer (*Rangifer tarandus*) were found in the northern zone, referred to as zone 1 (red).

the obex area were positive (optical density values 3.5 and 0.330), whereas all LNs were negative. The presence of CWD also was confirmed by the detection of PrP^{res} in MO by using the TeSeE western Blot test (Bio-Rad). Histopathologic examination of MO was inconclusive for spongiform degeneration due to autolysis. No intraneuronal vacuolation was observed.

Sections of MO, LNs, and tonsil were prepared for immunohistochemical examination to demonstrate accumulation of PrP^{res} as in Benestad et al. (2016) using the antibodies mAb L42, F99/97.6.1, and 12B2. In MO, F99/97.6.1 and L42 gave similar staining, with

coarse granular deposits in the neuropil, and perineuronal, intraneuronal, and linear staining in several nuclei and axonal tracts. The immunostaining obtained with 12B2 was weaker and gave no intraneuronal staining. The LNs and tonsils showed no PrP^{res} staining. Genomic DNA was extracted from brain tissue, and the prion protein gene open reading frame was amplified and sequenced (Applied Biosystems 3500×L genetic analyzer with Big Dye Terminator chemistry, Foster City, California, USA), showing glutamic acid (E) homozygosity in codon 226.

Based on brain tissue being found positive for PrP^{res} in three diagnostic tests, we concluded that this represented the first detection of CWD in wild red deer. Elk and red deer are closely related species (*Cervus* spp.). Experimentally, red deer were found to be susceptible to CWD after oral inoculation of prions from elk (Balachandran et al. 2010), accumulating PrP^{res} in brain, lymphoreticular, and other peripheral tissues. Natural CWD was reported in a young captive red deer from Minnesota having PrP^{res} positive brain tissue and LNs (Schwablander et al. 2013). This contrasts with our case, where no detectable PrP^{res} was found in the LNs and tonsils. However, some elk infected with CWD show deposits of PrP^{res} only in the brainstem and not in lymphoid tissues (Spraker et al. 2004, 2015).

Comprehensive surveillance was implemented after detecting CWD in Norway in 2016. Of 59,533 cervids tested from 2016 to 29 October 2018, CWD was diagnosed in 19 reindeer (all from Nordfjella mountain area), three moose (two from Selbu municipality, one from Lierne municipality), and the herein reported red deer from Gjemnes municipality (Fig. 1). In total, 11,398 wild red deer were tested, including 236 from Gjemnes (Norwegian Veterinary Institute 2018).

All of Norway is populated by one or more cervid species with varying degrees of overlapping distribution range. A relevant question is whether there is any migration of cervids between the CWD-positive populations. Gjemnes is located about 200 km north of Nordfjella, 170 km west of Selbu, and 335

km southwest of Lierne (Fig. 1). Seasonal migration is common, and distances may exceed 150 km, but migration distances for red deer in western Norway are usually in the range of 20–40 km (Meisingset et al. 2018).

The origin of CWD in Norwegian wild cervids is unknown. A major question is whether the CWD detected in reindeer, moose, and red deer represents different CWD strains. The CWD from moose in Norway was characterized as an atypical CWD identified as Nor16CWD (Pirisinu et al. 2018). In this context, to better understand the red deer isolate, further molecular analysis and bioassays are currently being done. The prevalence, epidemiology, and implications of the type of CWD discovered in this red deer remain to be determined.

LITERATURE CITED

- Austrheim G, Solberg EJ, Mysterud A. 2011. Spatio-temporal variation in large herbivore pressure in Norway during 1949–1999: Has decreased grazing by livestock been countered by increased browsing by cervids? *Wildl Biol* 17:286–298.
- Baeten LA, Powers BE, Jewell JE, Spraker TR, Miller MW. 2007. A natural case of chronic wasting disease in a free-ranging moose (*Alces alces shirasi*). *J Wildl Dis* 43:309–314.
- Balachandran A, Harrington NP, Algire J, Soutyrine A, Spraker TR, Jeffrey M, González L, O'Rourke KI. 2010. Experimental oral transmission of chronic wasting disease to red deer (*Cervus elaphus elaphus*): Early detection and late stage distribution of protease-resistant prion protein. *Can Vet J* 51:169–178.
- Benestad SL, Mitchell G, Simmons M, Ytrelhus B, Vikøren T. 2016. First case of chronic wasting disease in Europe in a Norwegian free-ranging reindeer. *Vet Res* 47:88.
- Canadian Food Inspection Agency. 2018. *Herds infected with chronic wasting disease in Canada—2018*. <http://www.inspection.gc.ca/animals/terrestrialanimals/diseases/reportable/cwd/herds-infected-in-2018/eng/1517600389905/1517600390422>. Accessed November 2018.
- Finnish Food Safety Authority Evira. 2018. *Moose found dead in forest with chronic wasting disease*. Finnish Food Safety Authority. https://www.evira.fi/en/animals/current_issues/2018/moose-found-dead-in-forest-with-chronic-wasting-disease/. Accessed October 2018.
- Haley NJ, Hoover EA. 2015. Chronic wasting disease of cervids: Current knowledge and future perspectives. *Annu Rev Anim Biosci* 3:305–325.
- Hamlin KL, Pac DF, Sime CA, DeSimone RM, Dusek GL. 2000. Evaluating the accuracy of ages obtained by two methods for Montana ungulates. *J Wildl Manage* 64:441–449.
- Meisingset EL, Loe LE, Brekkum Ø, Bischof R, Rivrud IM, Lande US, Zimmermann B, Veiberg V, Mysterud A. 2018. Spatial mismatch between management units and movement ecology of a partially migratory ungulate. *J Appl Ecol* 55:745–753.
- Norwegian Veterinary Institute. 2018. *CWD surveillance statistics*. <http://apps.vetinst.no/skrantesykestatistikk/NO/#omrade>. Accessed October 2018. [In Norwegian.]
- Pirisinu L, Tran L, Chiappini B, Vanni I, Di Bari MA, Vaccari G, Vikøren T, Madslén KI, Våge J, Spraker J, et al. 2018. Novel type of chronic wasting disease detected in European moose (*Alces alces*), Norway. *Emerg Infect Dis* 24:2210–2218.
- Schwabenlander MD, Culhane MR, Hall SM, Goyal SM, Anderson PL, Carstensen M, Wells SJ, Slade WB, Armien AG. 2013. A case of chronic wasting disease in a captive red deer (*Cervus elaphus*). *J Vet Diagn Invest* 25:573–576.
- Spraker TR, Balachandran A, Zhuang D, O'Rourke KI. 2004. Variable patterns of distribution of PrP(CWD) in the obex and cranial lymphoid tissues of Rocky Mountain elk (*Cervus elaphus nelsoni*) with subclinical chronic wasting disease. *Vet Rec* 155:295–302.
- Spraker TR, Gidlewski T, Powers JG, Nichols T, Balachandran A, Cummings B, Wild MA, VerCauteren K, O'Rourke KI. 2015. Progressive accumulation of the abnormal conformer of the prion protein and spongiform encephalopathy in the obex of nonsymptomatic and symptomatic Rocky Mountain elk (*Cervus elaphus nelsoni*) with chronic wasting disease. *J Vet Diagn Invest* 27:431–441.
- Statistics Norway. 2018. *Red deer hunting*. <https://www.ssb.no/en/jord-skog-jakt-og-fiskeri/statistikker/hjortejakt>. Accessed October 2018.
- Williams ES. 2005. Chronic wasting disease. *Vet Pathol* 42:530–549.

Submitted for publication 29 October 2018.

Accepted 19 December 2018.