

PATHOLOGY AND DISCRETE TYPING UNIT ASSOCIATIONS OF TRYPANOSOMA CRUZI INFECTION IN COYOTES (CANIS LATRANS) AND RACCOONS (PROCYON LOTOR) OF TEXAS, USA

Authors: Hodo, Carolyn L., Bañuelos, Rosa M., Edwards, Erin E., Wozniak, Edward J., and Hamer, Sarah A.

Source: Journal of Wildlife Diseases, 56(1): 134-144

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/2019-03-071

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PATHOLOGY AND DISCRETE TYPING UNIT ASSOCIATIONS OF TRYPANOSOMA CRUZI INFECTION IN COYOTES (CANIS LATRANS) AND RACCOONS (PROCYON LOTOR) OF TEXAS, USA

Carolyn L. Hodo,¹ Rosa M. Bañuelos,² Erin E. Edwards,^{1,3} Edward J. Wozniak,⁴ and Sarah A. Hamer^{2,5}

¹ Department of Veterinary Pathobiology, Texas A&M University, 4467 TAMU, College Station, Texas 77843-4467, USA

² Department of Veterinary Integrative Biosciences, Texas A&M University, 4458 TAMU, College Station, Texas 77843-4458, USA

³ Texas A&M Veterinary Medical Diagnostic Laboratory, PO 3040, College Station, Texas 77831-3040, USA

⁴ Texas Department of State Health Services, PHR1, 6302 Iola Ave, Lubbock, Texas 79424, USA

⁵ Corresponding author (email: shamer@cvm.tamu.edu)

ABSTRACT: Trypanosoma cruzi is a vector-borne, protozoal parasite of mammals. Infected humans, dogs (Canis lupus familiaris), and nonhuman primates may remain asymptomatic or may develop Chagas disease, most commonly characterized by lymphoplasmacytic myocarditis with myocardial degeneration and fibrosis, ultimately resulting in heart failure. Although wildlife species have important roles as sylvatic reservoirs, investigations into the pathology of T. cruzi in wildlife are limited to a few studies documenting histologic lesions in opossums (Didelphis spp.) and raccoons (Procyon lotor). Pathology in covotes (Canis latrans) has not, to our knowledge, been described, despite their recognition as a reservoir and close genetic relationship to domestic dogs. Our objectives were to perform a detailed, comparative cardiac pathology study of sympatric, naturally infected coyotes and raccoons, to characterize the overall *T. cruzi* infection prevalence in the heart and blood of each species via PCR, and to identify infecting discrete typing units (DTUs). We sampled hunter-harvested coyotes (n=120) and raccoons (n=24) in a 28 county region of central and south Texas, US. Raccoons were significantly more likely to have positive PCR results (P < 0.001) with a prevalence of 62% (15/24), comprising DTU TcIV exclusively, with mild to no evidence of cardiac pathology. In contrast, coyotes had a lower infection prevalence (8%, 10/120), comprising DTU TcI exclusively, with lymphoplasmacytic myocarditis observed in four of the six PCR-positive animals. Many raccoons had PCR-positive blood and heart tissue simultaneously, supporting previous reports that raccoons maintain parasitemia into chronic stages of infection; in contrast, none of the PCR-positive covotes were positive in both heart and blood. Our findings demonstrate marked differences in T. cruzi infection dynamics between coyotes and raccoons, with important implications for reservoir potential and their role in transmission cycles.

Key words: American trypanosomiasis, Chagas, coyotes, myocarditis, pathology, raccoons, *Trypanosoma cruzi*.

INTRODUCTION

Trypanosoma cruzi, the vector-borne protozoal agent of Chagas disease, is endemic across much of Latin America and is capable of infecting more than 200 mammalian species (Hoare 1972). The parasite multiplies in the hindgut of triatomine insect vectors (family Reduviidae, subfamily Triatominae), which pass infectious trypomastigotes in their feces. Chagas disease is a major public-health problem in endemic areas and is increasingly recognized as a threat to human and veterinary public health across the southern US, where sylvatic transmission cycles among vectors and wildlife reservoirs have been recognized for decades (Kofoid and McCulloch 1916; Kofoid and Donat 1933). Although many infected hosts may remain asymptomatic, some infected humans, dogs, and nonhuman primates develop cardiac disease, leading to sudden death or congestive heart failure (Rassi et al. 2010; Bern et al. 2011; Snowden and Kjos 2013).

Across the Americas, *T. cruzi* is maintained in complex transmission cycles involving diverse mammalian reservoir species and triatomine vector species. The complexity of those cycles and regional heterogeneity is one of the major challenges in Chagas disease control and prevention. In the US, more than 30 wildlife species have been identified as susceptible hosts, but the relative importance of those species as reservoirs (i.e., their contribution to the transmission and maintenance of the parasite in nature by serving as sources of infection to vectors) has been understudied (Hodo and Hamer 2017). Additionally, investigations into the pathology of T. cruzi in naturally infected wildlife have been conducted on only a limited basis and in only a few species (Packchanian 1942; Ryan et al. 1985; Barr et al. 1991a; Pietrzak and Pung 1998). An understanding of the degree to which various wildlife reservoirs are clinically affected by T. cruzi infection (i.e., their position on the spectrum from unaffected carriers of the parasite to severely diseased hosts) is necessary for predicting populationlevel effects of infection as well as for targeting interventions to manage zoonotic risk.

Trypanosoma cruzi is genotypically heterogeneous and is divided into seven discrete typing units (DTUs): TcI–TcVI and TcBat, which are associated with different geographical regions (Marcili et al. 2009; Zingales et al. 2012), mammalian hosts (Jansen et al. 2017), and vector species (Brenière et al. 2016). Further, there is evidence for associations between DTU and varying clinical outcomes in humans (Ramírez et al. 2010) and dogs (Barr et al. 1991b; Duz et al. 2014) as well as in other experimental animal models (Lisboa et al. 2007; Roellig et al. 2009b). The DTUs TcI and TcIV predominate in the US (Roellig et al. 2008; Bern et al. 2011; Hodo and Hamer 2017), and DTUs TcII, TcIV, TcV, and TcVI were identified in a few rodents in Louisiana (Herrera et al. 2015; Pronovost et al. 2019). In the US, only DTU TcI and an unresolved DTU TcII/V/VI group have been associated with autochthonous human infection (Roellig et al. 2013; Garcia et al. 2017). Regarding vertebrate host associations in the US, most sampled raccoons (Procyon lotor) were infected with DTU TcIV, whereas opossums (*Didelphis* spp.) were almost exclusively infected with DTU TcI (Roellig et al. 2008; Bern et al. 2011; Hodo and Hamer 2017).

Both DTUs TcI and TcIV have been identified in skunks (*Mephitis mephitis*), armadillos (*Dasypus novemcinctus*), woodrats (*Neotoma* sp.), and domestic dogs (*Canis lupus familiaris*; Roellig et al. 2008; Charles et al. 2013; Curtis-Robles et al. 2017a; Hodo and Hamer 2017). These DTU associations may be important in classifying wildlife species as important reservoirs of infection. Additionally, associations between strain type and pathology or clinical outcome in wildlife may be translatable for human and domestic animal health.

Raccoons are perhaps the best-studied reservoir species in the US, and T. cruzi infection dynamics and pathology are described in both naturally infected and experimentally infected animals (Pietrzak and Pung 1998; Roellig et al. 2009b; Bern et al. 2011). In contrast, coyotes (Canis latrans) are a relatively understudied host species, for which most previous studies report seroprevalence only (Hodo and Hamer 2017) and for which, to our knowledge, only a single, limited pathology study exists (Curtis-Robles et al. 2016). Raccoons and coyotes are abundant in both rural and urban settings, having the potential to bridge parasite infections from sylvatic to peridomestic habitats. Our objectives were to perform a detailed comparative cardiac-pathology study of naturally infected coyotes in central and south Texas and sympatric raccoons in central Texas and to characterize the overall T. cruzi infection prevalence in the heart and blood of each species via PCR with identification of infecting DTU.

MATERIALS AND METHODS

Sampling

We conducted a cross-sectional sampling effort in January 2016 at an annual recreational nuisance-animal hunt organized by private landowners in central Texas. Animals from that hunt have been studied in prior years in the context of *T. cruzi* infection, such that we anticipated finding infected animals (Curtis-Robles et al. 2016). Animals legally harvested under recreational permits by teams of hunters during a 24-h period were brought to a central check station. The area



FIGURE 1. Sampling locations of (A) coyotes (*Canis latrans*) and (B) raccoons (*Procyon lotor*) by county in Texas, USA, during January and March 2016. Hearts and blood of these hunter-harvested animals were tested via PCR for presence of *Trypanosoma cruzi*. Triangles mark counties from which *T. cruzi* PCR-positive animals were collected.

of harvest encompassed 25 counties in central Texas (Fig. 1) spanning five different ecoregions (Gould et al. 1960). Our team collected samples from coyotes and raccoons for which the county of harvest was known. Hearts and blood (when available) were collected in the field within 24 h of death. At the time of collection, each heart was briefly examined for gross lesions. Blood was collected from the axillary vasculature or the thoracic cavity, as available, during heart collection. Blood and hearts were transported to the laboratory on ice.

In the laboratory, blood samples were centrifuged, and serum was collected from those samples without extensive hemolysis. A 500- μ L volume of a blood clot or whole, hemolyzed blood was subsampled from each blood sample and stored at -20 C until DNA extraction. Hearts were stored at -80 C for 4–5 mo, then thawed and dissected to examine right and left atria and ventricles. Two tissue samples were taken from each of the four chambers. One section from each chamber was stored in 10% neutral-buffered formalin for histology, and the other section was minced. Minced samples from all chambers for each animal were pooled together for DNA extraction.

We also received samples from coyotes collected during an oral rabies-vaccine program surveillance in Webb County (south Texas) by the Texas Department of State Health Services in coordination with US Department of Agriculture– Animal and Plant Health Inspection, Wildlife Services, Texas branch, from 29 February 2016 to 1 March 2016 (Fig. 1). Heart apex, blood-soaked Nobuto filter paper (Advantec MFS, Dublin, California, USA), and whole blood (when available) were collected by Department of State Health Services personnel in the field. A section of the heart apex was subsampled and minced for DNA extraction.

Molecular work

From each animal, as available, one approximately 0.5-cm³ sample from each heart (representing all four chambers from central Texas animals and the apex only from south Texas animals), 500 μ L of blood, and one Nobuto strip were subjected to DNA extraction with the Omega[®] E.Z.N.A.[®] Tissue Extraction Kit (Omega Bio-tek, Norcross, Georgia, USA) according to the manufacturer's protocol for tissue extraction with an overnight lysis for the hearts and Nobuto strips and \geq 3 h lysis for blood.

After DNA extraction, samples were tested for infection with *T. cruzi* with a highly sensitive quantitative PCR to amplify a 166-base pair fragment of *T. cruzi* satellite DNA, as previously described (Piron et al. 2007; Curtis-Robles et al. 2016). DNA-negative controls (water) and a positive control of DNA extracted from a pure culture of Sylvio X10 CL4 (ATCC 50800, ATCC, Manassas, Virginia, USA; DTU TcI) were included in all reactions. Samples with a cycle threshold (Ct) value less than 36 were considered suspect positive and subjected to a multiplex quantitative PCR targeting the spliced leader intergenic region to confirm positivity and for determination of DTU, according to previously described protocols (Cura et al. 2015; Curtis-Robles et al. 2017a). The criterion for considering a sample positive on this assay was the detection of specific fluorescence to one or more DTU-specific probes within 40 cycles in a 45-cycle assay. Negative controls (water) and positive controls of DNA extracted from T. cruzi strain Sylvio X10 CL4 (DTU TcI, details presented earlier) and T. cruzi-infected Triatoma sanguisuga from Texas (DTU TcIV) were included in all reactions, and the positive control of DNA extracted from T. cruzi Y-strain (ATCC 50832, ATCC; DTU TcII) was added for reactions run later in the study. Samples were considered positive if they generated Ct values below the respective thresholds on both PCR assays, and animals were considered PCR positive if either blood or heart samples were positive.

Histopathology

Formalin-fixed heart tissue from central Texas coyotes and raccoons was processed routinely for histopathology and stained with H&E. Two slides from each animal, representing right and left atria and ventricles, were examined by light microscopy by a board-certified veterinary pathologist, blinded to the PCR status. Inflammation was semiquantitatively scored for each heart chamber on a numeric scale as normal (0), minimal (1), mild (2), moderate (3), or severe/marked (4). Additionally, the presence of fibrosis, cardiomyocyte degeneration, or necrosis and the distribution (focal, multifocal, focally extensive) and location (interstitial, myocardial, epicardial) of lesions were recorded. An inflammation index for each animal was calculated by adding the inflammation scores for each chamber. For analysis, animals were dichotomized by pathology status (significant lesions present or absent), in which significant was defined as an inflammation score ≥ 3 . The inflammation cutoff of 3 was chosen because it represented at least minimal inflammation in three heart chambers, at least mild inflammation in one section and minimal in another, or at least moderate inflammation in any one heart chamber. Slides from animals with an inflammation score \geq 3 were reexamined, and a descriptive morphologic diagnosis was recorded. Animals with lymphoplasmacytic myocarditis (consistent with expected lesions of T. cruzi infection) were included in the statistical analyses.

Statistical analysis

We tested for significant differences between the presence of T. cruzi DNA in samples (PCR status of heart and blood) and host attributes of species and sex using a Fisher's exact test. Further, we used the Fisher's exact text to compare the presence of *T. cruzi* DNA in samples with the presence of lymphoplasmacytic myocarditis for each species separately. Finally, the Mann-Whitney-Wilcoxon test was used to determine whether the inflammation scores differed between *T. cruzi*-positive coyotes and raccoons and whether the Ct values of PCR-positive blood samples were different between species. Statistical analyses were performed in R software (R Core Team 2018).

RESULTS

Sample population

We sampled 120 coyotes from 24 Texas counties and 24 raccoons from 14 counties (Fig. 1). Both males and females of each species were sampled. We collected hearts from 97 central Texas coyotes and 23 raccoons and heart apex from 23 south Texas coyotes. Blood was available for 92 coyotes and 18 raccoons from central Texas and 21 coyotes from south Texas. We also received bloodsoaked Nobuto filter paper strips from all 23 south-Texas coyotes.

PCR results

A total of 8% (10/120) of coyotes and 62% (15/24) of raccoons were confirmed to have positive PCR results for T. cruzi on two separate quantitative PCRs of either heart or blood (Table 1). Raccoons were significantly more likely to be *T. cruzi*-positive than were coyotes (P < 0.001). Sex was not associated with PCR status within either species. One coyote with PCR-positive blood was based on testing of a Nobuto strip (whole blood was not available for that animal). All other Nobuto strips tested had negative results, including three from coyotes with PCR-positive whole blood. The Ct values in T. cruzi-positive blood were significantly less for raccoons (mean=24.5, median=25.2, range, 18.7–27.4) than coyotes (mean=32.6, median=33.2, range, 31.1–33.86; P=0.002); thus, infected raccoons had a higher concentration of parasite DNA in blood than did coyotes did. All infected coyotes harbored DTU TcI,

| | | | Samples positive by PCR for <i>Trypanosoma cruzi</i> ^a | | | | | | | |
|----------|-----|-------|---|-------|-----------|-------|-----------|--|--|--|
| | | Heart | | Blood | | Total | | | | |
| Species | No. | % | Frequency | % | Frequency | % | Frequency | | | |
| Coyotes | 120 | 4 | 5/120 | 4 | 5/113 | 8 | 10/120 | | | |
| Raccoons | 24 | 52 | 12/23 | 44 | 8/18 | 62 | 15/24 | | | |

TABLE 1. Results of *Trypanosoma cruzi* PCR testing of heart and blood from hunter-harvested coyotes (*Canis latrans*) and raccoons (*Procyon lotor*) from central and south Texas, USA, collected in January and March 2016.

^a Frequency = no. positive/no. tested.

whereas all infected raccoons harbored DTU TcIV. No individual animals were found to harbor more than one DTU. Positive raccoons were identified in 13 of 14 sampled counties, and positive coyotes originated from five of 25 sampled counties (Fig. 1).

Pathology

Significant gross lesions, other than gunshot wound-associated trauma, were not observed in any of the hearts. Incidentally, adult heartworms (Dirofilaria immitis) were observed in six coyotes. Histologically, 62 of 120 animals had no lesions, 41 of 120 had minimal findings that were not considered significant (inflammation score <3), and 15 of 120 animals (11 coyotes and four raccoons) had lesions considered to be significant (inflammation score >3) for further characterization and inclusion in the significant-lesions group for analysis. One coyote was excluded from the histopathology analysis because of severe autolysis across all tissue sections examined. Four other coyotes that had severe autolysis of all sections, except for the left ventricle, were retained in the analysis and had no significant lesions in the assessed region of left ventricle.

Of the 11 coyotes with significant lesions, six (including four *T. cruzi*-positive animals) had mild to moderate, multifocal, lymphoplasmacytic myocarditis with varying degrees of myocardial degeneration and fibrosis (Fig. 2A–C), consistent with that described for *T. cruzi* infection in other species (Barr et al. 1991c; Andrade et al. 2009; Pisharath et al. 2013; Snowden and Kjos 2013). Another four of the 11 coyotes with significant lesions had severe locally extensive inflammation that was primarily histiocytic or pyogranulomatous, occasionally with visible intraleukocytic zoites, most consistent with *Hepatozoon americanum* infection (Davis et al. 1978). None of these animals had positive PCR results for *T. cruzi*. Mature cysts of *H. americanum* were observed in two of these animals, as well as in other coyotes without inflammation. Another coyote had mild, multifocal pyogranulomatous inflammation associated with microfilariae of *D. immitis*. Cysts of *Sarcocystis* sp. were observed in two coyote hearts, with no accompanying inflammation. Two of the six *T. cruzi*–positive coyotes had no apparent lesions in the sections examined.

Of the five raccoons with significant histologic lesions (including three T. cruzi-positive animals), three (two of which were T. cruzi positive) had minimal to mild lymphoplasmacytic myocarditis (Fig. 2D) considered possibly consistent with T. cruzi infection, which was accompanied by fibrosis in one animal. In a fourth, T. cruzi-positive raccoon, lymphoplasmacytic inflammation was accompanied by abundant eosinophils, which is not typically associated with T. cruzi. The fifth, T. cruzinegative raccoon with lesions had moderateto-severe multifocal histiocytic and lymphoplasmacytic perivascular myocarditis and epicarditis. A Sarcocystis sp. cyst was also observed in that animal but not in the area of inflammation. Trypanosoma cruzi amastigotes were not observed in any of the sections examined for either species. A total of 80% (12/15) of the T. cruzi-positive raccoons had no significant inflammation in the heart sections examined.



FIGURE 2. Photomicrographs showing myocarditis in coyotes (*Canis latrans*) and raccoons (*Procyon lotor*) collected from central Texas, USA in 2016, H&E stain. (A) Scale bar=500 μm, coyote C16-38, left ventricle; inflammation is present in multiple areas across the section. (B) Scale bar=50 μm, higher-magnification view of (A); cardiac myofibers are separated by a moderate amount of inflammation composed primarily of lymphocytes and plasma cells, with myocellular degeneration and loss. (C) Scale bar=100 μm, right ventricle, coyote C16-75; moderate lymphoplasmacytic inflammation is perivascular and infiltrates between cardiac myofibers, disrupting normal architecture. (D) Scale bar=50 μm, raccoon R16-23, left ventricle; mild lymphoplasmacytic myocarditis.

Table 2 shows the number of animals with lymphoplasmacytic myocarditis compared with *T. cruzi* status. The presence of lymphoplasmacytic myocarditis was significantly associated with *T. cruzi* positivity in coyotes (P<0.001) but not in raccoons (P=0.713). Additionally, the severity of lymphoplasmacytic myocarditis (as measured by the combined inflammation score) was greater for *T. cruzi*-positive coyotes than for negative coyotes (P<0.001) when other types of inflammation (attributed to other etiologic agents) were excluded.

DISCUSSION

In a region of central and south Texas known as a hotspot for *T. cruzi* transmission

(Curtis-Robles et al. 2016, 2017b, 2018), both raccoons and coyotes had T. cruzi-positive blood and cardiac tissue, sometimes associated with myocarditis. Thus, these species may not only be involved in the sylvatic transmission cycle, but also be negatively affected by the infection in some cases, especially coyotes. Although the overall *T. cruzi* prevalence was significantly higher in raccoons (62%; 15/24)than in coyotes (8%; 10/120), the infected coyotes exhibited more-severe histologic lesions than did infected raccoons. Finally, we found an association between parasite DTU and host taxa; all 10 coyotes for which DTU was determined harbored DTU TcI, whereas all 15 raccoons harbored DTU TcIV.

| | | | Presence of lesions ^a | | | | | |
|----------|-------------|-----|----------------------------------|-----------|--|-----------|--|--|
| | | | Lymphoplasmacytic myocarditis | | No myocarditis, or with other etiology | | | |
| Species | PCR results | No. | % | Frequency | % | Frequency | | |
| Coyotes | Positive | 6 | 67 | 4/6 | 33 | 2/6 | | |
| | Negative | 90 | 2 | 2/90 | 98 | 88/90 | | |
| | Total | 96 | 6 | 6/96 | 94 | 90/96 | | |
| Raccoons | Positive | 15 | 20 | 3/15 | 80 | 12/15 | | |
| | Negative | 8 | 25 | 2/8 | 75 | 6/8 | | |
| | Total | 23 | 22 | 5/23 | 78 | 18/23 | | |

TABLE 2. Two-by-two comparison of *Trypanosoma cruzi* PCR status and presence or absence of lymphoplasmacytic myocarditis in hunter-harvested coyotes (*Canis latrans*) and raccoons (*Procyon lotor*) collected from central Texas, USA, in January and March 2016.

^a Frequency = no. positive/no. tested.

The infection prevalence we found was consistent with previous reports in both raccoons and coyotes, where prevalences range from 20% to 90% in raccoons across the southern states (Hodo and Hamer 2017), and from 4% to 14% in coyotes in several southern states (Hodo and Hamer 2017). Specifically, a study that sampled animals during a previous year of the recreational hunt across many of the same central Texas counties reported prevalences of 70.0% in raccoons and 14.3% in coyotes (Curtis-Robles et al. 2016). The likely route of transmission in these animals is oral, through ingestion of infected kissing bug vectors. This is considered an important route in dogs (Barr 2009) and was effective in experimentally infecting raccoons and skunks (Davis et al. 1980; Roellig et al. 2009a). Transmission may also occur through ingestion of parasitemic vertebrates (Thomas et al. 2007; Rocha et al. 2013), although ingestion of infected meat did not infect raccoons (Roellig et al. 2009a).

Only minimal to mild lesions have been reported from natural infection with *T. cruzi* in raccoons (Pietrzak and Pung 1998; Charles et al. 2013; Curtis-Robles et al. 2016), consistent with our findings. Experimentally infected raccoons exhibited mild to severe lesions that varied with acuteness of infection and infecting DTU, exhibiting more-severe cardiac lesions during the acute stages of infection with DTUs TcI and TcII than with DTU TcIV (Roellig et al. 2009b). The specific lesions described in previous studies are very similar to those we observed: inflammation composed primarily of lymphocytes and plasma cells, with occasional myocardial degeneration or necrosis and infrequent observations of intramyocellular T. cruzi amastigotes. Significant histopathologic lesions in T. cruzi-infected coyotes have not previously been described, but our findings of multifocal lymphoplasmacytic inflammation with destruction of cardiomyocytes were consistent with lesions reported in infected dogs (Barr et al. 1991c; Snowden and Kjos 2013). A limitation of our study was that severely affected animals with advanced heart disease, including animals that died, are not represented in our study population because they would not be available for hunters to harvest.

Autolysis hindered interpretation of some histologic sections, but less than 5% of examined sections were affected to the extent that we could not determine the presence or absence of inflammation. Overall, only minimal artifactual change resulted from the single freeze-thaw cycle at -80 C, which was manifested mainly by artifactual separation of myofibers. The relative lack of freeze-thaw artifact observed should be considered in future sampling efforts for which histopathology may have otherwise been discounted because logistics would require freezing of samples.

Definitive diagnosis of *T. cruzi* based solely on histopathology is often difficult because the intracellular amastigote form is not commonly observed in chronic infections. Indeed, in our study, we did not observe any amastigotes in histologic sections. However, although not pathognomonic, the multifocal and infiltrative pattern of lymphoplasmacytic myocarditis is supportive of T. cruzi infection, especially when accompanied by positive PCR or serologic results. Studies in nonhuman primates identified an association between nonspecific lymphocytic myocarditis and the presence of T. cruzi DNA (Andrade et al. 2009; Mubiru et al. 2014). In our study, lymphoplasmacytic inflammation was associated with T. cruzi-positive PCR status in coyotes but not in raccoons. Inflammation was observed in the hearts of several PCRnegative coyotes and, in some cases, was explained by the presence of other parasites, including H. americanum and D. immitis and was characterized by the presence of macrophages and neutrophils. Other differentials for lymphoplasmacytic myocarditis in these animals included Bartonella sp. (Chomel et al. 2006) or Borrelia burgdorferi (Janus et al. 2014), although *Bartonella* sp. is most often associated with valvular endocarditis (Pesavento et al. 2005), and B. burgdorferi is uncommon in Texas (Bowman et al. 2009; Mitchell et al. 2016; Hodo et al. 2019).

Additionally, T. cruzi-negative status in the face of lesions suggestive of *T. cruzi* could also be explained by false-negative PCR results. Testing results from a single blood sample or small pieces of heart tissue do not necessarily reflect the true infection status of the individual. Thus, false-negative results could have resulted from sampling error because of the multifocal nature of the parasite distribution. This was supported by our histologic findings in which inflammation was focal to multifocal and not diffusely distributed throughout all sections. Other raccoon studies have reported PCR-positive results in some, but not all, sections of heart from the same animal (James et al. 2002; Curtis-Robles et al. 2016).

Eight of 11 T. cruzi-infected raccoons had PCR-positive blood, and all but one of those also had positive heart tissue (Table 1), suggesting chronic infection. Although PCR cannot confirm the presence of whole, viable parasites, the presence of parasite DNA in blood is suggestive of parasitemia and likely infectiousness to vectors. In contrast, we only detected T. cruzi DNA in the blood of five of 10 infected covotes, and none of these had PCR-positive heart tissue (Table 1). This suggested the possibility that, although raccoons appear to maintain long-term parasitemia with DTU TcIV, coyotes may only circulate DTU TcI during the acute stage of infection before the parasite localizes in tissues.

We found exclusive associations between host taxa and parasite DTU, in which raccoons were only infected with DTU TcIV and coyotes only with DTU TcI. The difference in degree of pathology observed between coyotes and raccoons could be explained by host specieslevel or DTU-level differences. In domestic dogs, experimental evidence supports differences in pathology related to different parasite strains. For example, dogs infected with T. cruzi isolates from an armadillo and opossum developed acute and chronic myocarditis, whereas dogs infected with an isolate from a dog did not develop disease (Barr et al. 1991c). In another study, DTU TcI infection resulted in more inflammatory cells in the hearts of infected dogs compared with DTU TcII infections (Duz et al. 2014). Raccoons are almost exclusively infected with DTU TcIV across multiple studies, with few reports of natural DTU TcI infection (Roellig et al. 2008; Bern et al. 2011; Curtis-Robles et al. 2016). Experimentally, raccoons were successfully infected with DTUs TcIV, TcI, and TcII, but developed longer-lasting parasitemia with DTU TcIV and more-severe cardiac lesions with DTUs TcI and TcII infections (Roellig et al. 2009b). Trypanosoma cruzi DTU TeIV may be host adapted to raccoons, supported by their high infection prevalence and lack of obvious pathology (Roellig et al. 2009b). No such association has been suggested for coyotes, and our findings of DTU TcI infection

with cardiac pathology may be evidence that DTU TcI is fundamentally a more cardiopathogenic strain than is DTU TcIV or simply that coyotes are more susceptible than raccoons to cardiac disease resulting from T. cruzi infection. In domestic dogs in the US, both DTU TcI and TcIV infections have been reported (Roellig et al. 2008; Hodo et al. 2019). Concerning other wildlife in central and south Texas, opossums harbor DTU TcI (Hodo et al. 2018), whereas both DTUs TcI and TcIV are found in skunks (Hodo et al. 2018) and woodrats (Charles et al. 2013). Interestingly, DTU TcIV has not been detected in autochthonous human infections of *T. cruzi* in the US (Roellig et al. 2008; Garcia et al. 2017), despite its widespread presence in triatomine vectors (Curtis-Robles et al. 2018). More research is needed on the comparative pathology of the different DTUs in canines as well as in humans. Although part of the response to T. cruzi infection is likely dependent on the individual host, evidence is accumulating that DTU may also have a role.

In conclusion, although coyotes had a lower T. cruzi infection prevalence than raccoons did, infection in covotes was associated with more-severe lesions and with DTU TcI. These findings may have important implications for the association of T. cruzi DTU with resulting pathology as well as for the reservoir potential of coyotes and raccoons. We also provide further support for the association of DTU TcIV with raccoons in the US. Although raccoons are known to maintain high levels of parasitemia into the chronic stages of infection (Roellig et al. 2009b), further investigation of parasitemia dynamics of coyotes is needed to determine their contribution to the reservoir community for T. cruzi. Additional considerations necessitating further research are the risks posed to hunters exposed to infectious wildlife as well as the effect of infection on wildlife populations.

ACKNOWLEDGMENTS

We thank the following individuals for help with field or laboratory work: Lisa Auckland, Rachel Curtis-Robles, Martha Hensel, Chloe

Goodwin, Jessica Rodriguez, Karen Snowden, Alyssa Meyers, Italo Zecca, Kimberley Doll, Amani Bourji, Beverley Finneburgh, Adam Curtis, and Elise Birkner. We acknowledge the hunters who donated animals to this study and thank Bob Woodward for his coordination and assistance. We are grateful to Justin Henefey and the US Department of Agriculture-Animal and Plant Health Inspection Wildlife Services personnel who were involved in collection of coyote samples in south Texas. Funding support was provided by the Department of Veterinary Integrative Biosciences and AgriLife Research. Student stipend support was provided by the National Institutes of Health (NIH) grants 2T32OD011083-06 to C.L.H. and NIH grant T35 2T35OD010991 to R.M.B.

LITERATURE CITED

- Andrade MCR, Dick EJ Jr, Guardado-Mendoza R, Hohmann ML, Mejido DCP, VandeBerg JL, DiCarlo CD, Hubbard GB. 2009. Nonspecific lymphocytic myocarditis in baboons is associated with *Trypanosoma cruzi* infection. Am J Trop Med Hyg 81:235– 239.
- Barr SC, Brown CC, Dennis VA, Klei TR. 1991a. The lesions and prevalence of *Trypanosoma cruzi* in opossums and armadillos from southern Louisiana. J *Parasitol* 77:624–627.
- Barr SC, Gossett KA, Klei TR. 1991b. Clinical, clinicopathologic, and parasitologic observations of trypanosomiasis in dogs infected with North American *Trypanosoma cruzi* isolates. Am J Vet Res 52:954– 960.
- Barr SC, Schmidt SP, Brown CC, Klei TR. 1991c. Pathologic features of dogs inoculated with North American *Trypanosoma cruzi* isolates. Am J Vet Res 52:2033–2039.
- Barr SC. 2009. Canine Chagas' disease (American trypanosomiasis) in North America. Vet Clin North Am Small Anim Pract 39:1055–1064.
- Bern C, Kjos S, Yabsley MJ, Montgomery SP. 2011. *Trypanosoma cruzi* and Chagas' disease in the United States. *Clin Microbiol Rev* 24:655–681.
- Bowman D, Little SE, Lorentzen L, Shields J, Sullivan MP, Carlin EP. 2009. Prevalence and geographic distribution of *Dirofilaria immitis*, *Borrelia burgdorferi*, *Ehrlichia canis*, and *Anaplasma phagocytophilum* in dogs in the United States: Results of a national clinic-based serologic survey. *Vet Parasitol* 160:138– 148.
- Brenière SF, Waleckx E, Barnabé C. 2016. Over six thousand *Trypanosoma cruzi* strains classified into discrete typing units (DTUs): Attempt at an inventory. *PLoS Negl Trop Dis* 10:e0004792.
- Charles RA, Kjos S, Ellis AE, Barnes JC, Yabsley MJ. 2013. Southern plains woodrats (*Neotoma micropus*) from southern Texas are important reservoirs of two genotypes of *Trypanosoma cruzi* and host of a

putative novel *Trypanosoma* species. Vector Borne Zoonotic Dis 13:22–30.

- Chomel BB, Boulouis H-J, Maruyama S, Breitschwerdt EB. 2006. Bartonella spp. in pets and effect on human health. Emerg Infect Dis 12:389–394.
- Cura CI, Duffy T, Lucero RH, Bisio M, Péneau J, Jimenez-Coello M, Calabuig E, Gimenez MJ, Ayala EV, Kjos SA, et al. 2015. Multiplex real-time PCR assay using TaqMan probes for the identification of *Trypanosoma cruzi* DTUs in biological and clinical samples. *PLoS Neglect Trop Dis* 9:e0003765.
- Curtis-Robles R, Auckland LD, Snowden KF, Hamer GL, Hamer SA. 2018. Analysis of over 1500 triatomine vectors from across the US, predominantly Texas, for *Trypanosoma cruzi* infection and discrete typing units. *Infect Genet Evol* 58:171–180.
- Curtis-Robles R, Lewis BC, Hamer SA. 2016. High *Trypanosoma cruzi* infection prevalence associated with minimal cardiac pathology among wild carnivores in central Texas. *Int J Parasitol Parasite Wildl* 5: 117–123.
- Curtis-Robles R, Snowden KF, Dominguez B, Dinges L, Rodgers S, Mays G, Hamer SA. 2017a. Epidemiology and molecular typing of *Trypanosoma cruzi* in naturally-infected hound dogs and associated triatomine vectors in Texas, USA. *PLoS Negl Trop Dis* 11: e0005298.
- Curtis-Robles R, Zecca IB, Roman-Cruz V, Carbajal ES, Auckland LD, Flores I, Millard AV, Hamer SA. 2017b. *Trypanosoma cruzi* (agent of Chagas disease) in sympatric human and dog populations in "colonias" of the Lower Rio Grande Valley of Texas. *Am J Trop Med Hyg* 96:805–814.
- Davis DS, Robinson RM, Craig TM. 1978. Naturally occurring hepatozoonosis in a coyote. J Wildl Dis 14: 244–246.
- Davis DS, Russell LH, Adams LG, Yaeger RG. 1980. An experimental infection of *Trypanosoma cruzi* in striped skunks (*Mephitis mephitis*). J Wildl Dis 16: 403–406.
- Duz ALC, Vieira PM de A, Roatt BM, Aguiar-Soares RDO, Cardoso JM de O, de Oliveira FCB, Reis LES, Tafuri WL, Veloso VM, Reis AB, et al. 2014. The TcI and TcII *Trypanosoma cruzi* experimental infections induce distinct immune responses and cardiac fibrosis in dogs. *Mem Inst Oswaldo Cruz* 109:1005– 1013.
- Garcia MN, Burroughs H, Gorchakov R, Gunter SM, Dumonteil E, Murray KO, Herrera CP. 2017. Molecular identification and genotyping of *Trypano-soma cruzi* DNA in autochthonous Chagas disease patients from Texas, USA. *Infect Genet Evol* 49:151– 156.
- Gould FW, Hoffman GO, Rechenthin CA. 1960. Vegetational areas of Texas. *Texas Agriculture Experimental Station leaflet* 492. Texas A&M University, College Station, Texas, 4 pp.
- Herrera CP, Licon MH, Nation CS, Jameson SB, Wesson DM. 2015. Genotype diversity of *Trypanosoma cruzi* in small rodents and *Triatoma sanguisuga* from a

rural area in New Orleans, Louisiana. *Parasit Vectors* 8:123.

- Hoare CA. 1972. The trypanosomes of mammals: A zoological monograph. Blackwell Scientific Publications, Oxford, UK, 749 pp.
- Hodo CL, Hamer SA. 2017. Toward an ecological framework for assessing reservoirs of vector-borne pathogens: wildlife reservoirs of *Trypanosoma cruzi* across the southern United States. *ILAR J* 58:379– 392.
- Hodo CL, Rodriguez JY, Curtis-Robles R, Zecca IB, Snowden KF, Cummings KJ, Hamer SA. 2019. Repeated cross-sectional study of *Trypanosoma cruzi* in shelter dogs in Texas, in the context of *Dirofilaria immitis* and tick-borne pathogen prevalence. J Vet Int Med 33:158–166.
- Hodo CL, Wilkerson GK, Birkner EC, Gray SB, Hamer SA. 2018. *Trypanosoma cruzi* transmission among captive nonhuman primates, wildlife, and vectors. *Ecohealth* 15:426–436.
- James MJ, Yabsley MJ, Pung OJ, Grijalva MJ. 2002. Amplification of *Trypanosoma cruzi-specific DNA* sequences in formalin-fixed raccoon tissues using polymerase chain reaction. J Parasitol 88:989–993.
- Jansen AM, Xavier SCC, Roque ALR. 2017. Ecological aspects of *Trypanosoma cruzi*: wild hosts and reservoirs. In: *American trypanosomiasis Chagas disease*, *One hundred years of research*, 2nd Ed., Telleria J, Tibayrenc M, editors. Elsevier, Amsterdam, the Netherlands, pp. 243–264.
- Janus I, Noszczyk-Nowak A, Nowak M, Cepiel A, Ciaputa R, Pasławska U, Dzięgiel P, Jabłońska K. 2014. Myocarditis in dogs: etiology, clinical and histopathological features (11 cases: 2007–2013). Ir Vet J 67: 28.
- Kofoid CA, Donat F. 1933. Experimental infection with *Trypanosoma cruzi* from intestine of cone-nose bug, *Triatoma protracta*. Proc Soc Exp Biol Med 30:489–491.
- Kofoid CA, McCulloch I. 1916. On Trypanosoma triatomae, a new flagellate from a Hemipteran bug from the nests of the wood rat Neotoma fuscipes. Univ Calif Publ Zool 16:113–126.
- Lisboa CV, Pinho AP, Monteiro RV, Jansen AM. 2007. *Trypanosoma cruzi* (kinetoplastida Trypanosomatidae): biological heterogeneity in the isolates derived from wild hosts. *Exp Parasitol* 116:150–155.
- Marcili A, Lima L, Cavazzana M, Junqueira ACV, Veludo HH, da Silva FM, Campaner M, Paiva F, Nunes VLB, Teixeira MMG. 2009. A new genotype of *Trypanosoma cruzi* associated with bats evidenced by phylogenetic analyses using SSU rDNA, cytochrome b and Histone H2B genes and genotyping based on ITS1 rDNA. *Parasitology* 136:641–645.
- Mitchell EA, Williamson PC, Billingsley PM, Seals JP, Ferguson EE, Allen MS. 2016. Frequency and distribution of Rickettsiae, Borreliae, and Ehrlichiae detected in human-parasitizing ticks, Texas, USA. *Emerg Infect Dis* 22:312–315.
- Mubiru JN, Yang A, Dick EJ Jr, Owston M, Sharp RM, VandeBerg JF, Shade RE, VandeBerg JL. 2014.

Correlation between presence of *Trypanosoma cruzi* DNA in heart tissue of baboons and cynomolgus monkeys, and lymphocytic myocarditis. *Am J Trop Med Hyg* 90:627–633.

- Packchanian A. 1942. Reservoir hosts of Chagas' disease in the state of Texas: Natural infection of ninebanded armadillo (*Dasypus novemcinctus texanus*), house mice (*Mus musculus*), opossum (*Didelphis virginiana*), and wood rats (*Neotoma micropus micropus*), with *Trypanosoma cruzi* in the state of Texas. Am J Trop Med Hyg 22 (Suppl 1):623–631.
- Pesavento PA, Chomel BB, Kasten RW, McDonald KA, Mohr FC. 2005. Pathology of *Bartonella* endocarditis in six dogs. *Vet Pathol* 42:370–373.
- Pietrzak SM, Pung OJ. 1998. Trypanosomiasis in raccoons from Georgia. J Wildl Dis 34:132–136.
- Piron M, Fisa R, Casamitjana N, López-Chejade P, Puig L, Vergés M, Gascón J, Gómez i Prat J, Portús M, Sauleda S. 2007. Development of a real-time PCR assay for *Trypanosoma cruzi* detection in blood samples. *Acta Trop* 103:195–200.
- Pisharath H, Zao C-L, Kreeger J, Portugal S, Kawabe T, Burton T, Tomaeck L, Shoieb A, Campbell BM, Franco J. 2013. Immunopathologic characterization of naturally acquired *Trypanosoma cruzi* infection and cardiac sequalae in cynomolgus macaques (*Macaca fascicularis*). J Am Assoc Lab Anim Sci 52:545–552.
- Pronovost H, Peterson AC, Chavez BG, Blum MJ, Dumonteil E, Herrera CP. 2019. Deep sequencing reveals multiclonality and new discrete typing units of *Trypanosoma cruzi* in rodents from the southern United States. J Microbiol Immunol Infect. In press.
- Ramírez JD, Guhl F, Rendón LM, Rosas F, Marin-Neto JA, Morillo CA. 2010. Chagas cardiomyopathy manifestations and *Trypanosoma cruzi* genotypes circulating in chronic Chagasic patients. *PLoS Negl Trop Dis* 4:e899.
- Rassi A Jr, Rassi A, Marin-Neto JA. 2010. Chagas disease. Lancet 375:1388–1402.
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project. org. Accessed June 2019.

- Rocha FL, Roque A, de Lima JS, Cheida CC, Lemos FG, de Azevedo FC, Arrais RC, Bilac D, Herrera HM, Mourão G, et al. 2013. *Trypanosoma cruzi* infection in neotropical wild carnivores (Mammalia: Carnivora): At the top of the *T. cruzi* transmission chain. *PLoS One* 8:e67463.
- Roellig DM, Brown EL, Barnabé C, Tibayrenc M, Steurer FJ, Yabsley MJ. 2008. Molecular typing of *Trypanosoma cruzi* isolates, United States. *Emerg Infect Dis* 14:1123–1125.
- Roellig DM, Ellis AE, Yabsley MJ. 2009a. Oral transmission of *Trypanosoma cruzi* with opposing evidence for the theory of carnivory. *J Parasitol* 95:360–364.
- Roellig DM, Ellis AE, Yabsley MJ. 2009b. Genetically different isolates of *Trypanosoma cruzi* elicit different infection dynamics in raccoons (*Procyon lotor*) and Virginia opossums (*Didelphis virginiana*). Int J Parasitol 39:1603–1610.
- Roellig DM, Savage MY, Fujita AW, Barnabé C, Tibayrenc M, Steurer FJ, Yabsley MJ. 2013. Genetic variation and exchange in *Trypanosoma cruzi* isolates from the United States. *PLoS One* 8:e56198.
- Ryan CP, Hughes PE, Howard EB. 1985. American trypanosomiasis (Chagas' disease) in a striped skunk. J Wildl Dis 21:175–176.
- Snowden KF, Kjos SA. 2013. American trypanosomiasis. In: *Infectious diseases of the dog and cat*, 4th Ed., Sykes JE, Greene CE, editors. Elsevier Inc., St. Louis, Missouri, pp. 722–730.
- Thomas ME, Rasweiler JJ IV, D'Alessandro A. 2007. Experimental transmission of the parasitic flagellates *Trypanosoma cruzi* and *Trypanosoma rangeli* between triatomine bugs or mice and captive neotropical bats. *Mem Inst Oswaldo Cruz* 102:559–565.
- Zingales B, Miles MA, Campbell DA, Tibayrenc M, Macedo AM, Teixeira MMG, Schijman AG, Llewellyn MS, Lages-Silva E, Machado CR, et al. 2012. The revised *Trypanosoma cruzi* subspecific nomenclature: Rationale, epidemiological relevance and research applications. *Infect Genet Evol* 12:240–253.

Submitted for publication 22 March 2019. Accepted 11 June 2019.